



Anti-Inflammatory Effect of Three Novel Hydroxy Flavones

K. Vijay Kumar^{a*‡} and Reena Verma^{b#}

^a Department of Pharmacology, RKDF Medical College, SRK University, Bhopal, India.

^b Department of Pharmacology, RKDF Medical College and Hospital, Bhopal, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i61B35511

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/83473>

Original Research Article

Received 14 October 2021

Accepted 25 December 2021

Published 28 December 2021

ABSTRACT

Aim: The aim of the present study was to evaluate the anti-inflammatory effect of three 7-hydroxy flavone derivatives; 7-HF, 7,3',4'-THF, and 7,8,3'-THF,

Materials and Methods: The acute anti-inflammatory effect of 7-hydroxy flavones was the effect of 7-hydroxy flavones on certain mediators of pain and inflammation like cyclooxygenases (COX-1 and COX-2), and pro-inflammatory cytokines (IL-1b and TNF- were investigated by using in vitro tests.

Results: The investigated 7hydroxy flavones produced a significant, the test compounds inhibited both the isoforms of cyclooxygenase, a higher degree of inhibition on COX-2 was evident. Concentration-dependent inhibition of other inflammatory cytokines like tumour necrosis factor- α and interleukin-1b was identified in the present study

Conclusion: conclusion of the present study reveal the pain and anti-inflammatory action of the investigated 7hydroxy flavones. The inhibitory effects of 7hydroxy flavone on various pro-inflammatory cytokines provide evidence for the mechanism of action of these compounds against pain and inflammation.

Keywords: Anti-inflammatory effect; cyclooxygenases; Interleukin-1b; flavone.

[‡]Research Scholar;

[#]Professor & HOD;

*Corresponding author: E-mail: vijay_kalwa2006@yahoo.com;

1. INTRODUCTION

Non-steroidal anti-inflammatory medicines (NSAIDs) are a class of pharmaceuticals that have anti-inflammatory, analgesic, and antipyretic properties. Glucocorticoids are also the cornerstone of treatment for a number of chronic inflammatory and autoimmune illnesses. However, many people are unable to use them because of their intrinsic side effects. As a result, anti-inflammatory medications that are both safe and effective are being aggressively sought all over the world. Flavonoids appear to be promising as safe and effective anti-inflammatory medications among numerous types of experimental agents. Despite the fact that flavonoids have a number of important pharmacological properties, their therapeutic utility is limited due to their low bioavailability. However, Flavones have a higher bioavailability than other derivatives, according to a recent study [1], implying that flavonoids could be used to treat a wide spectrum of illnesses. Flavonoids are polyphenolic chemicals found in a wide range of angiosperms that have a wide range of health benefits and pharmacological activities [2]. Many natural and synthetic flavone derivatives have been studied for their analgesic and anti-inflammatory properties in recent decades. In acute and chronic animal models of inflammation, flavonoids such as hesperidin, apigenin, luteolin, and gossypin [3], titionine [4] and its derivatives, silymarin [5], and naringin [6]. The anti-inflammatory effect of flavonoids was further validated by studies on a few monomethoxy flavones [7], monohydroxy flavones [8], and dihydroxy flavones [9,10]. A few structurally similar dimethoxy flavones were examined for their pharmacological activities in a recent study [11]. These dimethoxy flavones have been found to have potent antinociceptive action, especially in inflammatory pain models. As a result of this discovery, the anti-inflammatory potential of dimethoxy flavones was investigated, and the results were evaluated in this study. The effect of these compounds on important mediators of inflammation such as interleukin-1b (IL-1b), tumour necrosis factor-alpha (TNF- α), and the generation of prostaglandins were also investigated using appropriate in vitro techniques.

2. MATERIALS AND METHODS

Carrageenan (Sigma Chemical Co., USA) was used as a phlogistic agent and diclofenac sodium (Novartis) was used as a standard anti-

inflammatory agent. Diagnostic kits (Cayman., USA) were used for the in vitro assay of cyclooxygenases, IL-1b and TNF- α .

2.1 Effect of Dimethoxy Flavones on Cyclooxygenase (COX) Enzyme by Immunoassay

Cyclooxygenase assay was carried out using COX (ovine) inhibitor screening assay (Cayman, USA). This assay directly measures prostaglandin-F₂ (PGF_{2a}) which is derived from stannous chloride reduction of PGH₂ produced in the cyclooxygenase reaction. 50 μ of PGH₂ and various dimethoxy flavones dissolved and diluted in dimethyl sulfoxide (DMSO) to 5–50 IM concentration were incubated with reaction buffer and arachidonic acid. The reaction was allowed to proceed for 2 min and was quenched by addition of 50 μ of stannous chloride solution. The reaction was allowed to proceed for an additional 10 min. Further serial dilutions were made with the buffer and PGF_{2a} was measured using enzyme immuno assay (EIA). The amount of PGF_{2a} was directly proportional to the amount of COX-1 and COX-2 present in the well. Ibuprofen and celecoxib were used as standard COX-1 and COX-2 inhibitors respectively.

Effect of dimethoxy flavones on cytokines— Interleukin-1b (IL-1b) and Tumor necrosis factor-a (TNF-a).

Freshly heparinised human whole blood was used for the immunometric assay, which is based on a double antibody “sandwich” technique. Microwell plates supplied with the commercial kit (Cayman, USA) have been coated with monoclonal antibody specific for TNF-a or IL-1b that can capture any TNF- α or IL-1b introduced into the well. Various dimethoxy flavones dissolved and diluted in DMSO to different concentrations (20–500 IM) were added to the well. Acetylcholinesterase (AChE) 50 μ that binds selectively to a different epitope on the TNF-a or IL-1b molecule, was also added to the well. When TNF- α or IL-1b (standard or sample) is introduced, the two anti- bodies form a sandwich by binding on opposite sides of the TNF- α or IL-1b molecule. The concentration of the analyte was then determined by measuring the enzymatic activity of the acetylcholinesterase by adding Ellman’s reagent (which contains the substrate for AChE) to each well. The yellow color of the product of the AChE-catalyzed reaction was

read spectrophotometrically at 412 nm. The intensity of this color was directly proportional to the amount of bound conjugate which in turn was proportional to the concentration of the TNF- α or IL-1b. Dexamethasone was used as a standard for both TNF- α and IL-1b assays.

3. RESULTS

3.1 Effect of Trihydroxy Flavones on Cyclooxygenase – 1 (Table 1)

The investigated hydroxy flavones inhibited cyclooxygenase-1 activity in a concentration-dependent manner. Among the tested compounds, 7-hydroxy flavone produced a maximum inhibition of 86.23% in a 50 -M concentration. In a similar concentration 7,3'4'-THF produced inhibition of 78%, While 7,8,3', THF produced an inhibition of 67.24% (Table 1). The IC₅₀ values recorded for these compounds in increasing order are as follows: 7,3',4' – THF = 28.04 μ M, 7,8,3' – THF = 35.81 μ M, 7 – HF = 22.72 μ M. The IC₅₀ value of

ibuprofen in this assay procedure was found to be 14.68 μ M.

3.2 Effect of Trihydroxy Flavones on Cyclooxygenase 2 (Table 2)

Concentration-dependent inhibition of cyclooxygenase – 2 was clearly evident for all the tested trihydroxy flavones. A maximum of 71% inhibition was recorded for 7,3',4'-THF in a concentration of 50 μ M. In the same concentration maximum inhibition of 59.18.89%, and 67.9% was recorded for 7 – HF, 7,8,3'- THF respectively (Table 2). The IC₅₀ values recorded for the above compounds to inhibit cyclooxygenase – 2 are as follows: 7,8,3'–THF = 32.32 μ M, 7,3',4' – THF = 30.91 μ M , 7 – HF = 38.92 μ M. The IC₅₀ value for celecoxib in the same assay procedure was found to be 2.24 μ M.

It is pertinent to point out that all the investigated 7 hydroxy flavones almost equally inhibited both cyclooxygenase - 1 and cyclooxygenase – 2. The IC₅₀ values for inhibition of cyclooxygenase 1 and 2 for these compounds are also almost identical.

Table 1. Effect of trihydroxy flavones on cyclooxygenase – 1

Concentration (-M)	Treatment mg/Kg, sc and % Inhibition		
	7-HF	7, 3',4'- THF	7,8,3' – THF
10	27.5	24.3	19.12
20	45.97	36.23	32.55
30	69.3	54.68	40.55
40	73.87	70.78	55.79
50	86.23	77.89	67.24
IC ₅₀ (-M)	22.72565	28.04135	35.81687

Each value represents the mean of 3 observations

* IC₅₀ value calculated by linear regression analysis




Fig. 1. Effect of trihydroxy flavones on cyclooxygenase – 1

3.3 Effect of 7 Hydroxy Flavones on Interleukin – 1 β (Table 3)

Concentration-dependent inhibition of interleukin – 1 β activity was recorded for all the tested trihydroxy flavones. Nearly 80.56% inhibition was recorded for 7- HF in a concentration of 50 μ M. 7,8,3'- THF exhibited a maximum inhibition of 78.86% in the same concentration while that recorded for 7,3',4'- THF was 74%. The IC₅₀ values recorded for the above hydroxy flavones are as follows 7, 8, 3' – THF = 122.56 μ M, 7, 3', 4' – THF = 128.89 μ M, and 7 - HF = 138.01 μ M.

The IC₅₀ value of dexamethasone to inhibit interleukin 1 β in the same experimental procedure was found to be 101.23 μ M. It can be appreciated that the IC₅₀ value of all 7 HF is almost similar to that of dexamethasone,

whereas the compounds possessed a slightly higher IC₅₀ value (Table 3).

The investigated trihydroxy flavones exerted a potent inhibition of the activity of the tumour necrosis factor α . The inhibition was concentration-dependent (Table 4).

A fairly high degree of inhibition was recorded for 7 hydroxy flavones in concentrations ranging from 10 – 50 μ M. A maximum of 72.16% inhibition was recorded with 7-HF. In a similar concentration, 7,8,3'- THF produced a maximum inhibition of 64.32%. The inhibitory action of 7,3'4'- THF on tumour necrosis factor α was revealed only in higher concentrations. A maximum of 63.54% inhibition was observed. The IC₅₀ values calculated for the above compounds are as follows: 7– HF = 28.76 μ M, 7,8,3' – THF = 35.13 μ M and 7,3',4' – THF = 38.18 μ M.

Table 2. Effect of trihydroxy flavones on Cyclooxygenase –2

Concentration (μ M)	Treatment mg/Kg, sc and % Inhibition		
	7-HF	7, 3',4' – THF	7,8,3'– THF
10	17.48	24.52	25.36
20	30.97	39.91	39.54
30	45.97	49.76	46.43
40	50.48	59.48	58.68
50	59.18	71.18	67.9
IC ₅₀ (μ M)	38.9243	30.91239	32.32009

Each value represents the mean of 3 observations
 * IC₅₀ value calculated by linear regression analysis




Fig. 2. Effect of trihydroxy flavones on cyclooxygenase –2

Table 3. Effect of trihydroxy flavones on Interleukin – 1 β

Concentration (-M)	Treatment mg/Kg, sc and % Inhibition		
	7-HF	7,3',4'-THF	7, 8, 3'- THF
10	26.23	21.24	20.46
20	35.89	29.82	29.21
30	40.12	38.76	39.09
40	64.98	59.45	64.98
50	80.56	73.96	78.86
IC₅₀ (-M)	138.01	128.89	122.56

Each value represents the mean of 3 observations
 * IC₅₀ value calculated by linear regression analysis




Fig. 3. Effect of trihydroxy flavones on interleukin – 1 β

Table 4. Effect of trihydroxy flavones on tumor necrosis factor – α

Concentration -M	% Inhibition		
	7-HF	7,3',4' THF	7,8,3'THF
10	23.13	12.97	21.46
20	38.16	22.68	30.23
30	57.15	38.77	46.66
40	67.28	50.01	58.14
50	72.16	63.54	64.32
IC-50*	28.76 (μM)	38.18μM)	35.13(μM)

Each value represents the mean of three observations
 *IC – 50 value calculated by linear regression analysis

The IC₅₀ value for dexamethasone to inhibit tumor necrosis factor- α was recorded to be 25.33 μ M. It can be appreciated that the IC₅₀ values of 7- HF (27.08 μ M) and 7,8,3'- THF (34.54 μ M) are similar to dexamethasone value. The IC₅₀ values recorded for 7,3',4'- THF (38.13 μ M) are many-fold higher than the dexamethasone value.

4. DISCUSSION

The anti-inflammatory activity of flavonoids as reported by [10] revealed 50% inhibition of TNF- α with vitamin C, epicatechin, epigallocatechin, procyanidin B2, quercetin, and taxifolin. The same study has reported the inhibition of IL-1




Fig. 4. Effect of trihydroxy flavones on tumor necrosis factor – α

beta by epicatechin, epigallocatechin, procyanidin B2, quercetin was comparable to vitamin C [10]. In our study A concentration-dependent inhibition of interleukin- 1β activity was recorded for all the tested trihydroxy flavones. Nearly 78% inhibition was recorded for 7,8,3'- THF in a concentration of 50 μ M while that recorded for 7- HF was 80%. The IC₅₀ values recorded for hydroxy flavones were 122.56 μ M for 7, 8, 3'-THF, 128.89 μ M for 7, 3', 4'-THF, and 138.01 μ M for 7-HF. Our study also showed high degree of inhibition of TNF-alpha for 7-HF in concentrations ranging from 10 – 50 μ M. A maximum of 72.16% inhibition was recorded. In a similar concentration, 7,8,3'- THF produced a maximum inhibition of 64.32%. The inhibitory action of 7,3'4'- THF on tumor necrosis factor-alpha was revealed only in higher concentrations. A maximum of 72.16% inhibition was observed. The IC₅₀ values recorded for hydroxy flavones were 27.08 μ M for 7-HF, 38.13 μ M for 7,8,3'-THF and 34.54 μ M for 7,3',4'-THF. The COX forms prostaglandins and thromboxanes which causes inflammation [12,13]. Many flavonoids exhibited inhibition of COX-1 and COX-2 which were reported in previous research studies [14]. also reported the COX-1 and COX-2 inhibitory property of flavonoids [15]. In our study, flavones inhibited COX-1 activity in a concentration dependent manner. Among the tested compounds, 7-hydroxy flavone produced a maximum inhibition of 86.23% in 50 μ M concentration. In a similar concentration 7,3'4'- THF produced an inhibition

of 76%, While 7- HF produced an inhibition of 58.8%. The IC₅₀ values of COX-1 inhibition for hydroxy flavones were 27.61 μ M for 7,3',4'-THF, 28.21 μ M for 7,8,3'-THF, and 39.40 μ M for 7-HF. In contrary, revealed that flavones could not inhibit COX-1 [16]. The COX-2 inhibition was maximum (71%) for 7,3',4'-THF in a concentration of 50 μ M. In the same concentration maximum inhibition of 59.89%, and 67.9% with 7-HF, and 7,8,3'-THF respectively. The IC₅₀ values of COX-2 inhibition for hydroxy flavones were 31.98 μ M for 7,3',4'-THF, 28.46 μ M for 7,8,3'-THF, and 40.31 μ M for 7-HF.

5. CONCLUSION

In conclusion, the present study has identified three 7hydroxy flavone derivatives with good anti-inflammatory activity. The inhibitory effects of 7hydroxy flavone on various pro-inflammatory cytokines provide evidence for the mechanism of action of these compounds against pain and inflammation.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Li S, Min-Hsiung P, Chih-Yu L, Do T, Yu W, Fereidoon S, Chi-Tang H. Chemistry and health effects of polymethoxy flavones and hydroxylated polymethoxy flavones. *J Func Foods*. 2009;1:2–12.
2. Alcaraz MJ, Ferrandiz ML. Modification of arachidonic acid metabolism by flavonoids. *J Ethnopharmacol*. 1987;21:209–229.
3. Parmar NS, Ghosh MN. Antiinflammatory activity of gossypin, a bioflavonoid isolated from *Hibiscus vitifolius* Linn. *Ind J Pharmacol*. 1978;10:277–293.
4. Carvalho JC, Ferreira LP, Santos DSL, Correa MJ, Campos DOLM, Bastos JK, Sarti SJ. Antiinflammatory activity of flavone and some of its derivatives from *Virola Michelli* heckle (Myristicaceae). *J Ethnopharmacol*. 1999;64:173.
5. Gupta OP, Singh S, Bani S, Sharma N, Malhotra S, Gupta BD, Banergee SK, Handa SS. Antiinflammatory and anti-arthritic activities of silymarin acting through inhibition of 5-lipoxygenase. *Phytomed*. 2000;1:21–24.
6. Havsteen BH. The Biochemistry and medical significance of the flavonoids. *Pharmacol Ther*. 2002;96:67–202.
7. Muthiah NS, Viswanathan S, Thirugnana sambantham P, Reddy MK, Vijayasekaran V. Antiinflammatory activity of flavone and its mono-methoxy derivatives. A structure activity study. *Ind J Pharm Sci*. 1993;55:180–183.
8. Arivudainambi R, Viswanathan S, Thirugnana sambantham P, Reddy MK, Dewan ML, Sijheer JS, Gopalakrishnan C, Vijayasekaran V. Anti-inflammatory activity of flavone and its hydroxy derivatives. A structure activity study. *Ind J Pharm Sci*. 1996;58:18–21–177.
9. Umamaheswari S. Characterization of antinociceptive and anti-inflammatory effects of certain dihydroxy flavones. Ph.D. Thesis, Sri Ramachandra University, Chennai; 2008.
10. Vidyalakshmi K, Kamalakannan P, Viswanathan S, Ramasamy S. Anti-inflammatory effect of certain dihydroxy flavones and the mechanism involved. *Anti-inflammatory Anti-allergy agents Med Chem*. 2012;11:253–261.
11. Kamalakannan P, Vidyalakshmi K, Viswanathan S, Ramasamy S. Antinociceptive effect of certain dimethoxy flavones in mice. *Eur J Pharmacol*. 2014;727:148–157.
12. Kujubu DA, Fletcher BS, Varnum BC, Lim RW, Herschman HR. TIS10, a phorbol ester tumor promoter-inducible mRNA from Swiss 3T3 cells, encodes a novel prostaglandin synthase/cyclooxygenase homologue. *J Biol Chem*. 1991 Jul 15;266(20):12866-72.
13. Smith WL, DeWitt DL, Garavito RM. Cyclooxygenases: structural, cellular, and molecular biology. *Annu Rev Biochem*. 2000;69:145-82.
14. Madeswaran A, Umamaheswari M, Asokkumar K, Sivashanmugam T, Subhadradevi V, Jagannath P. Docking studies: In silico lipoxygenase inhibitory activity of some commercially available flavonoids. *||| Bangladesh Journal of Pharmacology|||*. 2011;6(2):133-8.
15. Ribeiro D, Freitas M, Tomé SM, Silva AM, Laufer S, Lima JL, Fernandes E. Flavonoids inhibit COX-1 and COX-2 enzymes and cytokine/chemokine production in human whole blood. *Inflammation*. 2015 Apr;38(2):858-70.
16. Valdés-Barrera ID, Cuca-Suarez LE, Coy-Barrera ED. Nectandra amazonum-derived flavonoids as COX-1 inhibitors: *In vitro* and docking studies. *Natural product communications*. 2014 May;9(5): 1934578X1400900515.

© 2021 Kumar and Verma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/83473>