



Study of Correlation and Path Coefficient Analysis for Different Morphological Characters in Forage Sorghum (*Sorghum bicolor* L. Moench)

**Nikhil ^a, S. K. Singh ^a, L. K. Gangwar ^a,
Alok Kumar Mourya ^{b*}, Pushendra ^a and Chiranjeev ^a**

^a Department of Genetics and Plant Breeding, Sardar Vallabh Bhai Patel University of Agriculture and Technology Meerut, (250110), India.

^b Department of Genetics and Plant Breeding, Chandra Shekhar Azad University of Agriculture and Technology Kanpur, (208002), India.

Authors' contributions

This work was carried out in collaboration among all authors. Author Nikhil conducted the research, collected and analyzed the data. Authors SKS and LKG supervised the study. Authors AKM prepared the manuscript. Author Pushendra reviewed the manuscript. All authors read and approved the final manuscript.

Article Information

DOI:10.9734/IJECC/2023/v13i113576

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108724>

Original Research Article

Received: 14/09/2023

Accepted: 19/11/2023

Published: 29/11/2023

ABSTRACT

This research was conducted to investigate the correlation and path coefficient analysis of thirty three genotypes for green fodder yield and its attributes of forage sorghum. The experimental material was planted in Completely Randomized Block Design (RBD) in three replication during *Kharif* in the year 2019-20 at Crop Research Centre, SVPUA&T, Modipuram, Meerut - 250110 (U.P.). The study of association and path analysis evaluation carried out using the ten green fodder yield traits, viz., days to 50% flowering, plant height, leaf length, leaf breadth, leaf area, stem girth, leaves per plant, leaf stem ratio, total soluble solids and green fodder yield per plant. There is significant and positive correlation with green fodder yield per plant was exhibited with stem girth,

*Corresponding author: E-mail: alokkumarmourya@gmail.com;
E-mail: nikhilahlawat136@gmail.com;

leaves per plant, leaf length, leaf breadth, leaf stem ratio and total soluble solids at genotypic and phenotypic level. While leaf area showed the maximal direct effect on green fodder yield per plant followed by stem girth, leaves per plant and leaf stem ratio. There for selection for these traits is advisable for improving the character of green fodder yield.

Keywords: Genetic variability; correlation; path coefficient; quantitative traits.

1. INTRODUCTION

“Sorghum (*Sorghum bicolor* L. Moench) is first important fodder crop, forasmuch of this has highly adaptation for the different ecological conditions, it has the wide range of resistant to drought and also suitable for low inputs requirement for cultivation [1]. “Sorghum is a C4 plant has the higher photosynthesis efficiency with more abiotic stress tolerance” [2,3]. “Mostly five races viz., *Bicolor*, *Guinea*, *Caudatum*, *Durra* and *Kafirare* found as cultivated sorghum and ten races are intermediate which corresponding to the pair wise combination of major races. All ten races are recognize by the phenotypic characteristics, specifically inflorescence related traits” [4]. “Sorghum is the fifth most important food crop globally after Rice, Wheat, Maize and Pearlmillet and is animal dietary staple feed of nearly 500 million people in 30 countries. This is growing area in India is nearly 5.14 million hectares while its annual production is 4.57 million tonnes and productivity is 889kg ha⁻¹. In India its major producer states are Maharashtra, Karnataka, Madhay Pradesh, Andhra Pradesh, Rajasthan and Gujarat” [5].

“The sorghum is majorly used for the purpose of food, feed, and forage crop. Alongside of sorghum also bestow the raw materials for the production of starch, fiber, dextrose syrup bio fuels, vinegar, alcohol and more products”. Karadi and Kajjidoni [6].

To develop sorghum as a viable and profitable crop, there is an incontinently need to begin research to develop varieties and hybrids with high grain and fodder yields, fast growth, early to medium maturity and good quality parameters. Improving grain and fodder yield potential in sorghum is a major concern for breeders. It has been observed that cultivated sorghum is more variable and suggests that, to increase yield, it is first necessary to know the nature and extent of genes present in the germplasm collection is a backbone. Estimating of genetic variability in the base population is the first step in any breeding program. It is necessary to study the relationships between traits to evaluate the

feasibility of joint selection for two or more traits. The path coefficient is just a standard partial regression coefficient that measures the direct effect of one variable on another variable and allows the correlation coefficient to be separated into components of direct and indirect effects of different characteristics on the dependent variable, thus making effective selection. To understand the effect of green fodder yield, significant genotypic correlations of various attributes with green fodder yield per plant were divided into their direct and indirect effects. This will provide more detailed information to select key conditions that can contribute more to green fodder yield. The genetic variability is the important phenomena for ecological balance and this study aims to assessment the present variation in outcomes between selected genotypes.

2. MATERIALS AND METHODS

This experimental material consisting of 33 advanced lines (Table 1) was grown in three replicates in a Randomized Blocked Design (RBD). The experimental material was planted in fields under irrigated conditions at the Crop Research Centre, Modipuram, SVPUA&T, Meerut - 250110 (U.P.) during the *Kharif* season 2019-20. All recommended agronomic practices were followed to produce a good sorghum crop with 4 rows of 5 meters long. The rowing space was 30 cm and the planting space was to be planted at 10 cm respectively. Observations were recorded on yield and yield attributing characters viz., days to 50% flowering (day), plant height (cm), leaf length (cm), leaf breadth (cm), leaf area (cm)², stem girth (mm), leaves per plant, leaf stem ratio, total soluble solids (% by refractometer reading), green fodder yield (g/plant). Flowering day was recorded as the number of days after sowing, 50 percent of plants in each plot flowered while leaf area (cm)² = leaf length (cm) x leaf breadth x 0.71 and leaf stem ratio calculated using the formula, Leaf stem ratio = Fresh weight of all the leaves/Fresh weight of stems. All the observations were taken from each plot, on randomly five selected plants from each genotype. The correlations at

Table 1. List of forage sorghum genotypes

Sl. No.	Genotype	Sl. No.	Genotype	Sl. No.	Genotype
1.	HC-541	12.	Pant Chari-3	23.	CSV-14
2.	Pant Chari-8	13.	Pant Chari-5	24.	CSV-13
3.	Pusa Chari-6	14.	HC-573	25.	Bundela Chari
4.	HC-171	15.	HC-308	26.	HJ-513
5.	HC-136	16.	Pusa Chari-9	27.	CSV-12
6.	Pratap Raj Jowar-1	17.	Pant Chari-4	28.	HC-260
7.	MP Chari	18.	SSG59-3	29.	CSV-28
8.	HJ-541	19.	Pusa Chari-23	30.	CSV-20
9.	Pusa Chari-615	20.	Versha	31.	PJ-1430
10.	Pant Chari-6	21.	CSV-15	32.	CSV-27
11.	Pant Chari-2	22.	Pant Chari-7	33.	CSV-23

genotypic, phenotypic and environmental levels were calculated from the analysis of variance and covariance as suggested by Searle [7] while path coefficient analysis was performed according to the method suggested by Dewey and Lu [8]

3. RESULTS AND DISCUSSION

Correlation coefficients are estimated between the yield and its contributed characteristics at the genotypic and phenotypic levels. Correlations at the genotypic level are higher than their corresponding phenotypic correlations. The results of genotypic and phenotypic correlation coefficients between the crop and its components are given in (Table 2). In most cases the direction and magnitude of genotypic and phenotypic correlations between different traits remained nearly identical.

In this study, green fodder yield is estimated as significant and positive correlation was exhibited with stem girth (0.68), leaves per plant (0.67), leaf length (0.64), leaf breadth (0.47), leaf stem ratio (0.27) and total soluble solids (0.20) at genotypic level grain yield per plant was found to be significantly and positively correlated with green fodder yield per plant was exhibited with stem girth (0.68), leaves per plant (0.67), leaf length (0.64), leaf breadth (0.47), leaf stem ratio (0.27) and total soluble solids (0.20) at genotypic level. While green fodder yield per plant showed negative significant relationship with days to 50% flowering (-0.25), green fodder yield per plant with plant height (-0.12) and leaf area (-0.19) A negative however no significant correlation is shown at the genotypic level. Hence, enhancement of green fodder yield can be achieved by improving these characters in forage sorghum. This result are ingenerally revealed with the finding of Dubey et al [9] recorded the leaf area, leaf length, number of leaves, grain weight showed significant positive correlation

with yield per plant. Jain et al [10] recorded plant height, stem girth and leaf length were positively and significantly associated with green fodder and dry fodder yield per plant. Prasad and Sridhar [12]. exhibited positive significant association green fodder yield with days to 50% flowering, days to maturity, number of leaves per plant, leaf length, leaf breadth, stem ratio and yield per plant. While Rohila et al. [11] recorded negatively associated the time of panicle emergence with green fodder yield.

The division of path coefficients into direct and indirect effects was done based on the results at the phenotypic level (Table 3). Critical observation of the results in the table revealed that leaf area (0.64) showed the maximum order of direct effect on green fodder yield, followed by stem girth (0.51), leaves per plant (0.49) and leaf stem ratio (0.17) was observed. The identical result reported by Damor *et al.* (2018) where a positive direct effect on grain yield for green fodder yield at both genotypic and phenotypic levels and Prasad and Sridhar [12] recorded positive direct effect on grain yield for fodder yield at both genotypic and phenotypic level and Prasad and Sridhar [12] found the plant height exhibited positive direct effect with yield per plant. While maximum negative direct effect was shown for plant height (-0.14), leaf length (-0.15) and leaf breadth (-0.19). Direct and indirect effects also recorded at the phenotypic level were generally similar to those shown at the genotypic level with little variation in Magnitude. Levels of residual effects were low at the phenotypic and genotypic levels, indicating that the characters included in the present Investigation were sufficient enough to account for the variability in the dependent attribute *i.e.* green fodder yield. The direct contribution of leaf area, stem girth, leaves per plant and leaf stem ratio with green fodder yield per plant observed in this study is also in confirmation with the findings Malaghan and Kajjidoni [13].

Table 2. Estimates of correlation coefficients for genotypic (G) and phenotypic (P) levels among different characters in forage sorghum (*Sorghum bicolor* L. Moench)

Characters		Days to 50% flowering	Plant Height (cm)	Leaf length (cm)	Leaf breadth (cm)	Leaf area (cm ²)	Stem girth (mm)	Leaves per plant	Leaf stem ratio	Total soluble solids (%)	Green fodder yield per plant(g)
Days to 50% flowering	G	1.00	0.12	0.11	0.30**	0.25**	0.25**	-0.02	-0.02	0.36**	-0.25**
	P	1.00	0.15	0.12	0.26**	0.22**	0.22**	-0.02	-0.02	0.31**	-0.23**
Plant Height (cm)	G		1.00	0.23**	0.33**	0.41**	-0.11	0.21*	0.20*	-0.33**	-0.12
	P		1.00	0.22**	0.39**	0.42**	-0.11	0.20*	0.20*	-0.33**	-0.12
Leaf length (cm)	G			1.00	-0.09	0.40**	0.23*	0.14	0.33**	0.30**	0.64**
	P			1.00	-0.09	0.39**	0.23*	0.13	0.32**	0.30**	0.65**
Leaf breadth (cm)	G				1.00	0.94**	0.23*	0.31**	-0.03	-0.10	0.47**
	P				1.00	0.94**	0.23*	0.29**	-0.03	-0.10	0.49**
Leaf area (cm ²)	G					1.00	0.14	0.34**	-0.01	-0.13	-0.19
	P					1.00	0.18	0.39**	-0.02	-0.13	-0.19
Stem girth (mm)	G						1.00	-0.06	0.51**	0.12	0.68**
	P						1.00	-0.06	0.50**	0.12	0.69**
Leaves per plant	G							1.00	0.21	-0.10	0.67**
	P							1.00	0.16	-0.10	0.69**
Leaf stem ratio	G								1.00	0.45**	0.27**
	P								1.00	0.44**	0.27**
Total soluble solids (%)	G									1.00	0.20*
	P									1.00	0.20*
Green fodder yield per plant (g)	G										1.00
	P										1.00

*** significant at 5% and 1% level respectively

Table 3. Path coefficient analysis showing the direct and indirect effect of ten characters on the green fodder yield at genotypic and phenotypic levels of forage sorghum (*Sorghum bicolor* L. Moench)

Characters		Days to 50% flowering	Plant Height (cm)	Leaf length (cm)	Leaf breadth (cm)	Leaf area (cm ²)	Stem girth (mm)	Leaves per plant	Leaf stem ratio	Total soluble solids (%)
Days to 50% flowering	G	0.10	-0.02	-0.01	-0.04	0.05	-0.13	-0.21	-0.06	0.01
	P	0.03	-0.01	-0.01	-0.05	0.05	-0.11	-0.14	-0.02	0.02
Plant Height (cm)	G	0.01	-0.15	-0.03	-0.05	0.08	0.06	-0.07	0.04	0.01
	P	0.04	-0.14	-0.03	-0.06	0.10	0.06	-0.05	0.03	0.03
Leaf length (cm)	G	0.01	0.23	-0.13	-0.02	0.08	0.04	-0.04	0.07	-0.02
	P	0.05	0.33	-0.15	-0.02	0.09	0.04	-0.03	0.08	-0.02
Leaf breadth (cm)	G	0.03	0.25	-0.02	-0.15	0.19	-0.12	-0.10	-0.08	0.02
	P	0.09	0.24	-0.02	-0.19	0.23	-0.12	-0.08	-0.09	-0.02
Leaf area (cm ²)	G	0.02	-0.06	-0.05	-0.14	0.62	-0.07	-0.09	0.07	-0.24
	P	0.07	-0.05	-0.06	-0.18	0.64	-0.07	-0.07	0.06	-0.24
Stem girth (mm)	G	0.02	0.21	0.01	-0.03	0.02	0.50	-0.11	-0.06	-0.04
	P	0.04	0.24	0.01	-0.04	0.03	0.51	-0.08	-0.05	-0.09
Leaves per plant	G	0.06	0.04	-0.02	-0.04	0.05	-0.17	0.44	0.02	0.03
	P	0.07	0.06	-0.02	-0.05	0.06	-0.16	0.49	0.01	0.07
Leaf stem ratio	G	-0.03	0.34	-0.04	0.06	0.06	0.15	-0.03	0.21	0.01
	P	-0.01	0.36	-0.05	0.07	0.08	0.15	-0.02	0.17	0.03
Total soluble solids (%)	G	0.03	0.02	-0.04	0.02	-0.02	0.05	-0.02	-0.02	-0.03
	P	0.05	0.02	-0.04	0.02	-0.03	0.05	-0.02	-0.01	-0.08

Residual values (G) = 0.17 residual values (P) = 0.28*, ** significant at 5%, 1% level, respectively Bold values indicate direct effects

4. CONCLUSION

Green fodder yield per plant had highly significant positive correlations with leaf length, leaf breadth, stem girth, leaves per plant, leaf stem ratio and total soluble solids at genotypic and phenotypic levels. Therefore, green forage

productivity can be increased by improving these characteristics in forage sorghum.

The genotypic and phenotypic path coefficient analysis of green fodder yield per plant and its component attributes showed that leaf area had the extremely positive direct on green fodder

yield per plant followed by stem girth, leaves per plant and leaf stem ratio, indicates that these traits could be considered as most important characters for improving fodder yield.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Doggett H. Sorghum, second edition. New York, USA: John Wiley and Sons. 1988;512.
2. Nagy ZZ, Tuba F, Soldos Z, Erdei L. CO₂ exchange and water relation responses of sorghum and maize during water and salt stress. *Journal of Plant Physiology*. 1995;145: 539-544.
3. Reddy BVS, Ramesh S, Reddy PS, Kumar AA. Genetic enhancement for drought tolerance in sorghum. *Plant Breeding Reviews*. 2009;31:189-222.
4. Harlan JR, de Wet JM. A simple classification of cultivated sorghum. *Crop Science*. 1972;12: 172-176.
5. Anonymous. Agriculture statistics at a glance, ministry of agriculture, Department of agriculture and cooperation, directorate of economics and statistics, New Delhi, India. 2018;99.
6. Karadi A, Kajjidoni ST. Genetic diversity and principal component analysis for grain quality and productivity traits in sorghum [*Sorghum bicolor* (L.) Moench]. *Int. J. Curr. Microbiol. App. Sci*. 2019;8(9): 839-850.
7. Searle SR. Phenotypic, genotypic and environmental correlations. *Biometrics*. 1961;17:474-480.
8. Dewey D, Lu KH. A correlation and path coefficient analysis in crested wheat grass seed production. *Agron. J.* 1959;51: 515-518.
9. Dubey MK, Pandey PK, Shrotria PK, Chauhan P, Pandey G. Assessment of variability, correlation and path analysis for yield and yield related traits in sorghum genotypes. *Green Farming*. 2016;7(6):1291-1295.
10. Jain SK, Elangovan M, Patel NV. Correlation and path coefficient analysis for agronomical traits in forage sorghum (*Sorghum bicolor* L. Moench). *Indian Journal Plant Genet. Resou.* 2017;23(1):15-1.
11. Rohila N, Arya S, Kumari P. Correlation and path coefficient analysis of various sorghum genotypes (*Sorghum bicolor* L. Moench) for quality traits. *Electronic Journal of Plant Breeding*. 2020;11(01):306-30.
12. Prasad BVV, Sridhar V. Assessment of variability correlation and path analysis for yield and yield related traits in yellow pericarp sorghum germplasm line [*Sorghum bicolor* (L.) Moench]. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(1): 870-873.
13. Malaghan S, Kajjidoni ST. Character association and path analysis of grain yield in rabi sorghum [*Sorghum bicolor* (L.) Moench]. *International Journal of Chemical Studies*. 2019;7(1):2309-2313.

© 2023 Nikhil et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/108724>