



# Effects of Auxin and Gibberellins Hormones on Regeneration of Pineapple Propagules from Crown Leaves (with Buds) in Kilifi County Kenya

Monthe Eunice <sup>a</sup>, Ndiso James <sup>a</sup>, Gogo Elisha Otieno <sup>a</sup>  
and Muti Simon <sup>a\*</sup>

<sup>a</sup> Pwani University, P.O. Box 19, Kilifi, Kenya.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Authors ME and MS designed and conducted the study, collected the data and performed the statistical analysis and inferences, then wrote the first draft of the manuscript. Authors NJ and GEO did statistical data analysis, did Manuscript proof reading and corrections and literature searches and citations. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JAERI/2024/v25i1575

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111586>

**Original Research Article**

**Received: 15/11/2023**  
**Accepted: 18/01/2024**  
**Published: 02/02/2024**

## **ABSTRACT**

Kilifi county has a large potential of over 2,000 Ha suitable for pineapple cultivation that has not been fully exploited due to lack of adequate clean planting material. Thus, the aim of the study was to evaluate the effects of auxin and gibberellin hormones on regeneration of pineapple propagules from pineapple crown leaves (with buds). The study was carried out at Pwani University Crop Science farm. A randomized complete block design with split plot arrangement replicated thrice

\*Corresponding author: E-mail: s.muti@pu.ac.ke;

was used. Treatments included auxin and gibberellins hormones where the cut crown leaves (with buds) were dipped and planted at a spacing of 4 cm by 6 cm in plastic containers filled with coir dust. Five plastic containers were used for each hormone treatment. In each container, three plants were tagged for data collection. The plastic containers (the plots) measured 30cm by 36cm by 15cm height. Data collected from the tagged plants included, time to shoot emergence, plant height, leaf length and breadth, number of leaves per plant, final root length and final number of roots. These were used to determine effects of auxin, gibberellins on growth of the propagules up to a height of 15-20 cm. Obtained data was subjected to ANOVA analysis using Genstat Statistical software 14<sup>th</sup> Edition and means obtained compared using Tukey's honest significant difference test ( $p \leq 0.05$ ). The results indicated auxin treated propagules had 50.7%, 50.6%, 64.2%, 52.6%, 24.4%, 10.4%, 39.0%, and 9.9% higher final values of plant weight, shoot weight, dry shoot weight, fresh root weight, plant height, leaf number, root number, dry root weight and root length, respectively, compared to those treated with gibberellins. The study also showed that auxin treated propagules had their initial roots originating from the base of the seedlings, while gibberellins treated propagules had their initial roots originating from the base of the cut crown leaves.

**Keywords:** *Pineapple; crown leaves; propagules; regeneration; auxin; gibberellins.*

## 1. INTRODUCTION

Pineapple (*Ananas Comosus* (L.)) is an important tropical fruit grown in Kenya for domestic and export market in processed form and as fresh fruits [1,2,3]. Main growing areas include Thika, Malindi/Magarini and Kisii where the climate is mild with temperatures ranging between 16 and 32°C. The major exporter of processed pineapple products in Kenya is Delmonte Kenya Ltd situated in Thika and grows pineapples year-round. Important varieties grown include Smooth Cayenne, Queen, and newly developed Delmonte varieties such as Del Monte Gold®, Extra Sweet, Honeyglow®, and Pinkglow® [1,3]. The functional benefits of eating pineapples include protection against macular degeneration, antioxidant protection and immune support, potential anti-inflammatory and digestive benefits, manganese and thiamin for energy production and antioxidant defenses [4]. The nutritional benefits include helping reduce malnutrition that is rampant in the area, especially for children under five years and the elderly [5,6]. It is rich in vitamins such as vitamins A, B, C and minerals that include Manganese which prevents osteoporosis. It has no fat and is therefore cholesterol free [4].

In Kilifi county, pineapple cultivation occurs in Malindi and Magarini sub-counties and is a major source of employment and income for majority of the communities in these areas of Kilifi County through sale of fresh fruits, pineapple juice and exports [7]. The variety grown is the smooth cayenne and is highly suitable for fresh market and canning. Other local varieties grown include Marafa Sweet and Mwanazi [3]. The demand for fresh sweet pineapples from Kilifi county is high

and the County has never met the increased demand yet there exists a large potential of over 2,000 ha for expansion of pineapple farming. Pineapple farming in Kilifi County is faced by a number of challenges which include inadequate availability of clean planting material, high initial investment costs, drought, high transport cost to markets, pests and diseases [8]. Of these challenges, availability of planting materials is the most limiting factor [8]. Currently, pineapple planting material is obtained from pineapple suckers, slips and sometimes from pineapple crowns. To get these materials ready for planting takes about 18-24 months to reach maturity and harvest [3]. This is a long period to wait. The current acreage of pineapple growing area in Kilifi county is estimated at 2,000 ha and 20,000ha for the larger Coastal region [7]. Using the required seed rate of 37,000-48,000 suckers per ha, over 80 million suckers are required for planting every growing season [9]. However, the current production of planting material in Kilifi County is much less, between 10,000-15,000 suckers per ha due to drought, disease infestation and poor soils, giving a deficit of 50 million suckers per planting season [7]. This is what limits expansion of pineapple farming in the county and Coastal region as a whole. In order to cover the deficit, there is need to explore other scientific methods of producing and increasing disease-free planting materials within the shortest time possible without compromising on the quality of final product.

### 1.1 Plant Hormones

Plant hormones are substances that regulate plant growth, development and response to

environmental stimuli when applied in small quantities [10]. Piloting of this study had indicated that growth and development of plantlets from crown leaf buds without use of any hormones took considerably a longer time (more than 3-4 months) for the shoots or roots to emerge. Hence auxin and gibberellins hormones were in cooperated in the study to evaluate their effect in regeneration of the propagules. Gibberellins hormones (GA) are responsible for inducing cell extension, expansion and differentiation thereby promoting root and shoot development. Auxin hormones induce and stimulate cell division and elongation in plants thereby promoting shoot and root growth [11]. In the elongation process, auxin alters the plant wall plasticity making it easier for the plant to grow upwards. Auxin also influences root formation. Gibberellins causes increased growth of the shoots, as a result of increased length of stem internodes and leaves and enhances apical dominance [12,11]. While GA has certain similarities to the auxins important differences are that: auxins greatly increases cell-extension in excised tissue sections, whereas GA has little effect; GA induces marked cell extension in shoots of some intact plants, whereas exogenous auxins have little effect; auxins strongly influence root growth, but GA does not. For some plant species, treatment with GA does not stimulate growth of intact roots, though some root sections do respond by increased growth. High concentrations of GA results in increased dry weight due to increased carbon fixation as a result of increased leaf growth. GA breaks dormancy of seeds, potato tubers and dormancy of shoot internodes and buds. Gibberellins act in plants by removing growth limitations by inducing degradation of the growth-inhibiting DELLA proteins [13].

In Kenya, there are limited studies on regeneration of pineapple planting material from pineapple crown leaves for rapid growth of pineapple plantlets in vivo [14]. Therefore, this study was conceived to evaluate the use of pineapple crown leaves treated with auxinand gibberellin hormones for mass regeneration of planting material for enhancing pineapple production in Kilifi County. The hypothesis under test was that, auxin and gibberellins hormones had no effect on generation and growth of pineapple propagules from crown leaves.

## 2. METHODOLOGY

### 2.1 Study Site

The study was conducted from January to August of 2022 at Pwani University Crop Science farm located 60 km from Mombasa lying between latitudes 3°S and 4°S and longitudes 39°E and 40° E, and at an altitude of about 30m above sea level (ASL) in Coastal lowland zone 3 (CL3) [15]. The average annual rainfall ranged between 600-1100 mm. Long rains in the region normally occur between April and July while the short rains occur from September to December. Annual mean temperatures ranged between 28°C - 32.0°C [16]. The pre-dominant of soil type in the study area was sandy-loam characterized by poor water holding capacity due to low organic matter content and poor in nutrients [17].

### 2.2 Material Used

These included: pineapple crowns which were obtained from local pineapple markets selected from big mature pineapple fruits with clean crown leaves of uniform size with a bud eye [3]. Plant hormones used were auxin under registered name as "Azatone" or "Rootex" (containing 33% auxin). It was sourced from Nairobi Agrovet Shops. Gibberellins was sourced from Amiran Kenya Ltd, in Nairobi under registered name as "Tivag 40 SL" (soluble liquid, gibberellins at 40g/L).

### 2.3 Experimental Design and Treatments

A completely randomized block design with split plot arrangement of treatments, replicated thrice was used (Fig. 1). Two treatments were involved, namely auxin (at 33%) and gibberellins (at 40gm per liter) hormones.

The treatments consisted of auxin and gibberellins. Allocation of the treatments in the plots was done randomly (Fig. 1). Each plot consisted of a plastic box measuring 30 cm by 36 cm by 15cm height, and filled with the coir media up to 10cm. Spacing between plots was 5 cm and between blocks 15 cm (Fig. 1). Spacing of the crown leaves (with buds) between rows was 6 cm and within rows was 4 cm, giving a total of 9 plants per plot. With a total of 30 plots, (15 for auxin and 15 for gibberellins) this gave a grand total of 270 plants for the whole experiment.

|                   | REP I | REP II | REP II |
|-------------------|-------|--------|--------|
| Auxin<br>A        | A1    | A5     | A3     |
|                   | A2    | A      | A      |
|                   | A     | A3     | A4     |
|                   | A3    | A      | A      |
|                   | A     | A4     | A1     |
|                   | A4    | A      | A      |
|                   | A     | A2     | A5     |
|                   | A5    | A      | A      |
| Gibberellins<br>G | G1    | A1     | A2     |
|                   | G     | G4     | G2     |
|                   | G2    | G      | G      |
|                   | G     | G3     | G5     |
|                   | G3    | G      | G      |
|                   | G     | G2     | G4     |
|                   | G4    | G      | G      |
|                   | G     | G5     | G1     |
|                   | G     | G      |        |
|                   | G5    | G3     |        |

**Fig. 1. Experimental plot layout in split-plot arrangement**

Key: Auxin = A; Gibberellins = G

#### 2.4 Preparation of Crown Leaf Cuttings, Treatment and Planting in Propagation Trays

One or two healthy pineapple crown leaves of more than 5 cm in length, were carefully and individually cut off from the stem of selected pineapple crowns, and checked for healthy eye buds. Crown leaf buds are shared by two leaves at leaf-overlap; hence either a single leaf or two leaves were hived off to obtain an intact bud (Plate 1). The hived off leaves were dipped in

70% alcohol for sterilization, and in a solution of Ridomil fungicide for treatment against fungal pathogens. The leaves were then dipped in auxin hormone powder (Azatone) or in gibberellins solution depending on the respective treatment (Plate 1).

At initial stages, the treated crown leaves were raised in large metallic propagation trays containing coir media to facilitate selection of healthy “germinating” pineapple propagules from crown leaf buds (Plate 2).



**Plate 1. Treatment of individual crown leaves (with a bud) with auxin hormone**



**Plate 2. Propagation tray, with coir media and planted crown leaf cuttings**

Before planting into propagation trays, the coir media was watered to saturation and left for 5 days to allow for free drainage and settling of the media. After planting in the propagation trays (measuring 90 cm by 60 cm) and irrigating uniformly to field capacity using Watering-Can, the propagation trays were transferred into propagation structures roofed with fiber glass for maintaining adequate lighting, constant temperature (of about 29-32°C) and high relative humidity (of around 60-80%) which is necessary for sprouting of the leaf buds (Plate 3). Regular watering was maintained every two days to ensure adequate moisture in the media. During the study period, no pests or diseases were noted.

### **2.5 Selection and Transplanting Germinated Propagules into Plastic Containers**

About one month after planting, healthy well sprouted and rooted propagules that had attained 3 cm in height were selected for transplanting into plastic containers filled with coir (Plates 3, 4 and 5).

### **2.6 Data Collection**

In each plot (plastic container) three plants were randomly selected and tagged (Plate 5). It is from these tagged plants where data was collected every 7 days after transplanting until the propagules attained 15-20 cm in height. Data collected included, plant height, plant weight, fresh and dry shoot weight, leaf length and width, number of leaves, stem diameter, root length, fresh and dry root weight and number of roots

per propagule. This experiment was conducted from January to August, 2022.

### **2.7 Determination of Parameters**

Plant height was determined using a meter rule, from the base of the propagule to the tip of the plant. The number of days to shoot emergence was determined from the day of planting until a new shoot emerged above media surface. Leaf length was determined using a ruler from its attachment to the stem to the tip of the leaf blade, while the leaf width, was determined using a ruler at its widest part across the midrib. The number of new leaves were determined by observing and counting the new leaves as they emerged above the media surface. Final root length was determined by uprooting and measuring the root length using a ruler. The final number of roots per propagule was determined by physically counting the number of roots per plant. Stem diameter was determined by placing a string around the stem and obtaining the length on a ruler and dividing the obtained circumference by value of pie (3.14). Shoots and root weights were determined by weighing the isolated samples using an electronic weighing balance.

### **2.8 Data Analysis**

The collected data for each variable was subjected to analysis of variance (ANOVA) using Genstat statistical software 14<sup>th</sup> edition and the means obtained subjected to Tukey's Honestly significant difference test for comparison at 5% level of significant difference.



Plate 3. One-month old germinated pineapple propagule seedlings



Plate 4. Rooted propagule at 3 cm in height ready for transplanting



Plate 5. Transplanted and three tagged propagules per plot for data collection

### 3. RESULTS


#### 3.1 Effects of Hormones on Growth Parameters of Pineapple Propagules

The results indicate that application of plant hormones on crown leaves (with buds) resulted in significant differences ( $P \leq 0.05$ ) in propagule growth parameters (Table 1 and Fig. 2). In general, auxin hormones resulted in higher values of plant parameters than gibberellins hormones (Table 1 and Fig. 2). However, they attained peak stages of growth at the same time (Fig. 2). One notable difference due to hormones treatment was on the place of origin of initial roots during the sprouting stage (Plate 6). In auxin treated crown leaves, initial roots originated from the base of the young germinating seedlings, while in gibberellins treated crown leaves, initial roots originated from

the base of the cut part of the crown leaves (Plate 6).

For auxins treated leaves (Plate 6a), roots originated from base of the sprouting seedlings, while for gibberellins treated leaves (Plate 6b), roots originated from the base of cut leaves. The results also indicated that auxin treated propagules had significantly 10.5%, 15.4%, 29.0%, 15.8% and 7.1% more leaves per plant, taller plants, larger stem diameter, longer leaf lengths and leaf widths respectively than gibberellins treated propagules (Tables 1 and 2).

Auxin hormones maintained significantly higher values for most tested parameters compared to gibberellins, for similar growth period. The growth pattern of most propagule parameters followed a parabolic (polynomial) trend (Fig. 2).



**Fig. 2. Effects of auxin and gibberellins on a) leaf length, b) plant height, c) leaf number, and d) leaf breadth. The period at which most growth parameters attained peak growth due to auxin and gibberellins hormones appear to coincide**



Plate 6. Site of origin of initial roots in hormone treated pineapple crown leaf

Table 1. Effects of hormones on growth parameters of pineapple propagule at 10th week after transplanting

| Hormones     | Propagule parameters |        |               |             |              |
|--------------|----------------------|--------|---------------|-------------|--------------|
|              | Leaf number          | Height | Stem diameter | Leaf length | Leaf breadth |
| Auxin        | 11.65a               | 17.29a | 6.66a         | 14.64a      | 2.26a        |
| Gibberellins | 10.43b               | 14.63b | 6.06b         | 12.33b      | 2.10b        |
| Mean         | 11.0                 | 16.0   | 6.4           | 13.5        | 2.2          |
| (P < 0.05)   | <.0001               | 0.000  | 0.000         | 0.000       | 0.012        |
| Sed          | 0.150                | 0.250  | 0.08          | 0.21        | 0.03         |
| CV%          | 1.4                  | 1.6    | 1.3           | 1.6         | 1.4          |

Means at the same column that do not share a letter are significantly different

Table 2. Effects of hormones on growth parameters of pineapple propagule at 17<sup>th</sup> week after planting

| Media types  | Plant weight | Shoot weight | Dry shoot weight | Fresh Root weight | Plant Height | Leaf number | Root number | Dry root weight | Root length |
|--------------|--------------|--------------|------------------|-------------------|--------------|-------------|-------------|-----------------|-------------|
| Auxin        | 78.9a        | 71.4a        | 12.0a            | 7.6a              | 26.2a        | 15.4a       | 30.8a       | 4.8a            | 27.2a       |
| Gibberellins | 38.9b        | 35.3b        | 4.3b             | 3.6b              | 19.8b        | 13.8b       | 18.8b       | 0.95b           | 24.5a       |
| Mean         | 58.9         | 53.35        | 8.15             | 5.6               | 23           | 14.6        | 24.8        | 2.9             | 25.85       |
| P-Value      | 0.000        | 0.000        | 0.000            | 0.000             | 0.000        | 0.006       | 0.000       | 0.000           | 0.070       |
| Sed          | 11.7         | 10.7         | 1.45             | 1.17              | 1.25         | 0.73        | 2.48        | 0.24            | 1.54        |
| % CV         | 19.9         | 20.1         | 17.8             | 20.9              | 5.4          | 5.0         | 10.0        | 8.3             | 6.0         |

Means in the same column that do not share a letter are significantly different

### 3.2 Effects of Hormones on Growth Parameters of Pineapple Propagules at 17<sup>th</sup> Week After Planting

Tables 1 and 2 shows that application of auxin hormones on crown leaves (with buds) resulted in significant differences ( $P \leq 0.05$ ) in propagule growth parameters compared to those treated

with gibberellins hormones. Auxin treated propagules occasioned significantly higher values of plant height, longer roots, higher number of leaves and roots per plant than gibberellins treated propagules (Table 2). Thus, auxin treated propagules had 50.7%, 50.6%, 64.2%, 52.6%, 24.4%, 10.4%, 39.0%, 9.9% and 5.2% higher values of plant weight, shoot



weight, dry shoot weight, fresh root weight, plant height, leaf number, root number, dry root weight, and root length, respectively, compared to those treated with gibberellins.

## 4. DISCUSSION

### 4.1 Effects of Hormones on Pineapple Propagule Growth Parameters

Auxin and gibberellins had significant different effects on propagule growth parameters, namely plant height, leaf length and width, number of leaves, stem diameter, root length and number of roots per propagule at 17<sup>th</sup> week after planting. Similar observations were by Suman et al. [18] and [19] who reported that, in pineapple species, auxin and gibberellins increase cell division, elongation and expansion activities. In this study, auxin hormones in particular resulted in higher number of leaves, taller plants, larger stem diameters, leaf lengths and leaf widths than gibberellins hormones. This could partly be attributed to the functional role and mode of action of each type of hormone on plant cells during growth processes. Auxins are generally known to promote cell elongation, while gibberellins are known to promote lateral cell expansion amongst other functions. This explains why auxin treated propagules had longer leaves, taller plants and wider stem diameter than those treated with gibberellins hormones. Initial roots establishment is key in water and nutrient uptake and this determines the rate of growth and assimilation of dry matter of the plantlets. da Costa [20] has pointed out involvement of auxin in development and initiation of adventitious and lateral roots. The adventitious roots appeared to develop from the base of the plantlets, most probably induced by auxin hormones. This perhaps could explain why auxin treated propagules exhibited faster growth rate than gibberellins hormones. In gibberellins treated propagules, the roots originated from the cut leaf base. Any absorption of water and nutrients were first processed in the callus tissue of the cut leaf, thus perhaps delaying nutrient transfer to the newly growing shoot unlike in auxin treated propagules. These results agree with similar observations by Khadr et al. [21] and [22] who reported that auxin primarily induced cell division and elongation, stimulated shoot elongation and root initiation and also vascular differentiation in carrots.

### 4.2 Effects of Hormones on Growth Parameters of Pineapple Propagules at 17<sup>th</sup> Week After Planting

Application of auxin hormones on crown leaf resulted in propagules that were taller and heavier in weight with longer roots, higher number of leaves and roots per plant than those treated with gibberellins. This would suggest that auxin hormones had a functional role of enhancing growth through increased cell division of most meristematic tissues and therefore faster accumulation of biomass by the pineapple propagules than gibberellins hormones. According to Gomes et al. [11] auxins greatly increases and mediates [23] cell-extension in excised tissue sections, whereas GA has little effect. Besides, during the elongation process, auxin alters the plant wall plasticity making it easier for the plant to grow upwards and influences root formation downwards. Brumos et al. [24] reported that Local auxin biosynthesis is a key regulator of plant development. This perhaps explains why auxin treated propagules were bigger and taller than those treated with gibberellins since the main function of auxin hormone is on growth regulation. Most literature cites the major role of gibberellins as being more of cell differentiation and therefore organ development than biomass accumulation [25]. This observation appears to agree with those reported by Rademacher [26], who observed that among the functions of gibberellins include delaying senescence in fruits, is involved in leaf expansion, break bud and seed dormancy, promote bolting in cabbages and beet, facilitate elongation of fruits such as apples and enhancing their shape, all of which point to developmental functions than growth.

## 5. CONCLUSIONS

This study has shown that it is possible to mass generate pineapple seedlings from crown leaves with eye-buds. The study has shown that auxin and gibberellin hormones can hasten regeneration of pineapple seedlings from crown leaves with eye-buds. Although both hormones influenced the growth parameters in a similar way, auxin hormones had profound effects on the different parameters compared to gibberellins. Auxin values for the different parameters were way above those of gibberellins. These findings indicate that farmers now have at their disposal a vast potential for mass generation of clean planting material using pineapple crown leaves and auxin hormones.

## 6. RECOMMENDATIONS

There is need to ascertain the mode of action of both hormones in terms of initiating root development either from the cut leaf base or from the base of the plantlets. There is also need to compare how long it takes for the pineapple propagules generated from crown leaf buds to reach flowering and produce a mature fruit, their shelf life and quality with planting material sourced from different parts of the plant such as suckers, slips, crowns, among others.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Bartholomew DP. MD-2'pineapple transforms the world's pineapple fresh fruit export industry. *Pine. News.* 2009;16:2–5:15.
2. FAO. Major Tropical Fruits: Preliminary results 2021. Rome; 2022.
3. Hossain MF. World pineapple production: An overview. *African Journal of Food, Agriculture, Nutrition and Development.* 2016;16(4):11443-11456.
4. Chang YK, d'Eeckenbrugge GC, Sanewski GM. Breeding and variety improvement. *The pineapple: botany, production and uses.* CABI Publishing, Wallingford, UK; 2003.
5. Muti SM, Kibe AM, Nge'tich W, Muindi E. Effects of Coir Dust Mulch on Evapotranspiration of PH4 Maize in Coastal Region of Kenya. *Journal of Agriculture and Ecology Research International.* 2017; 10(3):1-16.
6. Reinhardt DHR, Bartholomew DP, Souza FVD, Carvalho ACPPD, Pádua TRPD, Junghans DT, Matos APD. Advances in pineapple plant propagation. *Revista Brasileira de Fruticultura.* 2018;40.
7. Kilifi County profile. Kilifi Integrated Development Programme; 2020.
8. Muti, Simon, James Ndiso, Bahati Abdallah, Eunice Monthe, Serah Nzau, Doreen Kinoti, and Rebecca Amukhoye. "Status and Challenges Facing Pineapple (Ananas comosus) Production in Magarini Sub-County of Kilifi County in Coastal Kenya." *Journal of Agriculture and Ecology Research International.* 2024;25(1):1-18.
9. Hossain MF, Akhtar S, Anwar M. Nutritional value and medicinal benefits of pineapple. *International Journal of Nutrition and Food Sciences.* 2015;4(1):84-88.
10. Davies, Peter J, ed. *Plant hormones: physiology, biochemistry and molecular biology.* Springer Science & Business Media; 2013.
11. Gomes GLB, Scortecci KC. Auxin and its role in plant development: structure, signalling, regulation and response mechanisms. *Plant Biol (Stuttg).* 2021; (6):894-904.
12. Brian PW. Effects of gibberellins on plant growth and development. *Biological reviews.* 1959: 34:37-77.
13. Hedden P, Valerie S. A century of gibberellin research." *Journal of plant growth regulation.* 2015;34:740-760.
14. Chotangui AH, Kenhoung C, Mandou MS, Kouam EB. *Biosciences and Plant Biology.* International Journal of Current Research in Biosciences Plant Biology. 2019;6(11):1-8.
15. Jaetzold R, Hornetz B, Shisanya CA, Schmidt H. *Farm management handbook of Kenya Vol I-IV (Western, Central, Eastern, Nyanza, Southern Rift Valley, Northern Rift Valley and Coast).* Nairobi, Government Printers; 2012.
16. Mnyika AW, Muti SM, Gogo EO. Superabsorbent polymer and rabbit manure improve soil moisture, growth and yield of eggplant (*Solanum melongena* L). *NASS Journal of Agricultural Sciences.* 2020;2(1):12-20.
17. Daniel KA, Muindi EM, Gogo EO, Muti S. Performance of Brassica rapa and Brassica oleracea under Shade net within Coastal Environment. *Journal of Agriculture and Ecology Research International.* 2022;23(3):45-59.
18. Suman M, Sangma PD, Meghawal DR, Sahu OP. Effect of plant growth regulators on fruit crops. *Journal of Pharmacognosy and Phytochemistry.* 2017;6(2):331-337.
19. Zhou T, Geng Y, Ji C, Xu X, Wang H, Pan J, Lausch A. Prediction of soil organic carbon and the C: N ratio on a national scale using machine learning and satellite data: A comparison between Sentinel-2, Sentinel-3 and Landsat-8 images. *Science of the Total Environment.* 2021; 755:142661.
20. da Costa, Cibele Tesser, Remko Offringa, and Arthur Germano Fett-Neto. "The role of auxin transporters and receptors in

- adventitious rooting of Arabidopsis thaliana pre-etiolated flooded seedlings." Plant Science. 2020;(290):110294.
21. Khadr A, Wang GL, Wang YH, Zhang RR, Wang XR, Xu ZS, Xiong AS. Effects of auxin (indole-3-butyric acid) on growth characteristics, lignification, and expression profiles of genes involved in lignin biosynthesis in carrot taproot. Peer Journal. 2020;8:e10492.
  22. Brian PW, Hemming HG. A relation between the effects of gibberellic acid and indolyacetic acid on plant cell extension. Nature, Lond. 1957a; (179): 417.
  23. Du M, Spalding EP, Gray WM. Rapid auxin mediated cell expansion. Annual Review of Plant Biology. 2020;(71):1–24.
  24. Brumos J, Robles LM, Yun J, Vu TC, Jackson S, Alonso JS, Stepanova AN. Local auxin biosynthesis is a key regulator of plant development. Developmental Cell. 2018;(47):306–318.
  25. Enders TA, Strader LC. Auxin activity: past, present, and future. American Journal of Botany. 2015;(102):180–196.
  26. Rademacher W. Plant growth regulators: backgrounds and uses in plant production. Journal of plant growth regulation. 2015;34:845-872.

© 2024 Eunice et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/111586>