



# A Rare Case of Community-acquired Meningitis Caused by *Staphylococcus Aureus* at FANN Teaching Hospital, Dakar, Senegal

Daye KA <sup>a\*</sup>, Aboubakar Taha Ibrahim <sup>a</sup>,  
Aboubakar Sidikh Badiane <sup>a</sup>, Catherine Sarr <sup>a</sup>,  
Ndéye Marème Dieng <sup>a</sup>, Khardiata Diallo Mbaye <sup>a</sup>,  
Ndéye Aissatou Lakhe <sup>a</sup>,  
Viviane Marie Pierre Cisse Diallo <sup>a</sup> and Moussa Seydi <sup>a</sup>

<sup>a</sup> Department of Infectious and Tropical Diseases, Fann Hospital, Senegal.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ajrid/2024/v15i11387>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123894>

Case Report

Received: 30/07/2024  
Accepted: 01/10/2024  
Published: 04/10/2024

## ABSTRACT

Bacterial meningitis in Africa is caused mainly by three bacteria: *Streptococcus pneumoniae*, *Neisseria meningitidis* and *Haemophilus influenzae B*. *Staphylococcus aureus* is a rare cause of bacterial meningitis, with an estimated prevalence rate of between 0.89% and 5% of bacterial meningitis cases in Africa.

\*Corresponding author: E-mail: [dayeka10@gmail.com](mailto:dayeka10@gmail.com);

**Cite as:** KA, Daye, Aboubakar Taha Ibrahim, Aboubakar Sidikh Badiane, Catherine Sarr, Ndéye Marème Dieng, Khardiata Diallo Mbaye, Ndéye Aissatou Lakhe, Viviane Marie Pierre Cisse Diallo, and Moussa Seydi. 2024. "A Rare Case of Community-Acquired Meningitis Caused by *Staphylococcus Aureus* at FANN Teaching Hospital, Dakar, Senegal". *Asian Journal of Research in Infectious Diseases* 15 (11):13-17. <https://doi.org/10.9734/ajrid/2024/v15i11387>.

The aim was to highlight the clinical features, diagnostic challenges, and treatment outcomes of this rare case.

We report the case of a 36-year-old woman with dental caries admitted for acute meningitis. A cerebrospinal fluid (CSF) culture revealed the presence of *Staphylococcus aureus*. The patient had a good clinical evolution with intravenous antibiotic therapy. Following a review of the literature, we discuss the characteristics of *Staphylococcus aureus* meningitis.

**Keywords:** Meningitis; community; *Staphylococcus aureus*.

## 1. INTRODUCTION

Bacterial meningitis is a serious, life-threatening infection and a major public health problem in sub-Saharan Africa [1]. The most frequent causes of meningitis in Africa are: *Streptococcus pneumoniae*, *Neisseria meningitidis* and *Haemophilus influenzae* type B [2]. Meningitis due to *Staphylococcus* is a rare but a potentially fatal disease. It often occurs after neurosurgical procedures or in immunocompromised patients [3]. The most commonly used antibiotics include ceftriaxone and methicillin.

There are few data on its epidemiology. Nevertheless, its incidence is on the rise, particularly community-acquired meningitis. The prevalence in Africa is estimated to be between 0.89% and 5% of all causes of bacterial meningitis [2,4]. We report a rare form of community-acquired *Staphylococcus aureus* meningitis in a patient with no previous history of hospitalization or neurosurgical intervention.

## 2. CASE PRESENTATION

The patient was a 36-year-old female, unvaccinated against pneumococcus and meningococcus. She had no past medical history of diabetes, sickle cell anemia or any other medical or surgical history. She presented with an intense helmet-type headache associated with post prandial vomiting that did not relieve the headache. She also reported photophobia, vertigo and tinnitus. Physical examination revealed a fever of 39.5°C with no sweating or chills, and a stiff neck with Kernig's sign positive. Oral examination revealed 5 category I decayed teeth, 2 category IV decayed teeth and presence of tartar (Fig. 1). The genitourinary examination was normal, as was the rest of the physical examination.

Based on her clinical presentation, the following hypothesis were evoked: infectious meningitis (bacterial, viral), an intracranial abscess or a subarachnoid hemorrhage.

Her complete blood count showed a white blood cell count of  $15.82 \times 10^9/L$ , a haemoglobin level of 12.1g/dL and a platelet count of 249,000/mm<sup>3</sup>. C-Reactive Protein was 58.3 mg/L and procalcitonin 1.51 ng/mL (Table 1).

A brain CT scan (Computed Tomography Scan) showed bulging of the cerebellar tonsils in the foramen magnum (Chiari I malformation) and normal-sized ventricles. There was no cerebral expansive lesion, no mass effect and no peripheral contrast enhancement.

CSF analysis showed a clear macroscopic appearance. Cytological examination of the CSF revealed a pleocytosis of 580 leucocytes, 75% of which were lymphocytes, 20% unaltered polynuclear neutrophils and 5% altered polynuclear neutrophils. Biochemistry revealed an elevated CSF protein of 1.35 g/L, with a blood glucose/CSF glucose ratio of 0.66. Direct microscopic examination revealed the presence of Gram-positive cocci. Bacteriological culture of the CSF revealed methicillin-sensitive *Staphylococcus aureus*. The Xpert MTB Rif test on the CSF was negative, and PCR for viruses was not performed. Two pairs of blood cultures were collected, with negative results.

HIV (human immunodeficiency virus) serology was negative, serum protein electrophoresis showed an albumin/globulin ratio of 1.4 with no qualitative abnormalities. Fasting plasma glucose was 1.09 g/l and glycated hemoglobin was 6.1%.

A definitive diagnosis of methicillin-sensitive *Staphylococcus aureus* meningitis in an immunocompetent patient with a dental portal of entry was retained. The patient was hospitalized and treated with Ceftriaxone 4 grammes per day in a single dose. The Headache resolved within 5 days, and stable apyrexia was observed after two days of treatment. No episodes of vomiting were reported. All clinical signs resolved after 8 days of antibiotic therapy with normalization of the inflammatory biomarkers initially described.

No follow-up lumbar puncture was performed in view of the improvement in her clinical state.



**Fig. 1. Image of dental caries (oral cavity)**

**Table 1. Clinical Investigations Summary**

Investigation	Result
White Blood Cell Count	15.82 × 10 <sup>9</sup> /L
Hemoglobin Level	12.1 g/dL
Platelet Count	249,000/mm <sup>3</sup>
C-Reactive Protein	58.3 mg/L
Procalcitonin	1.51 ng/mL
CSF Appearance	Clear
CSF Pleocytosis	580 leukocytes
CSF Protein	1.35 g/L
Blood Glucose/CSF Glucose Ratio	0.66
CSF Culture	Methicillin-sensitive <i>Staphylococcus aureus</i>

### 3. DISCUSSION

Acute bacterial meningitis represents a heavy burden of morbidity and mortality in sub-Saharan Africa, with an incidence of up to 1,000/100,000 cases [2,5,6]. The three most frequently encountered germs are *Streptococcus pneumoniae*, *Neisseria meningitidis* and *Haemophilus influenzae B*. Staphylococcal meningitis accounts for no more than 10% of all bacterial meningitis [7]. The majority of these cases are secondary to neurosurgical procedures, and usually involve methicillin-resistant *Staphylococcus aureus* (MRSA) [7]. We describe a case of acute community acquired meningitis caused by methicillin-sensitive *Staphylococcus aureus*. This is the first case documented in the infectious and tropical

diseases department of CHNU Fann since 2018. This rarity is in line with data from existing literature. In a study by Aguilar J et al, there were only 17 cases of Staphylococcal meningitis out of a total of 668 cases of meningitis recorded over 9 years [8].

Most studies have focused on MRSA because of its severity and the difficulties involved in its management [8]. Moreover, a study conducted in Rabat from 2016 to 2018 revealed a significant resistance rate of 92% to penicillin G [9]. But with the current tragedy of antibiotic resistance, it would be pertinent to study the epidemiological characteristics of MSSA meningitis to prevent the selection of mutant strains.

In our patient, no immunosuppressive factors were found that could have predisposed her to a serious infection. Furthermore, at the time of diagnosis, a thorough history taking enabled us to determine the progressive nature of her symptoms, which was insidious and had been evolving for a fortnight. This clinical presentation is similar to that of cases reported by Pinet P et al [7], and Minatogawa A et al [10], who reported fifteen and seven days of evolution respectively. However, this is in contrast with most cases of acute bacterial meningitis, wherein the clinical signs develop abruptly over a period of a few days. In a large review of cases, Pintado Vicente et al found that clinical evolution was acute (≤ 7 days) in 89% of patients [11].

In contrast to the results of some studies which described frequent extra-neurological signs [7,12], associated with bacteremia in 75% of cases [13], our patient presented only neurological and general signs without bacteremia. This is perhaps one of the limitations of our clinical case. For, based on a single patient, it is difficult to generalize the data to other similar situations.

Traditionally, dental infections can be complicated by intracranial suppuration. However, in view of our patient's poor oral condition, and the germ isolated from the CSF, we wondered whether a meningitis-like complication was also possible. In the literature, it has been widely described that *Staphylococcus aureus* can establish cariogenic biofilms [14,15,16]. This can be the starting point for a meningitis-like complication. To the best of our knowledge, previous studies on *Staphylococcus aureus* meningitis have not focused on the germ's portal of entry.

Analysis of the CSF showed that the fluid was clear, predominantly lymphocytic. This is not in line with known data on acute bacterial meningitis, or even specifically *Staphylococcus aureus* meningitis [17], which is often predominantly neutrophilic. The antibiotic ceftriaxone at a dose of 2 g taken prior to admission could explain this discrepancy. This phenomenon is sometimes referred to as “decapitated meningitis”, where early administration of effective antibiotics suppresses the inflammatory response typical of bacterial meningitis, resulting in a decrease in CSF neutrophils and a relative predominance of lymphocytes.

The patient received a 10-day course of intravenous ceftriaxone-based antibiotics, adapted to her SASM microbiologic profile. The results of the antibiotic susceptibility test made it possible to maintain this initial treatment regimen.

#### 4. CONCLUSION

The bacteria most frequently associated with meningitis in Africa are *Streptococcus pneumoniae*, *Neisseria meningitidis* and *Haemophilus influenzae B*. However, in rare cases, other germs may be responsible, as was observed in our patient. Hence the importance of a systematic culture of the CSF in order to establish a precise diagnosis and institute a treatment adapted to the antibiogram.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

#### CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Meningococcal meningitis. [cited 15 July 2024]. Available: <https://www.who.int/fr/news-room/fact-sheets/detail/meningitis>
2. Oordt-Speets AM, Boliijn R, Hoorn RC van, Bhavsar A, Kyaw MH. Global etiology of bacterial meningitis: A systematic review and meta-analysis. PLOS ONE. 11 June 2018;13(6): e0198772.
3. van de Beek D, Brouwer MC, Koedel U, Wall EC. Community-acquired bacterial meningitis. Lancet. 25 Sept 2021;398(10306):1171-83.
4. O. Zender H, Olivier P, Genné D. Méningites bactériennes communautaires aiguës chez l'adulte. Rev Med Suisse. 7 Oct 2009;220(36):1968-74.
5. Daugla DM, Gami JP, Gamougam K, Naibei N, Mbainadji L, Narbé M, et al. Effect of a serogroup A meningococcal conjugate vaccine (PsA–TT) on serogroup A meningococcal meningitis and carriage in Chad: a community study. The Lancet. 4 Janv 2014;383(9911):40-7.
6. Paireau J, Chen A, Broutin H, Grenfell B, Basta NE. Seasonal dynamics of bacterial meningitis: a time-series analysis. Lancet Glob Health. 2016 Jun;4(6):e370-7.
7. Pinet P, Denes E, Garnier F, Durox H, Ducroix-Roubertou S, Weinbreck P. Méningites communautaires à *Staphylococcus aureus* méticilline sensible [Community-acquired meningitis due to methicillin susceptible *Staphylococcus aureus*]. Med Mal Infect. 2010 Mar;40(3):156-60.
8. Aguilar J, Urdy-Cornejo V, Donabedian S, Perri M, Tibbetts R, Zervos M. *Staphylococcus aureus* meningitis: case series and literature review. Medicine (Baltimore). 2010 Mar;89(2):117-125.
9. Al-Selwi AG, Barkat A. Antibiotic resistance of streptococcus pneumoniae, neisseria meningitidis, haemophilus influenzae and staphylococcus aureus in Morocco, National Data: Meta-Analysis. Biomed. Pharmacol. J. 2023 Mar;16(1): 251-63.
10. Minatogawa A, Ohara J, Horinishi Y, Sano C, Ohta R. Meningitis with staphylococcus aureus bacteremia in an older patient with nonspecific symptoms: A case report. Cureus. 2023 Jan 24;15(1): e34153.

11. Pintado V, Pazos R, Jiménez-Mejías ME, Rodríguez-Guardado A, Gil A, García-Lechuz JM, Cabellos C, Chaves F, Domingo P, Ramos A, Pérez-Cecilia E, Domingo D. Methicillin-resistant *Staphylococcus aureus* meningitis in adults: a multicenter study of 86 cases. *Medicine (Baltimore)*. 2012 Jan;91(1):10-17.
12. van Soest TM, Søndermølle MB, Brouwer MC, Chekrouni N, Larsen AR, Petersen A, van Sorge NM, Nielsen H, van de Beek D, Bodilsen J; DASGIB study group. Community-acquired *Staphylococcus aureus* meningitis in adults. *J Infect*. 2023 Mar;86(3):239-244.
13. Vestergaard HH, Larsen L, Brandt C, Hansen BR, Andersen CØ, Lüttichau HR, Helweg-Larsen J, Wiese L, Storgaard M, Nielsen H, Bodilsen J. Normocellular community-acquired bacterial meningitis in adults: A nationwide population-based case series. *Ann Emerg Med*. 2021 Jan;77(1):11-18.
14. Yadav K, Prakash S. Caries dentaires: une approche microbiologique. *J Clin Infect Dis Pract*. 2017 avr;2(1):1-5.
15. O'Donnell LE, Smith K, Williams C, Nile CJ, Lappin DF, Bradshaw D, Lambert M, Robertson DP, Bagg J, Hannah V, Ramage G. Dentures are a reservoir for respiratory pathogens. *J Prosthodont*. 2016 Feb;25(2):99-104.
16. Lewis N, Parmar N, Hussain Z, Baker G, Green I, Howlett J, Kearns A, Cookson B, McDonald A, Wilson M, Ready D. Colonisation of dentures by *Staphylococcus aureus* and MRSA in out-patient and in-patient populations. *Eur J Clin Microbiol Infect Dis*. 2015 Sep;34(9):1823-6.
17. Basri R, Zueter AR, Mohamed Z, Alam MK, Norsa'adah B, Hasan SA, Hasan H, Ahmad F. Burden of bacterial meningitis: A retrospective review on laboratory parameters and factors associated with death in meningitis, kelantan malaysia. *Nagoya J Med Sci*. 2015 Feb;77(1-2):59-68.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/123894>