



## **Ameliorative Efficacy of Phytochemical Content of *Corchorus olitorius* Leaves against Acute Ethanol-induced Oxidative Stress in Wistar Rats**

**Augustine I. Airaodion<sup>1,2\*</sup>, Emmanuel O. Ogbuagu<sup>3</sup>, Ogbonnaya Ewa<sup>4</sup>, Uloaku Ogbuagu<sup>1</sup>, Olaide O. Awosanya<sup>2</sup> and Olukule A. Adekale<sup>5</sup>**

<sup>1</sup>Department of Biochemistry, Federal University of Technology, Owerri, Imo State, Nigeria.

<sup>2</sup>Department of Premedical Science, Educational Advancement Centre, Ibadan, Oyo State, Nigeria.

<sup>3</sup>Department of Pharmacology and Therapeutics, Abia State University, Uturu, Nigeria.

<sup>4</sup>Department of Medical Biochemistry, Gregory University, Uturu, Abia State, Nigeria.

<sup>5</sup>Department of Botany, University of Ibadan, Oyo State, Nigeria.

### **Authors' contributions**

*This work was carried out in collaboration with all authors. Author AIA conceptualized and designed the study and also wrote the manuscript. Author EOO managed the analyses of the study. Authors OE and UO managed the literature searches. Author OOA wrote the protocol while author OAA performed the statistical analysis. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/AJBGMB/2019/v2i230054

#### Editor(s):

(1) Dr. Mohammed Rachidi, Director of Research, Molecular Genetics of Human-Diseases (MGHD), French Polynesia, University Paris Denis Diderot, Paris, France.

#### Reviewers:

(1) Mustafa Sevindik, Biology, Akdeniz University, Turkey

(2) Ochieng O. Anthony, Sumait University, Tanzania.

(3) Sangeetha A/P Arullappan, Universiti Tunku Abdul Rahman (UTAR), Malaysia.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49753>

**Original Research Article**

**Received 05 April 2019**

**Accepted 24 June 2019**

**Published 29 June 2019**

### **ABSTRACT**

**Aim:** This study is aimed at investigating the ameliorative efficacy of methanolic extract of *Corchorus olitorius* leaves against acute ethanol-induced oxidative stress in Wistar rats.

**Methods:** Fresh plants of *C. olitorius* were harvested from the Institute of Agricultural Research and Training, Ibadan. The leaves were dried and extracted using soxhlet apparatus and methanol as the solvent. The methanol was evaporated in a rotary evaporator at 35°C with a yield of 2.17 g which represents a percentage yield of 8.68%. Twenty adult male Wistar rats with body weight between

\*Corresponding author: E-mail: [augustineairaodion@yahoo.com](mailto:augustineairaodion@yahoo.com);

120 and 150 g were used for this study. They were randomly divided into four groups of five rats each. Animals in groups 1 and 2 were administered saline solution while those in groups 3 and 4 were administered *C. olitorius* extract for twenty-one days. The animals were administered the extract and saline solution at a dose of 4 mL per 100 g body weight 12 hourly via oral route of administration. At the end of the treatment, they were fasted overnight and animals in groups 2 and 4 were exposed to a single dose of 70% ethanol at 12 ml/kg body weight to induce oxidative stress. After 12 hours of ethanol administration, the animals were anaesthetized using diethyl ether and were sacrificed. Liver was excised, weighed and homogenized in 50 mmol/L Tris-HCl buffer (pH 7.4) and then centrifuged at 5000 × g for 15 minutes for biochemical analysis. Supernatants were immediately kept frozen for further analysis.

**Results:** Ethanol-induced oxidative stress significantly increased the activities of AST, ALT, LDH, LPO, CAT, SOD and GPX but decrease GSH. These effects were regulated by *C. olitorius* administration.

**Conclusion:** *C. olitorius* was able to remedy the effect of ethanol by regulating the oxidative stress biomarkers, thus possesses ameliorative efficacy against ethanol-induced oxidative stress and can protect the body against free radicals arising from oxidative stress.

**Keywords:** Ameliorative efficacy; *C. olitorius* leaves; ethanol; oxidative stress.

## 1. INTRODUCTION

*Corchorus olitorius* (Malvaceae) is a plant native to both tropical and subtropical regions throughout the world with mallow leaves commonly consumed as a leafy vegetable. The leaves have been reportedly used in ethnomedical practices to treat ache and pain, dysentery, malaria, enteritis, fever, gonorrhoea, pectoral pains and tumors [1]. *C. olitorius* is a green leafy vegetable popularly consumed among the Yorubas of southwestern Nigeria where it is commonly called Ewedu. Among the Igbos of southeastern, Nigeria, it is called Ahihara, while in English, the plant is known as jute mallow or bush okra. *C. olitorius* plant is not found in Nigeria only but also in other countries such as Egypt, Sudan, Malaysia, South America, and the Caribbean [2,3,4]. Nutritional substances; including calcium, potassium, phosphate, iron, ascorbic acid, carotene and large amount of mucilaginous polysaccharides have all been identified in the plant [5]. Medicinally, *C. olitorius* are used as a demulcent, diuretic, purgative, bitter tonic, laxative, refrigerant, carminative and lactagogue [6]. The leaves extract has given positive results in the management of chronic cystitis and dysuria. Its reported high antibacterial activity gives credence to its use traditionally for the treatment of dysentery, fever and gonorrhoea [5,6]. Its hypoglycemic and hypolipidaemic effect has also been reported [7].

Excessive acute or chronic alcohol consumption poses a serious health hazard and can result into several metabolic disorders in hepatic and extra-

hepatic diseases [8]. Alcohol is a commonly used hepatotoxin in experimental hepatopathy. Although the pathogenesis of alcohol-induced liver disease is not clearly defined, there is evidence that ethanol-induced liver injury is due to oxidative stress that leads to fibrosis and impaired liver functions [9,10]. Alcohol overuse is also characterized by central nervous system (CNS) intoxication symptoms, impaired brain activity, poor motor coordination, and behavioral changes [11]. Excessive alcohol consumption commonly causes hepatic, gastrointestinal, nervous and cardiovascular injuries leading to physiological dysfunctions [12]. Cellular disturbances resulting from excessive alcohol consumption results in increased formation of oxidative stress biomarkers such as malondialdehyde (MDA); reduction in the level of reduced glutathione level and a decrease in the activities of antioxidant enzymes [13,14]. Free radicals and reactive oxygen species (ROS) have been implicated in the oxidative damage of biomolecules and various organs of the body. Studies have shown the crucial role free radicals play in the pathogenesis of many human diseases namely, cardiovascular and pulmonary diseases, some types of cancer, immune/autoimmune diseases, inflammation, diabetes, cataracts and brain dysfunction such as Parkinson and Alzheimer [15]. However, the deleterious effect of free radicals can be corrected by antioxidants – both enzymatic and nonenzymatic. Oxidative stress is known to arise when there is an imbalance between free radical production (especially reactive oxygen species; ROS) and endogenous antioxidant defense system. This shift in balance is associated with

oxidative damage to a wide range of biomolecules including lipids, proteins, and nucleic acids, which may eventually impair normal functions of various tissues and organs [16].

There is an increasing global interest concerning the use of medicinal plants in the prevention and treatment of different pathologies [17,18]. The beneficial effects of plants are attributed to the presence of secondary metabolites such as polyphenols, tannins, terpenoids, alkaloids, flavonoids [19]. Considering the central role played by free radicals in the initiation and progression of many diseases, the use of natural products with antioxidant constituents has been proposed as an effective therapeutic and/or preventive strategy against diseases and the search for potent and cost-effective antioxidants of plant origin has since increased [20]. Many plants have been shown to possess antioxidant potentials [21,22]. This has thus raised interest in the investigation of commonly consumed plants for their phytochemicals with nutritional and chemotherapeutic potentials. Therefore, the need to argument synthetic chemotherapeutic compounds with natural products is the drive for the exploitation of natural products from plants; as they may have little or no side effects yet meeting the nutritional, chemotherapeutic and economic needs [23,24]. Moreover, despite the efforts of pharmaceutical companies in the production of synthetic antibiotics, there yet exists a marked increase in pathogen population exacerbated by multi drug resistant microorganisms. Consequently, there is increased research into phytochemicals for the effective therapeutics combat of this menace. The therapeutic effects of plant-based drugs have been documented to be due to the phytochemicals that constitute the plants [25,26]. These constituents selectively target toxins and pathogens without significant detrimental effect on the human host. This study therefore sought to investigate the ameliorative efficacy of *C. olitorius* leaves against acute ethanol-induced oxidative stress in Wistar rats.

## 2. MATERIALS AND METHODS

### 2.1 Collection and Extraction of Plant Material

Fresh plants of *C. olitorius* were harvested from the Institute of Agricultural Research and Training, Moor Plantation, Ibadan and were

identified by a botanist. The leaves were carefully removed from the stem and washed in running water to remove contaminants. They were air dried at room temperature in an open laboratory space for 14 days and milled into powder using an electronic blender (Moulinex). The extraction was done using soxhlet apparatus and methanol as the solvent according to the method described by Airaodion et al. [7]. About 25 g of the powder was packed into the *thimble* of the soxhlet extractor and 250 mL of methanol was added to a round bottom flask, which was attached to the soxhlet extractor and condenser on a heating mantle. The solvent was heated using the heating mantle and began to evaporate moving through the apparatus to the condenser. The condensate dripped into the reservoir housing the thimble containing the sample. Once the level of the solvent reached the siphon, it poured back into the round bottom flask and the cycle began again. The process was allowed to run for a total of 18 hours. Once the process was completed, the methanol was evaporated in a rotary evaporator at 35°C with a yield of 2.17 g which represents a percentage yield of 8.68%. The crude extract was preserved in the refrigerator until when needed.

### 2.2 Animal Treatment

Twenty adult male Wistar rats with body weight between 120 and 150 g were purchased from the Central Animal House, College of Medicine, University of Ibadan, Nigeria. They were housed in Imrat animal house, Ibadan. They were acclimatized for seven (7) days during which they were fed *ad libitum* with standard feed and drinking water and were housed in clean cages placed in well-ventilated housing conditions (under humid tropical conditions) throughout the experiment. All the animals received humane care according to the criteria outlined in the 'Guide for the Care and Use of Laboratory Animals' prepared by the National Academy of Science and published by the National Institute of Health. They were randomly divided into four groups of five rats each. Animals in groups 1 and 2 were administered normal saline solution while those in groups 3 and 4 were administered crude extract of *C. olitorius* leaves for twenty-one days. The animals were administered the extract and saline solution at a dose of 4 mL per 100 g body weight 12 hourly via oral route of administration. At the end of the treatment, they were fasted overnight and animals in groups 2 and 4 were exposed to a single dose of 70% ethanol at 12 ml/kg body weight to induce oxidative stress. The

dosage of ethanol used in this study has been documented to induce tissue toxicity and oxidative damage in rats [27]. After 12 hours of ethanol administration, the animals were anaesthetized using diethyl ether and were sacrificed. Liver was excised, weighed and homogenized in 50 mmol/L Tris-HCl buffer (pH 7.4) and then centrifuged at  $5000 \times g$  for 15 minutes for biochemical analysis. Supernatants were immediately kept frozen for further analysis.

## 2.3 Biochemical Analyses

### 2.3.1 Determination of hepatic marker enzymes activities

Alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine aminotransferase (ALT) and lactate dehydrogenase (LDH) activities were determined using Randox commercial Enzyme kits according to the method of Reitman and Frankel [28].

### 2.3.2 Determination of oxidative stress biomarkers

Determination of Lipid Peroxidation (LPO), Reduced Glutathione (GSH), Catalase (CAT), Superoxide Dismutase (SOD) and Glutathione peroxidase (GPX) were carried out according to the methods previously described by Airaodion et al. [29].

## 2.4 Statistical Analysis

Results are expressed as mean  $\pm$  standard error of the mean (S.E.M). The levels of homogeneity among the groups were assessed using One-way analysis of variance (ANOVA) followed by Turkey's test. All analyses were done using Graph Pad Prism Software Version 5.00 and p values  $< 0.05$  were considered statistically significant.

## 3. RESULTS

One major finding of this study was that *C. olitorius* leaves indeed unhinged and perturbed the activities of hepatic marker enzymes and oxidative stress biomarkers in the animals used. These perturbations were reflected as up/down regulation of the activities of these enzymes as shown in Tables 1 and 2.

## 4. DISCUSSION

Green leafy vegetables and fruits have been reported to have some health benefits [21,22]. Consequently, *C. olitorius* leaf is a green leafy vegetable popularly used as food and in traditional medicine for the management of some diseases [30]. However, there is dearth of information on the possible mechanisms of action by which these vegetables exert

**Table 1. Effect of *C. olitorius* on hepatic marker enzymes of experimental rats**

Hepatic marker enzymes	Control	70% Ethanol only	<i>C. olitorius</i> extract only	<i>C. olitorius</i> Extract + 70% Ethanol
AST (IU/L)	110.27 $\pm$ 3.22 <sup>ac</sup>	132.84 $\pm$ 5.78 <sup>b</sup>	101.22 $\pm$ 3.53 <sup>a</sup>	109.06 $\pm$ 3.83 <sup>c</sup>
ALT (IU/L)	46.85 $\pm$ 2.17 <sup>a</sup>	56.49 $\pm$ 3.64 <sup>b</sup>	44.45 $\pm$ 1.32 <sup>a</sup>	48.13 $\pm$ 2.92 <sup>c</sup>
ALP (IU/L)	15.60 $\pm$ 2.57 <sup>ac</sup>	22.79 $\pm$ 1.82 <sup>b</sup>	12.84 $\pm$ 1.23 <sup>a</sup>	17.41 $\pm$ 2.62 <sup>c</sup>
LDH (IU/L)	184.40 $\pm$ 9.13 <sup>ac</sup>	206.55 $\pm$ 10.10 <sup>b</sup>	197.18 $\pm$ 11.29 <sup>a</sup>	191.83 $\pm$ 14.05 <sup>c</sup>

Values are presented as Mean $\pm$ S.E.M, where n = 5. Values with different superscript along the same row are significantly different at p<0.05

Legend: AST = Aspartate Amino Transferase, ALT = Alanine Amino Transferase, ALP = Alkaline Phosphatase, LDH = Lactate Dehydrogenase

**Table 2. Effect of *C. olitorius* on oxidative stress biomarkers of experimental rats**

Oxidative stress biomarkers	Control	70% Ethanol only	<i>C. olitorius</i> extract only	<i>C. olitorius</i> extract + 70% ethanol
LPO (nmol MDA/mg protein)	11.76 $\pm$ 1.08 <sup>a</sup>	18.91 $\pm$ 2.27 <sup>b</sup>	10.02 $\pm$ 1.38 <sup>a</sup>	12.53 $\pm$ 1.27 <sup>a</sup>
GSH (mg/mL)	4.68 $\pm$ 0.10 <sup>a</sup>	3.06 $\pm$ 0.78 <sup>b</sup>	5.06 $\pm$ 0.57 <sup>a</sup>	4.28 $\pm$ 0.82 <sup>a</sup>
CAT (Mm H <sub>2</sub> O <sub>2</sub> /mg protein)	13.04 $\pm$ 1.04 <sup>a</sup>	22.00 $\pm$ 2.29 <sup>b</sup>	11.56 $\pm$ 2.43 <sup>a</sup>	17.34 $\pm$ 1.85 <sup>c</sup>
SOD (U/mg protein)	8.62 $\pm$ 1.21 <sup>a</sup>	13.93 $\pm$ 2.23 <sup>b</sup>	8.91 $\pm$ 1.28 <sup>a</sup>	9.63 $\pm$ 1.22 <sup>a</sup>
GPX (U/mg protein)	6.08 $\pm$ 0.16 <sup>a</sup>	8.73 $\pm$ 1.17 <sup>b</sup>	5.24 $\pm$ 0.84 <sup>a</sup>	6.11 $\pm$ 1.23 <sup>a</sup>

Values are presented as Mean $\pm$ S.E.M, where n = 5. Values with different superscript along the same row are significantly different at p<0.05

Legend: LPO = lipid peroxidation, GSH = Glutathione, CAT = Catalase, SOD = Superoxide Dismutase, GPX = Glutathione Peroxidase

their health benefits. Iseri et al. [31] reported that *C. olitorius* (jute) extract induced cytotoxicity and genotoxicity on human multiple myeloma cells (ARH-77). Acute and chronic alcohol exposures have been documented to increase the generation of reactive oxygen species (ROS). Many investigations have revealed a decreased level of antioxidants and increased production of free radicals in animals and humans following excessive ethanol exposure [32,33]. The protective effect of antioxidants against alcohol-induced liver injury in many studies further supports the involvement of oxidative stress [30].

Evaluation of liver function is very important when analyzing toxicity of drugs and plant extracts because of its relevance for the survival of the organism [34]. High levels of alkaline phosphatase (ALP), aspartate transaminase (AST), alanine transaminase (ALT) and lactate dehydrogenase (LDH) are an indicator of hepatotoxicity or liver diseases [35]. Studies on the alterations of these enzymes might reflect the metabolic abnormalities and cellular injuries in some organs. The liver and kidney have extremely important function in detoxification and excretion of metabolic wastes and xenobiotics [36]. Exposure to toxic chemicals causes alterations in some tissue enzyme activities [37,38]. Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) are distributed extensively in several different organs and have important roles in carbohydrate and amino acid metabolic pathways and their activities is established to change under several physiological and pathological circumstances [39].

In this study, the activities of AST, ALT, ALP and LDH were not significantly different upon treatment cell membrane of ethanol-induced animals without pretreatment. This effect was remedied by pretreatment with *C. olitorius* extract.

The elevation in the activities of markers such as ALT, AST and LDH in the liver tissue of animals without pretreatment with *C. olitorius* might be due to cellular necrosis of hepatocytes, which causes increase in the permeability of the cell. Lactate dehydrogenase (LDH) is an index of cell damage including hepatotoxicity and the endothelial disruption in blood vessel. The significant increase observed in the activity of LDH might be suggestive of the beginning of cytolysis, which is a possible indication of membrane damage including the endothelial

membranes of blood vessels. This disruption of endothelial membrane, directly or indirectly, as reported earlier, includes the generation of reactive oxygen species in endothelial cells [46]. Free radicals attack unsaturated fatty acids in the membranes resulting in membrane lipid peroxidation which decreases membrane fluidity, leakage of enzyme and loss of receptor activity as well as damage membrane proteins leading to cell inactivation [29]. As lipid peroxidation progressively increase, antioxidant defense system decrease equivalently resulting in oxidative stress [47]. This suggests that the administration of ethanol might have weakened the liver membrane of the rats with subsequent penetration and elevation of the hepatic biomarker enzymes.

Alcohol metabolism results in oxidative and nitrosative stress via elevation of NADH/NAD<sup>+</sup> redox ratios, induction of nitric oxide synthase (NOS) and NADPH/xanthine oxidase [48,49]. Lipid peroxidation, a primary mechanism of cell membrane destruction and cell damage is a common feature of both acute and chronic alcohol consumption [50]. The presence of a high concentration of oxidizable fatty acids and iron in liver significantly contributes to ROS production. A rise in lipid peroxidation level is only identified if there is oxidative damage due to the increase in free radical generation. Generally under normal conditions, the animals tend to maintain a balance between generation and neutralization of ROS in the tissues. But, when the organisms are subjected to xenobiotic stress, the rate of production of ROS including O<sub>2</sub><sup>-</sup>, H<sub>2</sub>O<sub>2</sub>, OH<sup>-</sup>, ROO<sup>-</sup>, exceeds their scavenging capacities. All the organisms have their own cellular antioxidant defense system composed of both enzymatic and non-enzymatic components. Enzymatic antioxidant pathway consists of SOD, CAT and GPX. Superoxide anion O<sub>2</sub><sup>-</sup> is dismutated by SOD to H<sub>2</sub>O<sub>2</sub>, which is reduced to water and molecular oxygen by CAT or is neutralized by GPX, which catalyzes the reduction of H<sub>2</sub>O<sub>2</sub> to water and organic peroxide to alcohols using GSH as a source of reducing equivalent. Glutathione reductase (GR) regenerates GSH from oxidized glutathione (GSSG), which is a scavenger of ROS as well as a substrate for other enzymes. Glutathione S-transferase (GST) conjugates xenobiotics with GSH for exclusion.

In this study, acute ethanol exposure significantly elevated the malondialdehyde (MDA) levels in the liver indicating enhanced peroxidation and breakdown of the antioxidant defense

mechanisms. Decomposition products of lipid hydroperoxide such as malonaldehyde and 4-hydroxynonenal, can cause chaotic cross-linkage with proteins and nucleic acids, which plays an important role in the process of carcinogenesis. In this investigation, hepatic lipid peroxidation (LPO) activities show significant increase due to ethanol intoxication. Furthermore, extensive damage to tissues in a free radical mediated LPO results in membrane damage and subsequently decreases the membrane fluid content. *C. olitorius* pretreatment significantly reversed these alterations causing a significant decrease in MDA levels, suggesting its protective effects against ethanolic-induced oxidative damage. This is consistent with the study of Oyenihini et al. [46] who reported the hepato- and neuro-protective effects of watermelon juice on acute ethanol-induced oxidative stress in rats. It is also in agreement with the report of Airaodion et al. [29] who studied the hepatoprotective effect of *Parkia biglobosa* on acute ethanol-induced oxidative stress in Wistar rats.

Glutathione (GSH) is a tripeptide (L- $\alpha$ -glutamylcysteinol glycine) which is highly abundant in all cell compartments and it is the major soluble antioxidant. Glutathione directly quenches ROS such as lipid peroxides, and also plays a major role in xenobiotic metabolism. Glutathione detoxifies hydrogen peroxide and lipid peroxide by donating electron to hydrogen peroxide to reduce it to water and oxygen protecting macromolecules such as lipids from oxidation. In this study, the decrease in the reduced glutathione level in animals treated with ethanol only is connected with ethanol-induced oxidative stress and direct conjugation of GSH with acetaldehyde and other reactive intermediates of alcohol oxidation. This result is in agreement with the finding of Pinto et al. [31] who reported that acute ethanol treatment caused reduction in the glutathione levels in different tissues. The significant increase ( $P < 0.05$ ) in the glutathione levels in the liver of *C. olitorius*-treated rats prior to ethanol-administration may be due to the direct ROS—scavenging effect of *C. olitorius* or an increase in GSH synthesis. This is consistent with the report of Airaodion et al. [29] who studied the hepatoprotective effect of *Parkia biglobosa* on acute ethanol-induced oxidative stress in Wistar rats.

Catalase (CAT) contributes to ethanol oxidation, by oxidizing a small amount of ethanol in the presence of a hydrogen peroxide ( $H_2O_2$ )

generating system to form acetaldehyde [51]. In this study, a significant increase was observed in the activity of catalase in control animals and those treated with *C. olitorius* extract only when compared with ethanol-induced animals with *C. olitorius* extract pretreatment. This contradicts the findings of Airaodion et al. [29] who reported a nonsignificant difference when animals were treated with *Parkia biglobosa*. The activity of catalase in animals pretreated with *C. olitorius* prior to ethanol induction was significantly reduced when compared with those without *C. olitorius* pretreatment. This might be that ethanol-induced oxidative stress generated elevated ROS in the liver which CAT tend to combat, thereby increasing its activity. *C. olitorius* was able to reduce the ROS generation with subsequent decrease in CAT activity due to its high phytochemical content as reported by Orieki et al. [52]. Increased CAT activity in this study following acute ethanol exposure suggests elevated ethanol oxidation and formation of oxidizing product-acetaldehyde. This is in agreement with the study of Airaodion et al. [29] and Oyenihini et al. [46] who reported a significantly higher CAT activity after ethanol treatment.

Superoxide dismutase (SOD) plays an important role in reducing the effect of free radicals attack, and SOD is the only enzymatic system quenching  $O_2^-$  to oxygen and  $H_2O_2$  and plays a significant role against oxidative stress [53]. These radicals have been reported to be deleterious to polyunsaturated fatty acids and proteins [54]. In this study, no significant difference was observed in the activity of SOD in control animals and those treated with *C. olitorius* extract only when compared with ethanol-induced animals with *C. olitorius* extract pretreatment. The activity of SOD in animals pretreated with *C. olitorius* prior to ethanol induction was significantly reduced when compared with those without *C. olitorius* pretreatment. This might be that ethanol-induced oxidative stress generated elevated ROS in the liver which SOD tend to combat thereby increasing its activity. *C. olitorius* was able to reduce the ROS generation with subsequent decrease in SOD activity due to its high phytochemical content as reported by Orieki et al [52]. The increased activity of SOD observed in ethanol induced animals contradicts the study of Halliwell and Gutterberidge [55] who reported that SOD activity was considerably reduced during ethanol intoxication.

Glutathione peroxidase (GPX) is another enzymic antioxidant that acts as a defense against oxidative stress. In this study, no significant difference was observed in the activity of GPX in control animals and those treated with *C. olitorius* extract only when compared with ethanol-induced animals with *C. olitorius* extract pretreatment. The activity of GPX in animals pretreated with *C. olitorius* prior to ethanol induction was significantly reduced when compared with those without *C. olitorius* pretreatment. This might be that ethanol-induced oxidative stress generated elevated ROS in the liver which GPX tend to combat thereby increasing its activity. *C. olitorius* was able to reduce the ROS generation with subsequent decrease in GPX activity due to its high phytochemical content as reported by Orieke et al. [54]. The increased activity of GPX observed in ethanol induced animals contradicts the studies of Airaodion et al. [29] who observed no significant difference in the activity of GPX in the study of hepatoprotective effect of *Parkia biglobosa* on acute ethanol-induced oxidative stress in Wistar rats and that of Yang et al. [56] who also observed no significant difference in GPX activities in rats hepatocyte exposed to varying concentrations of ethanol at an incubation time of 12 hours. The toxicity of ethanol is related to the product of its metabolic oxidation. Acetaldehyde and acetate, produced from the oxidative metabolism of alcohol are capable of forming adducts with cellular macromolecules, causing oxidative damage and affecting metabolic processes [51,57]. Catalase and glutathione peroxidase further detoxify H<sub>2</sub>O<sub>2</sub> into H<sub>2</sub>O and O<sub>2</sub> [54]. Thus, SOD, catalase and GPX function mutually as enzymatic antioxidative defense mechanism to counter the deleterious effect of ROS.

## 5. CONCLUSIONS

The results of this study indicated that ethanol induced oxidative stress as shown in the perturbation of the biomarkers. *C. olitorius* leaves was able to remedy this effect by regulating the oxidative stress biomarkers, thus possesses ameliorative efficacy against ethanol-induced oxidative stress and can protect the body against free radicals arising from oxidative stress. Regular consumption of this vegetable is hereby recommended.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Adedosu OT, Akanni OE, Afolabi OK, Adedeji AL. Effects of *Corchorus olitorius* Extract on Certain Antioxidants and Biochemical Indices in Sodium Arsenite Exposed Rats. American Journal of Phytomedicine and Clinical Therapeutics. 2015;3(03):245-256
2. Onyeka EU, Nwambekwe IO. Phytochemical profile of some green leafy vegetables in South East, Nigeria. Nigeria Institute of Food Science and Technology; 2007.
3. Ujah OF, Ipav SS, Ayaebene CS, Ujah C. Phytochemistry and hepato-protective effect of ethanolic leaf extract of *Corchorus olitorius* on carbon tetrachloride induced toxicity. European Journal of Medicinal Plants. 2014;4(8):882-892.
4. Nemba RM, Emadak A, Mouzong GC, Nemba CE. Qualitative and quantitative assesment of mineral elements in the leaves of *Corchorus fascicularis* and *Corchorus olitorius* harvested in Cameroon. J Curr Chem Pharm Sci. 2012;2(1):17-23.
5. Batran RA, Bayaty F, Abdulla MA, Obaidi MM, Hajrezae IM, Hassandarvish P, Fouad M, Golbabapour S, Talaee S. Gastro-protective effect of *Corchorus olitorius* leaf extract against ethanol-induced gastric mucosal hemorrhagic lesion in rats. J. gastroenterol hepatol. 2013;28(8):1321-1329.
6. Egua MO, Etuk EU, Bello SO, Hassan SW. Anti-diabetic activity of ethanolic seed extract of *Corchorus olitorius*. International Journal of Sciences: Basic and Applied Research. 2013;12(1):8-21.
7. Airaodion AI, Ogbuagu EO, Ogbuagu U, Awosanya OO, Airaodion EO. Effect of methanolic extract of *Corchorus olitorius* leaves on hypoglycemic and hypolipidaemic activities in albino rats. Asian Plant Research Journal. 2019;2(6):1-13.
8. Lieber CS. Ethnic and gender differences in ethanol metabolism, Alcohol. Clin. Exp. Res. 2000;24 (4):417-418.
9. Wu D, Cederbaum AI. Alcohol, oxidative stress, and free radical damage, Alcohol Res. Health. 2003;27:277-284.
10. Ronis MJ, Hakkak R, Korourian S, Albano E, Yoon S, Ingelman-Sundberg M, Lindros KO, Badger TM. Alcoholic liver disease in rats fed ethanol as part of oral or

- intra-gastric low-carbohydrate liquid diets. *Exp. Biol. Med.* 2004;229(4):351–360.
11. Gohlke JM, Griffith WC, Faustman EM. Computational models of ethanol-induced neurodevelopmental toxicity across species: Implications for risk assessment, *Birth Defect Res. B: Dev. Reprod. Toxicol.* 2008;83(1):1–11.
  12. Lieber CS. Hepatic and other medical disorders of alcoholism: From pathogenesis to treatment, *J. Stud. Alcohol.* 1998;59(1): 9–25.
  13. Nadro MS, Arungbemi RM, Dahiru D. Evaluation of *Moringa oleifera* leaf extract on alcohol-induced hepatotoxicity, *Trop. J. Pharm. Res.* 2007;5(1):539–544.
  14. Das SK, Vasudevan DM. Alcohol-induced oxidative stress. *Life Sci.* 2007;81(3):177–187.
  15. Rahman T, Hosen I, Islam MT, Shekhar HU. Oxidative stress and human health. *Adv Biosci Biotechnol.* 2012;3:997–1019.
  16. Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010;4:118–126.
  17. Gorinstein S, Yamamoto K, Katrich E, Leontowicz H, Lojek A, Leontowicz M, Trakhtenberg S. Antioxidative properties of Jaffa sweets and grapefruit and their influence on lipid metabolism and plasma antioxidative potential in rats, *Biosci. Biotechnol. Biochem.* 2003;67(4): 907–910.
  18. Ramana KV, Singhal SS, Reddy AB. Therapeutic potential of natural pharmacological agents in the treatment of human diseases, *Bio Med Res. Int.* 2014;1–4.
  19. Chikezie PC, Chiedozie O, Ibegbulem FN. Bioactive principles from medicinal plants. *Res. J. Phytochem.* 2015;9(3):88–115.
  20. Morales-González JA. Oxidative stress and chronic degenerative diseases-A role for antioxidants. In *Tech, Rijeka, Croatia.* 2013;500.
  21. Airaodion AI, Olatoyinbo PO, Ogbuagu U, Ogbuagu EO, Akinmolayan JD, Adekale OA, Awosanya OO, Agunbiade AP, Oloruntoba AP, Obajimi OO, Adeniji AR, Airaodion EO. Comparative assessment of phytochemical content and antioxidant potential of *Azadirachta indica* and *Parquetina nigrescens* leaves. *Asian Plant Research Journal.* 2019;2(3):1-14.
  22. Airaodion AI, Ibrahim AH, Ogbuagu U, Ogbuagu EO, Awosanya OO. Akinmolayan JD, Njoku OC, Obajimi OO, Adeniji AR, Adekale OA. Evaluation of phytochemical content and antioxidant potential of *Ocimum gratissimum* and *Telfairia occidentalis* leaves. *Asian Journal of Research in Medical and Pharmaceutical Sciences.* 2019;7(1):1-11.
  23. Airaodion AI, Obajimi OO, Ezebuoro CN, Ogbuagu U, Agunbiade AP, Oloruntoba AP, Akinmolayan JD, Adeniji AR, Airaodion EO. Prophylactic efficacy of aqueous extract of *Curcuma longa* leaf against indomethacin-induced ulcer. *International Journal of Research.* 2019;6(1):87-91.
  24. Airaodion AI, Olayeri IM, Ewa AO, Ogbuagu EO, Ogbuagu U, Akinmolayan JD, Agunbiade AP, Oloruntoba AP, Airaodion EO, Adeniji AR, Obajimi OO, Awosanya OO. Evaluation of *Moringa oleifera* leaf potential in the prevention of peptic ulcer in wistar rats. *International Journal of Research.* 2019;6(2):579-584.
  25. Airaodion AI, Ogbuagu U, Ogbuagu EO, Airaodion EO, Agunbiade AP, Oloruntoba AP, Mokelu IP, Ekeh SC. Investigation of aqueous extract of *Zingiber officinale* root potential in the prevention of peptic ulcer in albino rats. *International Journal of Research and Innovation in Applied Science.* 2019;4(2):64-67.
  26. Airaodion AI, Adekale OA, Airaodion EO, Ogbuagu EO, Ogbuagu U, Osemwowa EU. Efficacy of combined crude extract of *Curcuma longa* and *Moringa oleifera* leaf in the prevention of peptic ulcer in albino rats. *Asian Journal of Research in Medical and Pharmaceutical Sciences.* 2019;7(2):1-9.
  27. Han Y, Xu Q, Hu JN, Han XY, Li W, Zhao LC. Maltol, a food flavoring agent, attenuates acute alcohol-induced oxidative damage in mice, *Nutrients.* 2015;7(1):682–696.
  28. Reitman S, Frankel S. A colorimetric method for determination of serum glutamate oxaloacetate and glutamic pyruvate transaminase. *American Journal of Clinical Pathology.* 1957;28:56-58.
  29. Airaodion AI, Ogbuagu EO, Ogbuagu U, Adeniji AR, Agunbiade AP, Airaodion EO. Hepatoprotective effect of *Parkia biglobosa* on acute ethanol-induced oxidative stress in wistar rats. *International Research Journal of Gastroenterology and Hepatology.* 2019;2(1):1-11.
  30. Saravanan R, Pari L. Antihyperlipidemic and antiperoxidative effect of diasulin, a polyherbal formulation in alloxan induced



- hyperglycemic rats. BMC Complement. Alternative Med. 2005;5:14–34.
31. Iseri OD, Yurtcu E, Sahin FI, Mehmet H. *Corchorus olitorius* (jute) extract induced cytotoxicity and genotoxicity on human multiple myeloma cells (ARH-77). Pharmaceutical Biology. 2013;51(6):766-770.
  32. Airaodion AI, Airaodion EO, Ewa O, Ogbuagu EO, Ogbuagu U. Nutritional and anti-nutritional evaluation of garri processed by traditional and instant mechanical methods. Asian Food Science Journal. 2019;9(4):1-13.
  33. Airaodion AI, Airaodion EO, Ogbuagu EO, Ogbuagu U, Osemwowa EU. Effect of oral intake of African locust bean on fasting blood sugar and lipid profile of albino rats. Asian Journal of Research in Biochemistry. 2019;4(4):1-9.
  34. Owoade AO, Adetutu A, Airaodion AI, Ogundipe OO. Toxicological assessment of the methanolic leaf extract of *Bridelia ferrugelia*. The Journal of Phytopharmacology. 2018;7(5):419-424.
  35. Brautbar N, Williams JI. Industrial solvents and liver toxicity: Risk assessment, risk factors and Mechanisms. International J of Hygiene and Environmental Health. 2002; 25(6):497-492.
  36. Kaneko JJ, Harvey JW, Bruss ML. Clinical biochemistry of domestic animals. 5<sup>th</sup> Ed, Academic Press, London. 1999;829-44.
  37. Jarrar BM, Mahmoud ZN. Histochemical demonstration of changes in the activity of hepatic phosphatases induced by experimental lead poisoning in male white rats (*Rattus norvegicus*). Toxicol Ind Health. 1999;15:1–9.
  38. Gholipour-Kanani H, Shahsavani D, Baghishani H. Effect of exposure to sublethal levels of potassium cyanide on serum and tissue enzymes in roach fish (*Rutilus rutilus*). Online J Vet Res. 2013; 17(5):245-55.
  39. Al-Ghanim KA. Effect of cypermethrin toxicity on enzyme activities in the freshwater fish *Cyprinus carpio*. African Journal of Biotechnology. 2014;13(10): 1169–1173.
  40. Towle HC, Kaytor EN, Shih HM. Regulation of the expression of lipogenic enzymes by carbohydrates. Annual Review of Nutrition. 1997;17:405-433
  41. Zubay G, Schwartz D, Beckwith J. Mechanism of activation of catabolite-sensitive genes: a positive control system. Proceedings of the National Academy of Sciences. 1970;66(1):104-110.
  42. Todar K. Regulation and control of metabolism in bacteria: In: Todar's Online Textbook of Bacteriology. 2008;1-5.
  43. Johnson PJ. The assessment of hepatic function and investigation of jaundice. In: Marshall WJ, Bangert SK, editors. Clinical Biochemistry – Metabolic and Clinical Aspects. Churchill Livingstone, New York. 1995;217-36.
  44. Wright PJ, Plummer DT. The use of urinary enzyme measurement to detect renal damage caused by Nephrotoxic compounds. Biochem. Pharmacol. 1974;23:98-112.
  45. Akanji MA, Olagoke OA, Oloyede OB. Effect of chronic consumption of meta-bisulphite on the integrity of the rat kidney cellular system. Toxicology. 1993;81:173-179.
  46. Oyenihni OR, Afolabi BA, Oyenihni AB, Ogunmokin OJ, Oguntibeju OO. Hepato- and neuro-protective effects of watermelon juice on acute ethanol-induced oxidative stress in rats. Toxicology Reports. 2016;3: 288–294.
  47. Tiwari AK, Rao JM. Diabetes mellitus and multiple therapeutic approaches of phytochemicals: Present status and future prospects. Curr. Sci. 2002;83:30-38.
  48. Manzo-Avalos S, Saavedra-Molina A. Cellular and mitochondrial effects of alcohol consumption, Int. J. Environ. Res. Public Health. 2010;7(12):4281–4304.
  49. Sathivelu J, Senapathy GJ, Devaraj R, Namasivayam N. Hepatoprotective effect of chrysin on prooxidant-antioxidant status during ethanol-induced toxicity in female albino rats, J. Pharm. Pharmacol. 2009; 61(6):809–817.
  50. Ramezani A, Goudarzi I, Lashkarboluki T, Ghorbanian MT, Abrari K, Salmani ME. Role of oxidative stress in ethanol-induced neurotoxicity in the developing cerebellum, Iran. J. Basic Med. Sci. 2012;15(4):965.
  51. Agarwal DP. Genetic polymorphisms of alcohol metabolizing enzymes, Pathol. Biol. 2001;49(9):703–709.
  52. Orieko D, Ohaeri OC, Ijeh II, Ijioma SN. Identification of phytochemicals and acute toxicity evaluation of *Corchorus olitorius* leaf extract. European Journal of Medicinal Plants. 2018;23(1)1-16.
  53. Oh SI, Kim CI, Chun HJ, Park SC. Chronic ethanol consumption affects glutathione status in rat liver, J. Nutr. 1998;128(4): 758–763.

54. Murray RK, Granner DK, Mayes PO, Rodwell VW. Harper's illustrated biochemistry. 26<sup>th</sup> Edn., Appleton and Lange Medical Publication/McGraw Hill, USA. 2003;693. [ISBN-13: 9780071389013]
55. Halliwell B, Gutterberidge JMC. Free radicals in biology and medicine. 4<sup>th</sup> ed. Oxford, UK: Oxford University; 2006.
56. Yang SS, Huang CC, Chen JR, Chiu CL, Shieh MJ, Lin SJ, Yang SC. Effects of ethanol on antioxidant capacity in isolated rat hepatocytes, World J. Gastroenterol. 2005;11(46):7272.
57. Shirpoor A, Minassian S, Salami S, Khadem-Ansari MH, Yeghiazaryan M. Alpha-lipoic acid decreases DNA damage and oxidative stress induced by alcohol in the developing hippocampus and cerebellum of rat, Cell. Physiol. Biochem. 2008;22:(5-6):769-776.

© 2019 Airaodion et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/49753>