



An Overview on Fungal Diseases in Angiospermic Plants

Arti ^{a++} and Simerjit Kaur ^{a#*}

^a *Department of Life Sciences, Rayat Bahra University, Mohali, Punjab, India.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/APRJ/2023/v11i2207

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97839>

Review Article

Received: 04/02/2023

Accepted: 06/04/2023

Published: 12/04/2023

ABSTRACT

The main objective of this review study is to examine various fungal diseases on angiospermic plants which lower plant productivity and worsen the financial situation of flowers. Angiospermic flora are the most useful and marketable plants, employed for the majority of purposes, contributing to a better environment and spreading positive energy, but these flowering plants are presently damaged by unfavourable environmental conditions. The occurrence of fungal infection suppresses the beauty, growth, production and commercial values of flowers and also affect the living beings by increasing air pollution due to the infected plants. Fungicides, chemicals and molecular techniques are already in used to enhances the yield of crops and growth of plants, but still more research needs to be conducted to find the way for increasing the yield and preventing the plants more.

Keywords: Fungal diseases; flowering plants; pathogens; fungicides; pathological study; plant-pathogen interactions.

⁺⁺ *Research Scholar;*

[#] *Head;*

^{*}*Corresponding author: E-mail: dr.simar@rayatbahrauniversity.edu.in, dr.simer07@yahoo.com;*

1. INTRODUCTION

Angiosperms plants are classified as bearing seeds inside the ovary and producing fruit. They are true vascular plants since angiospermic plants have roots, stems, and leaves that are actually separate from one another. The angiosperms are diverse organisms. These plants are widespread around the world. Only angiospermic plants undergo double fertilisation or triple fusion. The globe over, flowers are utilised for a wide variety of functions and are a symbol of the natural beauty of the earth. The fungi are non-vascular, non-motile, heterotrophic and eukaryotic organisms. Fungi produces the number of different spores that causes the disease 's symptoms in the plants. It can cause the necrosis which is associated with some other symptoms like: leaf spot, canker, dieback, blight scab, anthracnose, mildew damping-off etc, the fungal symptoms can inhibit the growth of the plants or induce the abnormal or infected plants.

2. FUNGAL DISEASES IN ANGIOSPERMS

Initial leaf symptoms of the illness were a yellowish staining, which was followed by the emergence of variously sized grayish-white lesions on Tulip plant leaves. Blossom blight, a disease *Phytophthora cactorum* historically caused in Tulip plants, was discovered before the diseases mentioned above. *Pythium dissotocum*, *P. afertile*, *P. sylvaticum*, and other *Pythium* species were identified as the causal organisms for the current diseases [1]. Strawberry is a very delicious and popular fruit in the whole world and used for different purposes at different places, but certainly the productivity of strawberry fruit has been declined due to the infection caused by the fungi which causes the various diseases such as Botrytis gray mold rot (*Botrytis cinerea*), leafspot (*Ramularia tulasnei*), anthracnose (*Colletotrichum fragariae*), powdery mildew (*Sphaerotheca macularis*), crown rot & leather fruit rot (*Phytophthora cactorum*) and black root rot (*Pythium* spp., *Fusarium* spp.), etc and reduced the quality and yielding of the fruit. But now the several fungicides have been produced to control the infection and protect the plants [2].

Tagetes erecta is a most growing flowering plant in the world which are used for the different purposes and every occasion especially in India, but the yielding of the flowers has been less due to the fungal infection in the plant's parts mainly on flowers & leaves. The leaf spot and flower

blight diseases has been found due to their causal pathogens as *Alternaria tagetica*, which reduced the productivity of flowers and inhibit the growth as well [3]. *Hibiscus* plants are very useful for many purposes which belongs to Malvaceae family and these flowering plants are found in all over the world, but now the various diseases have been produced time to time due to the fungal causal organisms that inhibit the growth and development of the whole plant and shows the reduction in the quality and quantity as well [4].

To enhance the growth of Hibiscus plants, several cultivars were used, which affects the organisms and stop to produce more diseases in plants. The Root knot nematodes causes the growth suppression, poor yielding and decreases the harvesting efficiency of *Hibiscus* crops and the infection further increases in the plants due to the presence of causal fungus in the soil that produces more diseases in the plants [5]. The very serious diseases caused by the *Botrytis cinerea* in the many crops and mainly the development of this pathogen has been starts after yielding because of the weather conditions which are favourable for this organism and the gray mold disease were produced in the crops which affects the growth of the production and quality as well. But to prevent the crop from this disease, various fungicides was used which kill the organisms and helps to enhance the yielding of the crops [6]. *Silene latifolia* is a biennial or short-lived perennial plant which have its very high economic rates but due to some environmental factors the Anther smut disease has been appeared on the flower due to its causal pathogen *Microbotryum violaceum* and the genetic variations has been reported in this plant. The flowering plants was protected by the chemical treatment and molecular techniques [7].

The various diseases were found in the seedlings of *Hibiscus* plants due to the soil borne fungi, which deduce the growth, good quality of fibres and appearance of flowering plants, but now various chemical treatments with fungicides were used to stop this infection and to kill the identified organisms such as *Rhizoctonia solani*, *Pythium* spp., & *Fusarium* species and these chemicals also enhance the growth of the plants [8]. Every plant has their medicinal properties inside it and the extracts of the plant's material provide protection from the external causal organisms. In the crop of *Capsicum annum*, the number of fungal infections has been reported due to some environmental conditions and lack

of essential requirements in the plants. The main diseases of *Capsicum* are Anthracnose which is caused by the two main fungal species like as *Colletotrichum capsici* and *Gloeosporium piperatum*. The both fungi produced the conidial spores on the plants and mycelial would be grown on the plant's parts. In the recent research, the leaf extracts and some fungicides were used to inhibit the growth of conidia on the plants and prevent the plants from the further infections [9].

The fungal contaminants were detected in the most of angiosperms with the help of PCR primers technique. From the *Dahlia* (Asteraceae) flowering plant, a number of DNA samples were extracted to identify the fungal contamination in Mexico. A simple screening test were used for detecting the fungal infection in angiosperm that reduce the possibility of gathered damaged leaves. More techniques were also used for the identification of fungal infection in angiosperm's DNA [10]. The most effective chemical treatment was developed as *Serratia plymuthica* HRO-C48 to control the infection of diseases Verticillium wilt caused by *Verticillium dahlia* and *Phytophthora* root rot caused by *Phytophthora cactorum*. This biological product was used for the Strawberry plants for stop the growth of the pathogens and kill them and this was so effective which were reported in the nursery, green houses and fields of Germany [11].

Many surveys were conducted in USA to investigate the fungal pathogens which was associated with *Eichhornia crassipes* in rivers and lakes. The outcome of the surveys showed that the different fungal organisms has been identified from this hydrotropic plant which causes the distinct diseases such as leaf lesions caused by *Colletotrichum*, *Stagonospora*, *Myrothecium*; colonising leaf tissue caused by *Mycosphaerella*, *Didymella*, etc. [12]. The several kinds of spores were reported on the fruiting and flowering plants in Taiwan and these different types of spores causes the brown root rot diseases in the plants due to the inoculation of fungal organism *Phellinus noxius* because of the bad environmental conditions. These diseases affect the plants and inhibit the growth as well, but the various chemical treatments have been used to deduce the infection and protect the plants [13].

Catharanthus roseus is a very popular flowering plant as well as medicinal plant, which is found in all over the world. The several alkaloids are

obtained from the extract of *Vinca* which are used for the anticancerous purposes such as vincristine, vindesine, vinorelbine, vinblastine, and the most novel alkaloid is vinflunine. This is a host specific plant of endophytic fungi and the number of fungi were isolated from the different parts and their tissues of *Vinca* plant from two separate regions in North India. The most common fungal pathogens are *Alternaria*, *Cladosporium*, *Aspergillus*, etc. Here, fungi also play a positive role as mutual symbionts by producing natural bioactive compounds that helps to prepared the medicines and cure the several diseases [14]. Previously in 1987, *Plasmopara halstedii* pathogen were infect *Helianthus annuus* plants and caused a harmful downy mildew disease in the region of USA and Europe. Another outbreak of downy mildew disease also caused in *Helianthus annuus* plants by *Plasmopara halstedii* in the commercial field of Thrace, part of Marmara (Turkey) during spring season of 2007 & 2008. This disease was commonly produced in the raining season with low temperature range and decreased the yield of this flowering plants [15].

In the commercial nursery of Poland, the various symptoms of growth abnormalities were seen as shoot proliferation, leaf resetting and malformation in the 3year old Ash tree plants. These symptoms were identical to another ash yellow disease connected with the group of *Phytoplasma* like; *Candidatus Phytoplasma fraxini* in USA. These pathogens have been commonly found in the ornamental crops as *Magnolias*, *Roses* ash leaf maple tree that related with the growth abnormalities and this is the first data on the appearance of *Candida Phytoplasma asteris* in Poland [16]. *Tithonia diversifolia* is the maximum producing flowering plant in the world and this is present throughout the year shows their perennial nature. At presently, the number of plants were identified as infected by the occurrence of powdery mildew diseases from the region of Meghalaya in June-July 2007. The pathogenicity has been confirmed when the formation of different kinds of spores were reported on the plants [17]. The disease on leaf was observed from the rose plants by using the molecular techniques as PCR, BLAST analysis during June 2008 in Gorakhpur (India) and the leaves of the *Rosa* plant become yellowish and the growth has been stopped of the plants due to the fungal infection of *Candidatus asteris* that inhibit the production and induce the abnormal parts of the plants. This fungal pathogen causes the various diseases in

other plants as well like Sesame, Desert rose (*Adenium obesum*) along with the *Rosa alba* [18].

The extract of these three plants (*Lawsonia inermis*, *Punica granatum*, *Hibiscus sabdariffa*) were used to control the fungal, bacterial contamination in flowering plants. The material is used in their natural form for herbal medicines or purified form in pharmaceutical industry. These 3 different extracts were used to check their ability to inhibit the growth of fungal and bacterial pathogens which are becoming a threat to living beings [19]. *Jasminum sambac* is the most growing flowering plant because of its attractive fragrance and appearance. Nowadays the yielding of various flowers was less due to the occurrence of infection by pathogens as *Colletotrichum truncatum*, *C. siamense*, which causes the diseases leaf and fruit anthracnose in the plants. These diseases were identified by using the molecular phylogeny techniques and the techniques also helps to inhibit the further growth of the pathogens in the plants [20].

Chrysanthemum plants have very elegant flowers of golden colour which belongs to the Asteraceae family and these plants grows around the world due to their maximum uses and yielding. However, now it is too hard to preserve the flowers for long period of time because the flowers get damaged by the fungal organisms. The several diseases such as leaf blight, leaf spot, rust, wilt and bacterial blight has been found in the flowers due to their specific pathogen and the diseases causes the maximum loss in the growth and quality of the flowers also affected. To deduced the infection in the crops, the biofungicides and bioagents was used and protect the plants from another diseases [21]. Nowadays to decrease the infection in the crops, the number of fungicides are used, but now the diseases are controlled by the association of bioagents and some medicinal botanical plants. The most effective bioagent is Trichoderma which is consists of 7 different forms such as T1, T2, T3, T4, T5, T6 and T7 and out of these 7 agent's forms, the maximum inhibition of wilt diseases was observed by T3 form and the more effective botanical plant were *Mentha arvensis* (MA) and stop the growth of *Fusarium oxysporum chrysanthemi* (Foc) [22].

The flower blight disease in *Chrysanthemum* was the most critical disorder which reduce the quality and quantity of the flowers. This disease was caused by the many fungal pathogens such as

Alternaria alternata, *Botrytis cinerea*, *Itersonilia perplexans*, *Mycosphaerella lingulicola* and *Stemphylium vesicarium*. Due to the environmental conditions, the fungi grown very quickly on the field crops and the different kinds of spores has been produced on the flowers which inhibit the growth of the flower that reported in the fields of New Zealand. The various fungicidal treatments were applied to identify the symptoms of diseases and prevent the plants from the disorders. These chemical fungicides control the spore germination and kill the germinated spores that act as protectants of the flowering plants [23].

The number of airborne mycosporophytes was spreading on the *Helianthus annuus* crop due to the external impact of environmental parameters. These fungal spores were reported in the different classes of fungi, each class having individual quantity, which was found in the parts of *Helianthus* plants and the most common fungal spore is *Cladosporium*, that infects the flowering plant and causes the diseases. These tiny mycospores were observed by using the special equipment volumetric Tilak Air Sampler. This air sampler has great spore collection capacity and provides nonstop volumetric information of air sampling for 1 week and allow to analyse the microbial population in both quantitative and qualitative ways [24].

Ornamental plants are the beauty of nature that provides a lot of positive vibes in religious events. The better production of these flowering plants increases the economic status and its commercial values in a whole world. But, with the passage of time, various infections have been caused due to their causal organisms and leads to the problems in the growth of the plants. The main reason of the entry of fungal components is through infected dormant flowering buds and the fungal pathogens produces a number of conidia spores in the form of chain structure which are rapidly dispersed by wind and spread the different diseases in plants. The several chemical agents can be used to stop the infection and prevent the plants from diseases [25]. *Catharanthus roseus* is the perennial flowering plant and this is also known as *Vinca rosea*. this plant contains the ornamental properties as well as medicinal properties which helps to cure the various diseases in human beings and plants as well. But nowadays the properties have been reduced due the formation of fungal organisms which affects the whole plant and causes the several diseases. The main causal pathogen is

Colletotrichum gloeosporioides causes the twig blight disease in the plants and the dark brown to black colour lesions were appeared as symptoms after the rainy season. These symptoms were identified by the morphology of fungal spores, which were formed on the parts of plants [26].

The Human fungal pathogen *Candida albicans* affects the number of plants which belongs to the Apocynaceae family. This fungal organism inhibits the growth and development of the plants and causes the diseases in individual parts of the plants. Every plant has their medicinal values by which treat the causal diseases with the help of additional techniques and chemicals. A Microtiter Plate Assays technique were used to evaluate the causal organisms and also using some extracts of plant's parts to identify the pathogens. The most effective leaf extracts of *Catharanthus roseus* are petroleum ether and distilled water, both shows their positive effects to inhibit the growth of *Candida albicans* [27]. *Hibiscus sabdariffa* is a cosmopolitan flowering as well as medicinal plant. This is commonly cultivated in Upper Egypt having about 94% yielding of its total production. But, the productivity get decreased due to the occurrence of some fungal diseases, mainly root rot, damping-off and wilt diseases that stop the growth of the whole plant from root to leaves by the formation of these several pathogenic soil borne fungi linked with the rotted roots were *Fusarium oxysporum*, *F. solani*, *F. equiseti*, *Rhizoctonia solani* [28].

The *Fusarium* wilt fungal disease is a major problem for the crop of *Chrysanthemum* flowers. Due to the interference of the pathogen in the plants and their parts, the plants do not grow as well and the heavy loss has been occurred in the flower production. The disease was spread very quickly when the environmental condition was favourable for the pathogens and the moderate temperature, more humidity helps to growth and multiplication of the pathogens stopped the vascular system of plants without any external support [29]. In the cultivation of Tulip bulbs, the *Fusarium oxysporum* is a major obstacle because this fungal species infects the plant and inhibit their growth. Various tests were performed for the detection of the pathogens in the different species of Tulip plant and the main evaluation was done by the help of soil infection test and spot inoculation test with the addition of suitable markers and resist the plants to the *Fusarium oxysporum* causal organism [30].

Tagetes flowering plants are very popular and growing in around the world and its parts are also used as medicinal purpose. In Bangladesh, the several diseases were reported in each part of the *Tagetes* plant, especially in two species of *Tagetes* as *T. erecta* and *T. patula*. Commonly leaf spot, foliage blight, grey mold, powdery mildew and anthracnose diseases are found in the most of the plants, but the foliage blight disease is caused due to the causal agent *Aspergillus fumigatus* in *Tagetes*. The two methods (tissue planting and blotter method) were used by using various chemical agents for the detection of diseases [31]. The three main species of *Stagonosporopsis*, which are morphologically identical, but phylogenetically different were affecting the *Chrysanthemum* and *Pyrethrum* flowering plants and caused the ray blight disease in both plants. Ascomata group provided help to the fungal pathogens in their survival and dispersal on the ray florets of *Chrysanthemum* and leads to necrosis of ray florets. Ray blight disease was also found in the commercial *Pyrethrum* in Kenya and Tanzania, while the plants were most infected in Australia and reduce the quantity as well as quality as yielding [32].

Sorghum bicolor is a very popular cereal which is grown in the world and produced a number of natural products by using its high rich sources. Nowadays the occurrence of diseases is so common in every plant and causes a lot of side effects. In *Sorghum bicolor*, leaf spot diseases were caused by the fungal pathogen *Bipolaris sorghicola* and infect the whole plant. To reduce the variations of the pathogen in Sorghum plant, molecular markers (RAPD, RFLP, SSR, ISSR, URP-PCR) were using. These three main RAPD, ISSR, URP-PCR molecular markers help to identify the causal organism and stop the growth of pathogen by the help of their primers [33]. *Chrysanthemum* is the mature flowering plant which is grown for the commercial purposes in the world. These ornamental plants obtained maximum yielding because of their better growth, but due to the weather conditions, the plants were infected by the fungi that causes the diseases in the plants and reduced the production as well as quality of the crop that leads towards the more loss in the economic status. In *Chrysanthemum*, the following diseases were identified in the different parts of the plants like leaf blight, leaf spot, rust wilt, bacterial blight which are resistant to the biotic and abiotic stresses. The foliar blight diseases caused by *Alternaria chrysanthemi*, which

produces the number of spores on the plants that causes the weakness in the plants and the plants became died. These several diseases have been inhibited by using the various alternate fungicides [34].

Tagetes erecta is a demanding flowering plant which is used for every purpose to grace the events. This plant is growing very easily along with its good adaptability in the habitats and its commerciality increases the economic status with high yielding. But since few decades, the several diseases were found in the Marigold plant due to some causal pathogens such as flower blight caused by *Alternaria zinnia*, wilt & stem rot caused by *Phytophthora* spp., *Cercospora* leaf spot caused by *Cercospora megalopotamica*. To control the efficacy of these diseases, a field experiment was conducted in the laboratory by the Pathology Department in 2014-2015 by using some fungicides and biofungicides that control the effects of pathogens and prevent the plant to less yielding. Bavistein & Captan gave the best result to inhibit the variation of disease & its pathogens [35].

Presently, the Orchids are produced in very huge amount as exotic ornamental plant in whole world and its products are transported to the other countries as well. But nowadays, the occurrence of diseases is very common in plants and the disease also moves into the other places along with the plants, where this fungus has not been found previously and spreading very quickly in maximum number. In Orchids, the *Fusarium* species is identified as pathogen that infects the plants and reduce the quality and yielding of the flowers and causes the diseases as leaf and flower spot, leaf and sheath blight, root rots, wilting. Molecular techniques were used to identify the causal agents and the treatment also [36].

In January 2018 to April 2019, three trials were carried out for the investigation and survey on Rose's diseases from the different places. Most of diseases were identified in Rose plants such as, leaf blight, leaf spot, black spot, powdery mildew, mosaic, rust, flower blight, dry brown spot, dieback, foot rot, etc. The major disease of Rose plant is powdery mildew which is caused by *Podosphaera pannosa*, recently recorded in Jashore (district of Bangladesh) and the highest number of leaf spot disease were found in Panisara, while the black spot disease was maximumly identified from Singair. In addition, Researchers conducted a survey on farmer's

socio-economic conditions and issues with growing Roses in Bangladesh and noted their personal opinion for cultivation of Rose plants [37].

3. FUNGAL INFECTION IN PLANTS- A SERIOUS CONCERN

The occurrence of diseases in plants is very serious for the farmers and others as well. Disease caused the various side effects on the plant's parts and decreases the quality and yielding of the crops. But, now a number of alternative methods was used to reduce the power of the pathogens prevent the crop for damaging. In this paper, researchers extract some bacterial strains from the bacteria *Bacillus subtilis* (*spizizenii*) were used for the controlling of leaf blight caused by *Alternaria alternata* in *Tagetes erecta*. This bacterial strain MM19 is most effective to inhibit the further progress of diseases in crops [38]. A field trial was conducted at the experimental farm of Agriculture Department during the two alternate seasons in 2011 & 2012 to test the impacts of the culture filtrates that applied on the Rose crops for their protection from the black spot & powdery mildew diseases which are caused by their specific pathogens *Diplocarpon rosae wolf* & *Sphaerotheca pannosa*. Two culture filtrates prove very effective in controlling the progress of diseases in Rose plants which were named as *Pseudomonas fluorescence* & *Trichoderma viridi*. The sprinkling of these bioassay chemicals on the *Rosa* plants enhance quality of flower and increases their production [39].

Downy mildew is very serious disease in plants, which is formed by their pathogen *Peronospora Danica* and this was investigated in the *Chrysanthemum* plants of Japan. The disease affects the production of the flowers and expands the whole plant. The fungal infection was decreased by using the thermotherapy means treatment with hot water and hot air and biofungicides as spray on the unhealthy plants. A major difference has been appeared between healthy and infected plants [40]. Ornamental plants are very essential for the whole world which provide the grace to each part of earth, but now the huge reduction has been observed in the production and actual quality of the flowers because of the occurrence of several diseases in the plants by their specific causal organisms. These diseases affect the development of crops and create more loss in economy. The most common disease is *Fusarium oxysporum*

chrysanthemi, *Alternaria* leaf spot, sooty mould found in *Chrysanthemum* plants and these diseases were also identified in other crops as well [41].

Powdery mildew is very serious diseases which occurred in the plants especially in *Tabernaemontana divericata* flowering shrub plant. The number of infected leaves were identified due to the diseases Powdery mildew caused by fungal pathogen *Erysiphe elevata* that produces the conidial spores on the plants which inhibit the growth of the plants. To stop the infection and enhances the growth of the plants, various fungicides and molecular techniques were used [42]. *Triticum aestivum* is a well growing cereal crop which is used to make the number of products for edible purpose. But, in the past the various diseases caused by their causal organisms such as *Puccinia triticina* in the wheat crop and the diseases reduce the purity of wheat grains and leads to the economic losses. To control the spreading of diseases from one plant to the other plant, some resistant cultivars were extracted that provide protection against the popular races of pathogens and cure the wheat crops. Several advanced techniques were used to identify and characterized the diseases and its pathogens, like transcriptomics, RNA sequencing, bimolecular fluorescence complementation, virus-induced gene silencing and other approaches [43].

Ornamental plants are playing a major role to enhance the glory of the entire world, but fungal infection causes the diseases in flowering plants that reduce the aesthetic values of the plants and damaged the crops leads to the losses in the production. This is very critical problems to face the pathogenic infection and reduction in the yielding of the crop. But recently, the advance methods of molecular biology tools as spectroscopic and imaging technologies are investigated. These modern equipments helps to identify the causal symptoms and diseases in Rose plants by using fluorescence sensors, spot spectroscopy, hyperspectral, thermal imaging and also decreases the pathogenic effects from the plants [44]. The several diseases were observed from the ornamental plants in US, by which a lot of abnormalities has been produced in the flowering plants and their production. The main fungal component *Phytophthora drechsleri* was reported from the flowers of *Chrysanthemum* that causes the diseases leaf blight, dieback, flower blight, foliar wilt in the rainy season along with high range of

temperature because in this weather, the fungi produced the number of sporangia on the flower which stop the further development of plant. The different kinds of molecular techniques have been applied on the flowering plants to identify and control the diseases and some phytohormones was used to enhance the yielding and quality of flowering production [45].

The epidemic blast diseases infect the rice crops due to its fungal pathogen *Magnaporthe oryzae* and spoil the crops. For the weeds, diseases and pathogens management, several herbicides are widely used but these are not much effective. Among the herbicides, a new chemical agent Glyphosate was withdrawn which was used in the cultivation and this broad-spectrum chemical has antifungal property against the phytopathogens and its diseases to defence the transgenic rice crops. This well deployed chemical also has anti-sporulation power to reduce the spore formation in the rice plants and killed them [46]. The various fungal and bacterial diseases are controlled by using the Plant Growth Promoting Rhizobacteria (PGPR) as a manure and fungicides that helps to inhibit the growth of pathogens in the rice crops especially basmati rice crop. The production and yielding of rice grains has been reduced due to the infection of bacterial and fungal organisms in every part of the plant and causes the several diseases as bacterial leaf blight, brown spot, blast, sheath blight by *Pseudomonas*, *Bacillus*, *Chryseobacterium*, *Streptomyces*. PGPR helps to boost the development of the plant growth traits [47].

Microorganisms are play a vital role in the growth of plants, but these are harmful as well because these infects the root system of plants and inhibit the growth due to the formation of fungal pathogens in soil and causes the diseases in *Abelmoschus* plants. The volatile organic compounds secrets from the *Trichoderma viride* biofungicide in the soil were used to identify the effect of causal organisms on plants and stop the further growth of collar rot pathogen *Sclerotium rolfsii*. The strain BHU-V2 of the most popular biofungicides *T. viride* were used to reduce the effects of causal organisms under the ground and it also helps to enhance the development of the plant and it's all parts [48]. *Vitis vinifera* is the most materialistic valuable edible fruit grown in the whole world. But, number of diseases are produced by their causal organisms in the grapevine crops that deduce the productivity and quality of the crop. The most common diseases

are Downy mildew and Powdery mildew which are caused due to their specific pathogens like as *Plasmopara viticola* belongs to oomycetes class and *Erysiphe necator* belongs to ascomycetes class, which suppress the yield of the crop. These diseases were identified with the help of different classes of molecular defensive genes through the genome wide identification method. These defensive genes are very useful to protect the grapevine production from the powdery and downy mildew diseases and helps to improve the quantity as well as quality of the crops [49].

The most common diseases occurred in the flowering plants such as stem canker, foot rot, wilting, dieback, leaf drop, etc. The stem canker disease is caused by its specific causal organism as *Diaporthe tulliensis* in the *Jasminum officinale* and the symptoms were identified by using the such chemicals in in-vitro condition. Firstly, chlorosis symptoms were appeared on the flowers and also showed that leaves become fall from the branches of the plant. The diseases were controlled by the help of molecular techniques and the chemical treatments [50]. *Tagetes erecta* is commonly known as the African marigold flowering plant which is cultivated as cosmopolitanly. But nowadays the flowering production have been decreased due the interference of diseases causal pathogens on the flowers of the marigold plants. The white cottony mycelial growth appeared on the flower as a whole covering that caused by fungal organism *Sclerotium rolfsii* and its association with other species as *S. sclerotiorum* and the diseases white mould rot is spreading day by day, which stop the growth of the flower and reduce the quality or yielding as well. The diseases were identified on the basis of the external morphological characters of the plants and molecular traits [51].

4. CONCLUSION

Following a review of many studies conducted by scientists from India and other nations, it can be said that the interaction of fungal pathogens with diseased plants lower plant productivity and worsen the financial situation of flowers. This has a negative impact on nature by lowering air quality, which is bad for people and the splendour of the entire planet. Though to improve crop output and plant development, various chemicals and molecular approaches are already employed; however, further research is required to determine how to increase

productivity and prevent plants from growing further.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mukobata H, Yamamoto T, Nahata K, Suzui T. Occurrence of *Pythium* blight caused by *Pythium* spp. in Tulipa. Japanese Journal of Phytopathology. 1989;55(5):594-602.
2. Paulus AO. Fungal diseases of Strawberry. Hort Science. 1990;25(8):885-9.
3. Dhiman JS, Arora JS. Occurrence of leaf spot and flower blight of Marigold (*Tagetes erecta*) in Punjab. Journal of Research-Punjab Agricultural University. 1990;27(2): 231-6.
4. Biswas AC, Sultana K, Begum HA, Iqbal S, Farukuz zaman AK. Diseases of kenaf (*Hibiscus cannabinus*) and mesta (*Hibiscus sabdariffa*) recorded in Bangladesh. Bangladesh Journal of Jute and Fibre Research; 1992.
5. Cook CG, Mullin BA. Growth response of Kenaf cultivars in Root-knot Nematode/Soil-Borne fungi infested soil. Crop Science. 1994;34(6):1455-7.
6. Elad Y, Evensen K. Physiological aspects of resistance to *Botrytis cinerea*. Phytopathology. 1995;85(6):637-43.
7. Shykoff JA, Kaltz O. Effects of the Anther smut fungus *Microbotryum violaceum* on host life-history patterns in *Silene latifolia* (Caryophyllaceae). International Journal of Plant Sciences. 1997;158(2):164-71.
8. Rebanales CR. Chemical seed treatment to enhance emergence and establishment of kenaf (*Hibiscus cannabinus*) under field conditions. Mississippi State University; 1998.
9. Gomathi V, Kannabiran B. Inhibitory effects of leaf extracts of some plants on the anthracnose fungi infecting *Capsicum annum*. Indian Phytopathology. 2000; 53(3):305-8.
10. Saar DE, Polans NO, Sørensen PD, Duvall MR. Angiosperm DNA contamination by endophytic fungi: Detection and methods of avoidance. Plant Molecular Biology Reporter. 2001; 19(3):249-60.

11. Kurze S, Bahl H, Dahl R, Berg G. Biological control of fungal Strawberry diseases by *Serratia plymuthica* HRO-C48. Plant disease. 2001;85(5):529-34.
12. Evans HC, Reeder RH. Fungi associated with *Eichhornia crassipes* (Water hyacinth) in the upper Amazon basin and prospects for their use in biological control. In ACIAR Proceedings. 2000;9(62-70):1998.
13. Ann PJ, Chang TT, Ko WH. Brown root rot of fruit and ornamental trees in Taiwan. Plant Disease. 2002;86(8):820-6.
14. Kharwar RN, Verma VC, Strobel G, Ezra D. The endophytic fungal complex of *Catharanthus roseus* G. Don. Current science. 2008;25:228-33.
15. Göre ME. Epidemic outbreaks of downy mildew caused by *Plasmopara halstedii* on Sunflower in Thrace, part of the Marmara region of Turkey. Plant Pathology. 2009; 58(2).
16. Kamińska M, Berniak H. 'Candidatus *Phytoplasma asteris*' in *Fraxinus excelsior* and its association with ash yellows newly reported in Poland. Plant Pathology. 2009; 58(4).
17. Baiswar P, Kumar R, Chandra S, Ngachan SV. First report of Powdery mildew on Mexican Sunflower in India. Plant Pathology. 2009;58(2).
18. Chaturvedi Y, Singh M, Rao GP, Snehi SK, Raj SK. First report of association of 'Candidatus *Phytoplasma asteris*' (16Srl group) with little leaf disease of Rose (*Rosa alba*) in India. Plant Pathology. 2009;58(4).
19. Elmanama AA, Alyazji AA, Abu-Gheneima NA. Antibacterial, antifungal and synergistic effect of *Lawsonia inermis*, *Punica granatum* and *Hibiscus sabdariffa*. Annals of Alquds Medicine. 2011;7.
20. Wikee S, Cai L, Pairin N, McKenzie EH, Su YY, Chukeatirote E, Thi HN, Bahkali AH, Moslem MA, Abdelsalam K, Hyde KD. *Colletotrichum* species from Jasmine (*Jasminum sambac*). Fungal Diversity. 2011;46(1):171-82.
21. Kumar GA, Kamanna BC, Benagi VI. Management of *Chrysanthemum* leaf blight caused by *Alternaria alternata* Keissler under field condition. Plant Archives. 2011;11(1):553-5.
22. Singh PK, Vijay K. Biological control of *Fusarium* wilt of *Chrysanthemum* with *Trichoderma* and Botanicals. Journal of Agricultural Technology. 2011;7(6):1603-13.
23. Singh G, Milne KS. Field evaluation of fungicides for the control of *Chrysanthemum* flower blight. New Zealand Journal of Experimental Agriculture. 1974;2(2):185-8.
24. Kshirsagar JJ, Pande BN. Prevalence of *Cladosporium* spores over Sunflower fields at Rajuri (N) MS, India. Science Research Reporter. 2012;2(1):66-8.
25. Singh VK, Singh Y, Kumar P. Diseases of Ornamental plants and their management. Eco-friendly innovative approaches in plant disease management. 2012:543-72.
26. Pankaj S. New record of twig blight on *Catharanthus roseus* in India. African Journal of Microbiology Research. 2013; 7(38):4680-2.
27. Wankhede SB, Routh MM, Rajput SB, Karuppayil SM. Antifungal properties of selected plants of Apocynaceae family against the human fungal pathogen *Candida albicans*. International Current Pharmaceutical Journal. 2013;2(7): 122-5.
28. Hassan N, Shimizu M, Hyakumachi M. occurrence of root rot and vascular wilt diseases in Roselle (*Hibiscus sabdariffa*) in Upper Egypt. Mycobiology. 2014;42(1): 66-72.
29. Singh PK, Kumar V. Fusarium wilt of *Chrysanthemum*—problems and prospects. Plant Pathology & Quarantine. 2014;4(1): 33-42.
30. Tang N, van der Lee TA, Jalink H, van der Schoor R, Shahin A, Bijman PJ, van Tuyl JM, Arens PF. Genetic mapping of resistance to *Fusarium oxysporum* in *Tulip*. Gewasbescherming. 2014;45(1):12.
31. Aktar M, Shamsi S. Blight of two species of Marigold (*Tagetes erecta*) caused by *Aspergillus fumigatus*. Bangladesh J. Plant Pathol. 2015;31(1&2):1-6.
32. Vaghefi N, Pethybridge SJ, Hay FS, Ford R, Nicolas ME, Taylor PW. Revisiting *Stagonosporopsis* species associated with *Chrysanthemum* and *Pyrethrum* ray blight. Australasian Plant Pathology. 2016;45(6): 561-70.
33. Kandan A, Akhtar J, Singh B, Pal D, Chand D, Rajkumar S, Agarwal PC. Genetic diversity analysis of fungal pathogen *Bipolaris sorghicola* infecting *Sorghum bicolor* in India. Journal of Environmental Biology. 2016;37(6):1323.
34. Kavitha M, Kumari KL. Management of foliar blight (*Alternaria chrysanthemi*) of *Chrysanthemum*.

35. Chandel S, Kumar V. Evaluating fungicides and biofungicide for controlling *Cercospora* leaf spot on Marigold. International Journal of Current Microbiology and Applied Sciences. 2017; 6(5):2072-7.
36. Srivastava S, Kadooka C, Uchida JY. *Fusarium* species as pathogen on Orchids. Microbiological research. 2018;207:188-95.
37. Chhanda ST. Investigation on nursery and field diseases of rose (*Rosa spp.*) in selected areas of Bangladesh (Doctoral dissertation, Department of Plant Pathology).
38. Priyanka R, Nakkeeran S, Pravin IA, Moorthy AK, Sivakumar U. Antifungal activity of *Bacillus subtilis* subsp. *spizizenii* (MM19) for the management of *Alternaria* leaf blight of Marigold. Journal of Biological Control. 2018;32(2):95-102.
39. Amin F, Qazi NA, Banday S, Dar SH, Shahnaz E. Biological control of Powdery mildew and Black spot diseases of Rose. Journal of Pharmacognosy and Phytochemistry. 2018;7(3):2826-8.
40. Yoshida K, Asano S, Sumikawa Y. Occurrence of *Chrysanthemum* downy mildew in Nara prefecture and its control with thermotherapy and fungicide. Annual Report of The Kansai Plant Protection Society. 2019;61:79-84.
41. Borah M, Rajkhowa M, Ali S. Occurrence of diseases in floricultural crops in and around Jorhat, Assam. Int. J. Econ. Plants. 2019;6:54-63.
42. Xu D, Zeng Y, Zhang J, Xu J, Qiao F. First report of powdery mildew of crape Jasmine (*Tabernaemontana divaricata*) caused by *Erysiphe elevata* in China. Plant Disease. 2021;105(4):1203.
43. Prasad P, Savadi S, Bhardwaj SC, Gupta PK. The progress of leaf rust research in Wheat. Fungal biology. 2020;124(6): 537-50.
44. Traversari S, Cacini S, Galieni A, Nesi B, Nicastro N, Pane C. Precision agriculture digital technologies for sustainable fungal disease management of ornamental plants. Sustainability. 2021;13(7):3707.
45. Krasnow CS, Rechcigl NA, Olson JD, Schmitz LT, Jeffers SN. First report of Stem and Foliage Blight of *Chrysanthemum* Caused by *Phytophthora drechsleri* in the United States. Plant Disease. 2021;105(11):3765.
46. Mehta S, Kumar A, Achary VM, Ganesan P, Rathi N, Singh A, Sahu KP, Lal SK, Das TK, Reddy MK. Antifungal activity of glyphosate against fungal blast disease on glyphosate-tolerant OsmEPSPS transgenic Rice. Plant Science. 2021; 311:111009.
47. Jasrotia S, Salgotra RK, Sharma M. Efficacy of bioinoculants to control of bacterial and fungal diseases of Rice (*Oryza sativa*) in northwestern Himalaya. Brazilian Journal of Microbiology. 2021;52(2):687-704.
48. Singh J, Singh P, Vaishnav A, Ray S, Rajput RS, Singh SM, Singh HB. Belowground fungal volatiles perception in Okra (*Abelmoschus esculentus*) facilitates plant growth under biotic stress. Microbiological Research. 2021;246:126721.
49. Goyal N, Bhatia G, Garewal N, Upadhyay A, Singh K. Identification of defense related gene families and their response against Powdery and Downy mildew infections in *Vitis vinifera*. BMC genomics. 2021;22(1):1-6.
50. Hsu CC, Hsiao HY, Huang TC, Shen YM. First report of stem canker caused by *Diaporthe tulliensis* on Jasmine in Taiwan. Plant Disease. 2022;13(ja).
51. Kumar P, Sharma S, Singh R, Singh P, Kumar A. First report of *Sclerotinia sclerotiorum* causing white rot of Marigold in Punjab, India. Journal of Plant Pathology. 2022;104(1):435.

© 2023 Arti and Kaur; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/97839>