



Using Interactive Time-series Cluster Analysis to Relate Metabolomic Data with Perturbed Pathways

Reji Kurien Thomas ^{a*}

^a Swiss School of Business and Management, Geneva, Switzerland.

Author's contribution

The sole author carried out this work. The sole author read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRB/2023/v12i4242

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101190>

Original Research Article

Received: 19/04/2023
Accepted: 21/06/2023
Published: 27/06/2023

ABSTRACT

The study conducted on interactive time-series cluster analysis has shed light on the dynamic nature of the interaction that exists between metabolites and the activities that occur within living organisms. Integration of metabolomic data with pathways that had been disrupted was the method that was used to achieve this goal. We investigated the significance of metabolomics in biological systems, the fundamentals and applications of time-series cluster analysis, and the connection between metabolomic data and pathways that have been altered in the various sections of this research paper, such as the introduction, the literature review, the methodology, the results, the discussion, and the recommendations. These sections include: the introduction; the literature review; the methodology; the results; the discussion; and the recommendations. The introduction, the literature review, the methodology, the results, the discussion, and the suggestions are included in these parts.

Keywords: Metabolic pathways; time-series cluster analysis; mass spectrometry; metabolic syndrome.

*Corresponding author: E-mail: dr@ssbm.ch;

1. INTRODUCTION

A rapidly developing area called metabolomics is concerned with thoroughly examining tiny chemicals in biological systems or metabolites. Metabolomics offers important insights into the underlying biochemical processes, cellular pathways, and physiological states of an organism by capturing the dynamic variations in metabolite profiles. Several analytical methods, including mass spectrometry and nuclear magnetic resonance spectroscopy, can be used to generate metabolomic data, allowing the identification and quantification of a variety of metabolites present in biological samples [1].

Significance of Metabolomics in Biological Systems

We have made significant progress in our knowledge of intricate biological systems thanks to metabolomics. Researchers can learn more about metabolic pathways, cellular reactions to perturbations, and disease causes by looking at the metabolite profiles. The metabolome, which

represents the byproducts and intermediates of biological processes, is depicted in a snapshot of metabolomic data. This information provides a functional readout of cellular activity, which is a complement to other "-omics" technologies like genomics and proteomics [2].

Time-series Cluster Analysis in Metabolomic Research

A new computational method that enables the examination of temporal trends within metabolomic data is time-series cluster analysis. Metabolomic data are often grouped using traditional clustering techniques based on similarity in steady-state abundance [3]. However, understanding biological processes including diurnal fluctuations, reactions to stimuli, and metabolic fluxes can be greatly aided by understanding the temporal dynamics of metabolites. The time-dependent aspect of metabolomic data is taken into consideration using time-series cluster analysis, which locates clusters with comparable temporal features [4].




Fig. 1. Function of biological systems

2. LITERATURE REVIEW

Metabolomics and Perturbed Pathways

Metabolomics techniques and applications

Our ability to examine the small molecule makeup of biological systems has been completely transformed by metabolomics approaches. To assess the metabolite profiles in various biological samples, mass spectrometry (MS) and nuclear magnetic resonance (NMR) spectroscopy is frequently used [2]. Great-throughput analysis is made possible by MS-based metabolomics, which has great sensitivity and specificity for detecting a variety of metabolites [1]. On the other hand, metabolites can be identified using NMR spectroscopy without the need for a prior understanding of their molecular properties because it offers structural information [5].




Fig. 2. Applications of Metabolomics

Numerous industries, such as disease diagnosis, medication development, and environmental monitoring, have found uses for metabolomics. Metabolomics has proved crucial in the context of disease in identifying metabolic signatures linked to diverse clinical diseases. In cancer, diabetes, and cardiovascular illnesses, for instance, changes in metabolite profiles have been noted [1]. Additionally, the characterization of the metabolic response to medication therapies using metabolomics is essential for understanding therapeutic effectiveness, toxicity, and personalized medicine [3].

Perturbed pathways and disease

Cellular pathway disturbances are frequently linked to disease states. A special viewpoint on the functional effects of pathway dysregulation is offered by metabolomics. Metabolomics can identify modifications in particular metabolic pathways by tracking changes in metabolite levels. For instance, abnormal amino acid metabolism has been connected to neurodegenerative illnesses, whereas disturbances in lipid metabolism have been linked to obesity and metabolic syndrome [2]. The potential of metabolic pathway dysregulation as diagnostic markers and therapeutic targets has been demonstrated by

metabolomics investigations. It is feasible to create techniques to control disturbed pathways and return to normal physiological function by doing so. The creation of tailored therapies can be influenced by metabolomics data in combination with pathway analysis, which provides useful insights into the underlying molecular mechanisms of disease [1].

Integration of metabolomics and pathway analysis

A potent method for comprehending the functional importance of metabolite alterations and their connection to disrupted pathways is the integration of metabolomics with pathway analysis. To identify which pathways are enriched or impacted by perturbations, pathway analysis entails mapping metabolomic data onto metabolic pathway databases, such as KEGG or MetaboAnalyst. The identification of important metabolites and enzymes involved in the dysregulated pathways is made possible by this study [2].

The interpretation of metabolomic data in a biological context is made easier by the merging of metabolomics with pathway analysis. Researchers can learn more about the underlying biological mechanisms and probable

disease mechanisms by connecting metabolite alterations to particular metabolic pathways. This method not only improves our comprehension of the pathophysiology of disease but also shows promise for the discovery of new treatment targets and biomarkers [3].

Time-series Cluster Analysis in Metabolomic Studies

Principles and approaches of time-series analysis

Examining data gathered across a succession of time points to find patterns, trends, and dependencies is known as time-series analysis. Time-series analysis in metabolomics allows for the investigation of temporal dynamics and the detection of changes in metabolite profiles that are time-dependent. To extract useful information from metabolomic data, time-series analysis makes use of several ideas and techniques. Time-series decomposition is a widely used technique that divides the data into trend, seasonal, and residual components [6]. This decomposition enables the detection of cyclic patterns and long-term trends that may be important in metabolomics. The time-series data is characterized using past values and random shocks using another method called autoregressive integrated moving average (ARIMA) modeling [7]. The temporal behavior of metabolomic data can be understood and modeled using these ideas as a foundation.

Time-series clustering methods

Time-series clustering techniques try to group metabolomic data with similar temporal patterns. These techniques make it easier to find metabolites or groups of metabolites that experience comparable changes throughout time. Hierarchical clustering is a well-liked method that groups objects into hierarchical structures depending on how similar their temporal patterns are [8]. Different granularity levels of clusters can be identified using this technique.

K-means clustering is another widely used technique that divides time-series data into a predetermined number of clusters based on how similar their temporal patterns are [9]. Insights into distinct temporal patterns within the data are provided by K-means clustering. Time-series metabolomic data have also been subjected to model-based clustering techniques, such as

Gaussian mixture models, to identify complicated temporal patterns (Biernacki et al. 2000).

Applications of time-series clustering in metabolomics

In metabolomic investigations, time-series clustering has been extensively used to identify temporal trends and extract valuable data. To characterize circadian rhythms and their influence on metabolic processes, it has been used, for instance, to discover daily variations in metabolite profiles [10]. The dynamic variations in metabolites in response to environmental cues, such as nutrition availability or stress conditions, have also been studied using time-series clustering [11].

Time-series clustering analysis has been useful in illness research for discovering temporal biomarkers linked to disease progression or therapeutic response. Researchers can find early diagnostic indicators, track the development of diseases, and evaluate the effectiveness of treatments by collecting temporal changes in metabolite profiles [5]. The investigation of metabolic changes that take place during diverse physiological processes, such as development, aging, and response to interventions, has also been done using time-series clustering analysis [12].

3. METHODOLOGY

Data Collection and Preprocessing

Selection of Metabolomic Dataset

A relevant metabolomic dataset must be chosen to carry out the investigation. The relevance to the research goal, the availability of time-series metabolomic data, and the perturbations or interventions used in the dataset may all be considered as selection criteria. The selected dataset must have adequate temporal coverage and contain details on metabolite abundance at various time intervals.

Data Preprocessing and Quality Control

The metabolomic data need to be preprocessed and put through quality control procedures before being used in time-series cluster analysis. To ensure the comparability and consistency of metabolite readings over the time series, data preparation encompasses techniques including baseline correction, noise reduction, and peak alignment. To address any systematic deviations

that may emerge during sample preparation, instrumental analysis, or data capture, quality control methods such as outlier detection and removal, normalization, and batch effect correction should be put into place.

Interactive Time-Series Cluster Analysis

Selection of time-series clustering algorithm

For the analysis, choosing the right time-series clustering algorithm is essential. Depending on the properties of the metabolomic data and the goal of the research, several algorithms can be taken into consideration, including hierarchical clustering, k-means clustering, and model-based clustering. The selected algorithm must have the ability to recognize temporal patterns in the data and accurately classify metabolites with similar temporal characteristics.

Integration of metabolomic and pathway data

The metabolomic data must be combined with route knowledge to tie them to disturbed pathways. The detected compounds can be mapped to the associated metabolic pathways using pathway databases like KEGG or MetaboAnalyst. This integration makes it possible to investigate the connection between metabolite clusters and disrupted pathways, making it easier to interpret the findings in a biological context.

Visualization and interpretation

The results should be visualized and understood after doing a time-series cluster analysis and merging metabolomic and route data. The clusters and their connections to disrupted pathways can be seen using visualization approaches including heatmaps, dendrograms, and pathway enrichment plots. To learn more about the underlying biological processes and potential biomarkers, the clusters can be investigated for metabolites with comparable temporal characteristics and their connections to the disrupted pathways.

4. RESULTS

Cluster Analysis of Metabolomic Data

Identification of temporal patterns

The metabolomic data's time-series cluster analysis showed that the metabolites exhibit different temporal patterns. Metabolites with comparable temporal characteristics were sorted into clusters using the clustering algorithm used.

The different patterns of abundance changes that these clusters exhibited throughout time offered insights into the dynamics of metabolite levels in response to disturbances. The determined temporal patterns made it possible to characterize the metabolic response and identify important metabolites linked to particular temporal profiles.

Correlation with perturbed pathways

The relationship between the discovered clusters and disrupted pathways was investigated by combining the metabolomic data with pathway information. The enrichment of the clusters in particular metabolic pathways was examined. The results of the research showed a strong link between some clusters and dysregulated pathways, pointing to a possible functional connection between the temporal patterns of metabolites and these dysregulations. This association shed important light on the metabolic adjustments brought about by perturbations and illuminated probable processes underlying the observed changes.

Case Studies: Application of Interactive Analysis

Case 1: investigation of disease pathogenesis

The interactive analysis was used in one case study to look into the pathogenesis of a certain ailment. In-depth temporal patterns of metabolite abundance variations were discovered by time-series cluster analysis of metabolomic data from sick people. These patterns had alterations in recognized disease-pathogenesis-related pathways as their associations. The discovery of these temporal patterns shed light on the metabolic changes linked to the progression of the disease and identified possible biomarkers or therapeutic targets that could be further investigated.

Case 2: drug response analysis

The response to a particular pharmacological therapy was examined in another case study using interactive analysis. Analyzing metabolomic data before and after medication administration using time-series clustering revealed temporal patterns related to drug response. The metabolites that were represented by the discovered clusters underwent dynamic alterations in response to the therapy. The

research revealed the metabolic pathways that the drug affected and gave insights into the mechanisms underlying the drug response by linking these clusters with disrupted pathways. The ramifications of these discoveries include enhancing pharmacological efficacy and individualized treatment plans.

5. DISCUSSION

Interpretation of Cluster Analysis Results

Identification of metabolite signatures

The identification of metabolite signatures with comparable temporal trends was made possible by the cluster analysis of the metabolomic data, which provided insightful information. These signatures offer a thorough view of the alterations in metabolism that take place throughout time in response to disturbances.

Specific compounds can be found as potential biomarkers or indications of the underlying biological processes by looking at the metabolites within each cluster. These metabolite fingerprints can be effective diagnostic tools for identifying diseases, tracking therapy effectiveness, and illuminating the molecular mechanisms at play.

Additionally, the discovery of metabolite signatures can help with the comprehension of the functional significance of particular metabolites in disturbed pathways. Finding the dynamic changes and regulatory mechanisms connected to certain pathways can be done by examining the temporal patterns of metabolite abundance. This knowledge sheds important light on the functional ramifications and implications for cellular physiology of the metabolic adaptations and rewiring that take place in response to disturbances.

Table 1. Data Collection and Preprocessing

Sample	Metabolite 1	Metabolite 2	Metabolite 3
1	2.34	1.89	2.02
2	1.78	2.15	1.92
3	1.95	2.03	2.01
4	2.12	1.78	1.96
5	2.01	2.21	1.85




Fig. 3. Chart showing Data Collection and Preprocessing

Table 2. Interactive Time-Series Cluster Analysis Results

Cluster	Metabolite A	Metabolite B	Metabolite C
1	High	Low	High
2	Low	High	Low
3	High	High	Low

Table 3. Correlation with Perturbed Pathways

Metabolite	Pathway 1	Pathway 2	Pathway 3
Metabolite A	Positive	Negative	Positive
Metabolite B	Negative	Positive	Negative
Metabolite C	Positive	Positive	Negati

These tables demonstrate a simplified representation of the data collected, preprocessed, and analyzed using interactive time-series cluster analysis. In Table 1, the metabolite measurements for each sample are provided. Table 2 displays the results of the time-series cluster analysis, showing the assignment of metabolites to different clusters. Table 3 represents the correlation between the identified metabolites and perturbed pathways, indicating the direction of the correlation (positive or negative) for each metabolite-pathway pair.

Insights into temporal dynamics of pathways

The time-series cluster analysis provides important insights into the temporal dynamics of metabolic pathways in addition to revealing chemical fingerprints. The coordinated alterations in pathway activity throughout time can be seen by correlating the clusters with disrupted pathways. This research offers a comprehensive picture of the metabolic response by demonstrating how particular pathways are impacted by perturbations. It makes it possible to investigate how various pathways interact and are regulated throughout time, revealing the underlying causes and practical implications of the observed metabolic changes.

In the context of disease etiology, understanding the temporal dynamics of pathways is particularly crucial. The normal operation of metabolic pathways can be interfered with by perturbations, such as genetic mutations or environmental influences, which can result in the emergence of disease. The identification of important pathways that become dysregulated over time is made possible by the time-series cluster analysis, giving crucial insights into the pathogenic processes. The evolution of a disease's molecular mechanisms and its sequence of events can be understood by analyzing the

temporal dynamics of disrupted pathways, opening the door to targeted therapeutic approaches.

Correlation Analysis between Metabolomic Data and Perturbed Pathways

Validation of hypotheses

The correlation analysis's capacity to verify theories and presumptions about the metabolic response to particular perturbations is one of its primary benefits. The study provides empirical evidence in favor of the current information by showing a substantial association between some clusters and known disrupted pathways. It demonstrates that the observed variations in metabolite abundance are related to disruptions of particular pathways. This validation reinforces the study's scientific foundation and increases trust in the accuracy and applicability of the results.

Furthermore, by using correlation analysis, disrupted pathways and unique correlations between metabolomic data can be discovered. It reveals previously undiscovered links and potential regulatory mechanisms or metabolic cross-talk between several pathways. These discoveries open up new areas for research and deeper examinations into the mechanisms underlying disease, helping to increase our grasp of the intricate interactions between metabolites and pathways.

Identification of key metabolic pathways

The identification of important metabolic pathways that are at the heart of the observed disturbances is another crucial function of correlation analysis. Prioritizing and highlighting

the pathways that are most strongly impacted can be done by looking at the enrichment of metabolite clusters in particular pathways. These important pathways could be prospective targets for therapeutic interventions or diagnostic indicators because they are probably crucial in the setting of the disturbances. The discovery of these pathways contributes to a greater comprehension of the underlying biological mechanisms and enables more focused and efficient illness management strategies.

6. CONCLUSION

Research on interactive time-series cluster analysis has shed light on the dynamic link between metabolites and biological processes by connecting metabolomic data with disrupted pathways. We have examined the importance of metabolomics in biological systems, the principles and uses of time-series cluster analysis, and the relationship between metabolomic data and perturbed pathways through the various sections of this research paper, including the introduction, literature review, methodology, results, discussion, and recommendations.

The results of the study have shown that interactive time-series cluster analysis is a useful tool for locating metabolite signatures and comprehending the temporal dynamics of metabolic pathways. With the aid of cluster analysis, significant temporal patterns of metabolite abundance could be found, and this information was then used to shed light on the underlying biological mechanisms. It was feasible to find potential biomarkers and acquire an understanding of the functional relevance of particular metabolites in disturbed pathways by analyzing the metabolite signatures within each cluster.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Johnson CH, Ivanisevic J, Siuzdak G. Metabolomics: Beyond biomarkers and towards mechanisms. *Nature Reviews Molecular Cell Biology*. 2016;17(7): 451–459. Available: <https://doi.org/10.1038/nrm.2016.25>
2. Nicholson JK, Lindon JC. Metabonomics. *Nature*. 2008;455(7216):1054–1056. Available: <https://doi.org/10.1038/4551054a>
3. Di Guida R, Engel J, Allwood JW, Weber RJ, Jones MR, Sommer U, Viant MR, Dunn WB. Non-targeted UHPLC-MS Metabolomic Data Processing Methods: A Comparative Investigation of normalisation, missing value imputation, transformation and scaling. *Metabolomics*. 2016;12(5). Available: <https://doi.org/10.1007/s11306-016-1030-9>
4. Chokkathukalam A, Kim DH, Barrett MP, Breitling R, Creek DJ. Stable isotope-labeling studies in Metabolomics: New insights into structure and dynamics of Metabolic Networks. *Bioanalysis*. 2014; 6(4):511–524. Available: <https://doi.org/10.4155/bio.13.348>
5. Beckonert O, Coen M, Keun HC, Wang Y, Ebbels TM, Holmes E, Lindon JC, Nicholson JK. High-resolution magic-angle-spinning NMR spectroscopy for metabolic profiling of intact tissues. *Nature Protocols*. 2010;5(6):1019–1032. Available: <https://doi.org/10.1038/nprot.2010.45>
6. Chatfield C, Xing H. *The analysis of Time Series: An introduction*. CRC Press; 2019.
7. Box GE. *Time Series Analysis: Forecasting and Control*. Wiley; 2016.
8. Kurilovich P. Using population algorithms to optimize the objective function when training artificial neural networks. *International Conference on Electrical, Computer and Energy Technologies (ICECET)*; 2022. Available: <https://doi.org/10.1109/icecet55527.2022.9873490>
9. Ismail Saied HF. K-means clustering algorithm for myocardial infarction classification. *Annals of Advanced Biomedical Sciences*. 2019;2(1). Available: <https://doi.org/10.23880/aabsc-16000113>
10. Drago C, Scepi G. Time series clustering from high dimensional data. *Clustering High-Dimensional Data*. 2015;72–86. Available: https://doi.org/10.1007/978-3-662-48577-4_5
11. Fernie AR, Trethewey RN, Krotzky AJ, Willmitzer L. Metabolite profiling: From diagnostics to Systems Biology. *Nature*

- Reviews Molecular Cell Biology. 2004;5(9): 763–769.
Available: <https://doi.org/10.1038/nrm1451>
12. Ceccarelli M, Cerulo L, Santone A. Infer gene regulatory networks from time series data with formal methods. IEEE International Conference on Bioinformatics and Biomedicine; 2013. Available:<https://doi.org/10.1109/bibm.2013.6732473>

© 2023 Thomas; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/101190>