



Efficacy of Salicylic Acid in Response to Plant Stress Tolerance, Growth and Productivity: A Review

Zuby Gohar Ansari ^{a++}, Hridesh Harsha Sarma ^{b*},
Anasuya Boruah ^{c#}, Gangadhara Doggalli ^{d†}, Raj Bahadur ^{e‡},
Utkarsh Tripathi ^{f^} and Doppalapudi Vijaya Rani ^{g##}

^a Department of Plant Physiology, College of Agriculture Indore under RVSKVV, Gwalior, India.

^b Department of Agronomy, Institution: Assam Agricultural University, India.

^c Assam Agricultural University, Jorhat, Assam, India.

^d Department of Genetics and Plant Breeding, University of Agricultural Sciences, Dharwad, Ettinagudda Road, Krishi Nagar, Dharwad, India.

^e ANDUA&T, Ayodhya, UP, India.

^f Department of Linguistics, College of Agriculture, ANDUAT, Kumarganj, Ayodhya, India.

^g Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, India.

Author's contribution

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113573

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109071>

Review Article

Received: 08/09/2023

Accepted: 15/11/2023

Published: 29/11/2023

⁺⁺ Contractual Teacher;

[#] M. Sc. Scholar;

[†] PhD Scholar;

[‡] Asoc. Prof. Agronomy;

[^] Associate Professor, (English);

^{##} Ph.D. Research Scholar;

*Corresponding author: E-mail: hrideshsarma.official@gmail.com;

ABSTRACT

Plant stress generated by various climatic and anthropogenic activities has exacerbated the degradation of agricultural systems and productivity throughout the developmental era, and it is currently recognized as the world's most significant and possibly deadly danger impacting crop plant economic output. Nonetheless, phytohormones have been found as a potent tool for minimizing the detrimental effects of stressors in agricultural plants in a sustainable manner. Salicylic acid (SA), one of the most important phytohormones, is required for the regulation of plant growth, development, ripening, and defensive responses. A lot of interest has been generated by SA's defensive mechanism. Major agricultural crops' ability to withstand stress has been said to be much improved. SA utilization is influenced by the amount of SA applied, the application method, and the state of the plants (such as their developmental stage and level of acclimation). In this review, we have focus on the efficacy of SA on mitigating various plant biotic and abiotic stress under adverse environmental conditions and its role on plant growth, development and productivity.

Keywords: Salicylic acid; stress tolerance; defence mechanism; growth; development and productivity.

1. INTRODUCTION

Current climate circumstances and anthropogenic activity are among the most damaging elements causing massive agricultural losses throughout the world. The aforementioned challenges, such as high temperatures, salinity, drought, and cold, are significant variables that limit agricultural output. With the environment being altered and rare natural resources being exploited more economically, the global agricultural goal of increasing food production by 70% by the year 2050 for approximately 2.3 billion new-born people is facing serious challenges [1]. "Impact of climate change on ecosystems have had a serious negative influence on the gross agronomy, making these efforts ineffective" [2]. Crops are being subjected to an increase in stressors (biotic and abiotic stress) as a consequence of the impact of human activity and the intensification of climate change, which has become the main factor in the deterioration of agricultural systems and the loss in grain output [3]. As a result, developing plants to withstand or tolerate stressors (biotic and abiotic) using previously discovered resistance mechanisms will be part of the sustainable development approach. Salicylic acid (SA), for instance, has been shown to be crucial in resistance and defensive signaling under both biotic and saline stress [4].

"Salicylic acid (SA) is a naturally occurring small-molecule phenolic compounds that possesses an aromatic ring and a hydroxyl group. It functions as a signal sensor to control plant response. Through the control of procedures including

antioxidant defense, nitrogen metabolism, photosynthesis, and water stress, it shields plant cells from the toxicity of ion buildup and cell death" [5]. Because of its numerous physio-biochemical and developmental activities in hormonal interactions as well as its antioxidant characteristics, salicylic acid (SA), a common plant phenolic molecule, is an efficient stress reliever [6]. It has been shown to increase tolerance and defense against pathogen attack [7], as well as to mitigate the harmful effects of several environmental stresses on plants, including low temperature and chilling [8,9], high temperature and drought [10], and salinity [11]. Additionally, it improves seedling emergence and germination rates and percentages, ensuring adequate crop stand under a variety of environmental circumstances [12].

Salicylic acid is regarded as a crucial signalling molecule that contributes to local and endemic disease resistance in plants as a result of diverse pathogenic assaults [13]. In addition to conferring disease resistance, SA may be able to regulate how plants respond to various oxidative stresses. Significant advancements have been achieved in pinpointing the essential elements and comprehending the functions of SA and nutrients connected to plant defense pathways. In this review, the multifunctional roles of salicylic acid in crops have been summarized and illustrated. These roles include defense mechanisms, tolerance to a variety of biotic and abiotic stresses, biosynthesis, signaling, effectiveness on bio-productivity, growth, activities of various enzymes in plants, as well as the necessary space constraints.

2. SALICYLIC ACID (SA) BIOSYNTHESIS PATHWAY

The endogenously produced phytohormone SA (2-hydroxybenzoic acid) that is present in plants is a phenolic substance with a seven-carbon (C) backbone. Two primary secondary metabolite-producing pathways—the shikimic acid system and the malonic acid pathway—produce these plant phenolic compounds. However, the majority of phenolic compounds in plants are created via the shikimic acid system, which does so by converting aromatic amino acids, such as phenylalanine, which is the precursor of SA, into precursors of carbohydrates acquired through glycolysis and the pentose phosphate pathway. Although another system termed the isochorismate pathway also accumulates SA in plants, the phenylalanine pathway is responsible for producing the majority of the SA there is [14,15]. The phenylalanine ammonia-lyase (PAL) enzyme acts on the phenylalanine to remove ammonia, which results in the formation of trans-cinnamic acid in the phenylalanine pathway. Cinnamic acid serves as a precursor for two different chemicals, one of which is coumaric acid, which is created when an enzyme called cinnamate-4-hydroxylase (C₄H) hydroxylates cinnamic acid at the C4 position. The subsequent oxidation of its side chain and hydroxylation of coumaric acid lead to the production of SA.

Cinnamic acid is converted into a second product, benzoic acid, which is then subjected to ortho-positional hydroxylation, which results in the creation of SA [15].

By virtue of the activity of the enzyme isochorismate synthase (ICS), chorismate serves as a precursor for isochorismate in the isochorismate pathway (Fig. 1). The enzyme isochorismate pyruvate lyase converts isochorismate into SA [15]. The ICS level increases in plant cells in response to biotic or abiotic stressors, which results in a high level of SA in plant defense mechanisms [16]. The majority of the SA in plants still exists in its inactive, glycosylated form. The enzyme UDP-glucose SA-glucosyltransferase glycosylates SA. However, SA glucosidase changes SA from its inert state into one that is active. These two enzymes also function as regulators, determining the degree of SA in plant cells under various environmental conditions. Arabidopsis under stress uses isochorismate to create SA mostly in chloroplasts. Other crops may have other pathways for producing SA. Under typical development conditions, rice accumulates significant amounts of SA, around tenfold higher than that of Arabidopsis leaves [17]. SA biosynthesis is carefully regulated during biotic and abiotic stress.




Fig. 1. Biosynthesis of salicylic acid
(Source: Yang et al., [18])

3. SIGNALLING OF SALICYLIC ACID (SA)

Regulation of the SA biosynthesis pathway occurs at both the transcriptional and posttranscriptional stages. Plant transcriptional analysis has revealed the presence of MYBs (MYELO BLAST; MYB 96 and MYB30), WRKYs (WRKY28 and WRKY46), and WIPK (Wound-Induced Transcription Factors of Mitogen-Activated Protein Kinase) genes. According to Vidhyasekaran [19], these genes increase SA synthesis in plants by favorably regulating the isochorismate synthase pathway of the ICS gene encoding. An important part in establishing and signaling a defense response against diverse stressors is played by the well-known naturally occurring signaling molecule known as SA. For the downstream reaction to occur, the SA that plants make must be detected by its receptors. There are a number of SA-binding proteins (SABPs) that can detect SA. By influencing physiological and biochemical processes, salicylic acid functions as an endogenous natural signal molecule and is essential to defensive systems [20]. According to Kawano *et al.* [14], salicylic acid produced by cells can readily enter and exit cells, tissues, and organs; and this movement is carefully controlled by ROS and Ca²⁺ [21]. Additionally, SA was shown to engage in signalling and the control of gene expression during the senescence of *Arabidopsis* leaves Morris *et al.* [22].

Salicylic acid functions as a signalling molecule and controls the formation of chloroplasts [23], photosynthetic activity [24], gravitropism [25], and the prevention of fruit ripening [26]. Most genes and signaling pathways that are sensitive to salt and are in charge of cell death in response to salt stress are made more active by salicylic acid. These responsive genes produce the enzymes sinapyl alcohol dehydrogenase (SAD), cinnamyl alcohol dehydrogenase (CAD), and cytochrome P₄₅₀, which are all involved in secondary metabolite pathways. They also produce heat shock proteins (HSPs), chaperones, and chaperone-related proteins [27]. Ascorbate peroxidase (APX), glutathione reductase (GR), and superoxide dismutase (SOD) are part of the antioxidant system that SA activates to protect cells from stress [28]. By speeding up the rate of CO₂ absorption due to an increase in stomatal conductance, the exogenous application of SA (500 µM) relieved the drought stress in *Hordeum vulgare* [29]. The AsA-GSH pathway's enzymatic and nonenzymatic components are regulated by SA

supplementation, which also helps plants subjected to drought to overcome oxidative stress [30]. According to Miura *et al.* [31], the expression of the *SIZ1* gene in *Arabidopsis thaliana* caused endogenous buildup of SA, which is crucial for stomatal closure under drought. Under low temperature stress, exogenous application of 0.5 mM SA also altered the activity of antioxidant enzymes such APX, SOD, GPOX, GSH-reductase, and GR and increased the chlorophyll fluorescence in *Z. mays* [32]. In citrus fruits under cold stress, 2.0 mM SA was found to increase phenolic accumulation and PAL activity [33]. Under heat stress, *Cucumis sativus* produced more PSII, Fv/Fm, and quantum yields of the PSII electron transport when 1.0 mM SA was added [34]. This was accomplished by reducing electrolyte leakage and oxidative stress. According to a study by Khan *et al.* [35], the exogenous injection of 0.5 mM SA reduced the stress-induced synthesis of ethylene and hence the harmful consequences of heat stress.

“Salicylic acid is not the mobile signal itself, although SAR is still shown to occur in other parts of the plant. Furthermore, recent studies indicate that the volatile SAR signal produced by the methyl ester methyl salicylate is transmitted to remote parts of the plant, including neighboring plants” [36]. “Salicylic acid carboxyl methyltransferase (16 µM) converts salicylic acid (SA) to methyl salicylate (MeSA) in plants at low SA concentrations” [37]. “Only the methylated form (MeSA) of SA has been demonstrated to travel regionally and systemically in plant tissue following pathogen infections” [38]. Therefore, MeSA was assumed to be the long-distance signaling molecule via phloem translocation, which travels from afflicted to unaffected leaves.

4. SALICYLIC ACID (SA) ROLES IN PLANT STRESS RESPONSES

The survival and development of plants are significantly impacted by a variety of stress elements in the plant growth environment, such as pathogen infection, dehydration, salt stress, etc. Plants have developed a number of stress-tolerance strategies involving the control of several cell signaling pathways [39]. Salicylic acid (SA) is essential for plants to be able to withstand stress (Fig. 2).

The direct involvement of SA in the activation of plant defense systems has been widely documented and examined (Tables 1 & 2). SA

may increase the activity and expression of antioxidant enzymes including superoxide dismutase (SOD), catalase (CAT), and peroxidase (POD) while it activates the appropriate plant defense mechanism [40]. A plant's resistance to biotic stressors brought on by pathogens, fungi, and pests can be increased while SA induces the production of active anti-pathogenic compounds [41].




Fig. 2. SA main roles in plant tolerance to stresses

(Source adapted from Khoshru et al., [39])

SA serves several functions in improving plant stress resistance, according to recent studies. The following categories may be used to group these roles of SA: immunological response [42], antioxidative defense [40], salt tolerance [43], and drought tolerance [44]. Numerous studies have shown that SA plays a key signaling role in plants' immunological responses to biotic and abiotic stress [45]. In order to increase plant tolerance to diseases, it can control cell wall strengthening [46], trigger the expression of a broad variety of disease resistance genes [47], and promote hormone synthesis [48]. SA can also increase a plant's resilience to oxidative stress. For instance, studies have demonstrated that SA can increase the production of antioxidant enzymes, improving the plant's capacity to neutralize reactive oxygen species and lessen cellular oxidative damage [49]. "SA is a crucial signaling molecule that activates responses to a variety of abiotic stressors, such

as cold, heat, drought, salt, heavy metals, and others" [50]. It also takes part in developmental signaling pathways necessary for optimal plant growth, such as blooming and senescence.

5. SALICYLIC ACID (SA) IN PLANT GROWTH AND PRODUCTIVITY

It has been demonstrated that SA has the ability to control the healthy growth and development of plants. Plants' physiological and biochemical processes are known to be affected by salicylic acid and other salicylates, which may be important in controlling plant growth and productivity. Salicylic acid plays a crucial role in plant physiological functions, such as increasing the plant's response to stress conditions (biotic and abiotic), increasing the plant's resistance to System Acquired Resistance (SAR), and stimulating or altering the internal endogenous signaling to withstand a variety of stresses. "Additionally, it has the capacity to form conjugates with certain amino acids, including proline and arginine, increasing the plant's ability to withstand environmental challenges and maintaining systemic acquired resistance. The generation of antioxidants is one of SA most significant impacts. The SA also plays a number of significant physiological activities, including promoting flowering, ion absorption, nutrition transfer, raising CO₂ representation, regulating stomatal movement, photosynthesis, gas exchange, and protein synthesis. In addition, it speeds up the production of different plant pigments and raises their levels, such as chlorophyll and carotene, and inhibits the representation of ethylene gas, which is in opposition to the function of ABA, which is responsible for the fall of leaves. Additionally, it plays a significant part in boosting metabolic rates, which helps the plant save energy by using alternate routes and changing its amount of nucleic and amino acids" [51].

When wheat seeds were treated with salicylic acid prior to planting, greater germination and seedling development were observed [69]. According to Fariduddin et al. [24], when lower amounts of salicylic acid were sprayed on Brassica juncea, the dry matter buildup was dramatically increased. Higher SA concentrations exhibited an inhibitory impact, though. Laishram et al. [6] reported significant growth and development of lentil under rainfed conditions owing to watersoluble antioxidant compound of SA which regulate the plant growth and development. The number of leaves and the

Table 1. Response of SA to plant biotic and abiotic stress

Stress Factors	Plant Species	SA Roles	References
Multiple pathogens	General plants	Activation of SAR pathways; defense response against biotrophic pathogens	Khoshru <i>et al.</i> [39]
Multiple biotic stresses	General plants	Activation of MAPK, CDPK, and other PKs; enhancement of SM biosynthesis	Anjali <i>et al.</i> [52]
Pathogens	<i>Arabidopsis</i>	Modulation of SA biosynthesis by SABC1's role as a molecular switch in balancing plant defense and growth	Liu <i>et al.</i> [53]
Bacterial pathogens	Jujube	Indication of antimicrobial activity	Aldhanhani <i>et al.</i> [54]
Insect herbivores and pathogens	General plants	Defense responses of SA induction to pathogens or aphid feeding	Musaqaf <i>et al.</i> [55]
Fungal pathogens	Apple	Significant upregulation of NPR1 and PR1; enhancement of endogenous SA; initiation of SA signaling pathway by (E)-2-Hexenal-based coating	Wang <i>et al.</i> [56]
Weeds	Crops	Alteration of crop growth by weeds through increment of SA signaling process	Horvath <i>et al.</i> [57]
Pests	Grapevines	Reduced injury by JA and SA application; reduction of number of eggs laid by <i>D. suzukii</i> females in JA- and SA-treated plants	Hussain <i>et al.</i> [58]

Abbreviations: SA (salicylic acid), SAR (systemic acquired resistance), SM (secondary metabolite), MAPK (mitogen-activated protein kinase), CDPK (calcium-dependent protein kinase), SABC1 (salicylic acid biogenesis controller 1), JA (jasmonic acid).

Table 2. Response of SA to plant abiotic stress

Stress Factors	Plant Species	SA Roles	References
Heat, cold	Citrus plants	Involvement in fruits' heat-induced cross-adaptation to chilling	Lafuente <i>et al.</i> [59]
Salt, drought	Rice	Regulation of stomatal aperture through the OsWRKY45-reactive oxygen pathway; regulation of adaptation to soil salinity and drought stress	Xu <i>et al.</i> [60]
Heavy metals	General plants	Induction of acclimatization impacts; improvement of tolerance against heavy metal stress	Rahman <i>et al.</i> [61]
	Common bean	Improvement of growth traits and photosynthetic indexes; enhancement of antioxidant enzyme activities and proline accumulation	Khalil <i>et al.</i> [62]
Heat	Maize	Significant elevation of transcripts of SA biosynthesis and signaling, and heat stress-responsive genes in SO ₂ pretreated seedlings; improvement of seedlings' thermotolerance	Li <i>et al.</i> [63]
Drought	Wheat	Substantial reduction of drought influence, and enhancement of grain yield and water use efficiency by co application of K ⁺ and SA	Munsif <i>et al.</i> [64]
Dry climatic conditions	Wheat	Attenuation of deficit irrigation, and improvement of growth and production by co-application of essential plant nutrients and SA	Alotaibi <i>et al.</i> [65]
Cadmium	Tomato	Significant reduction of cell wall Cd accumulation; changes in Cd distribution	Jia <i>et al.</i> [46]
Waterlogging stress	Soybean	Reduction of some physiological indexes by SA and KN; enhancement of antioxidant defense by SA and KN; effective improvement of ROS metabolism and waterlogging stress tolerance	Hasanuzzaman <i>et al.</i> [49]
Water deficit and nutrient deprivation	<i>Sempervivum tectorum</i> L.	Strong positive correlation between SA and some physiological indexes; possible antagonistic modulation of ABA and SA on water deficit-induced morphological changes	Villadangos <i>et al.</i> [66]
Abiotic stresses	Horticultural crops	Provision of abiotic stress tolerance; elicitation of various physiological and morphological responses to stress; regulation of stress-responsive genes' expression; direct interaction with various hormones, proteins, and enzymes involved in abiotic stress tolerance	Yang <i>et al.</i> [67]
Deficient irrigation	Potato	Significant enhancement of growth characteristics, yield components, and photosynthetic attributes	Desoky <i>et al.</i> [68]
Saline stress	General plants	Induction of saline stress tolerance	Yang <i>et al.</i> [67]

Abbreviations: SA (salicylic acid), ABA (abscisic acid), ROS (reactive oxygen species), Cd (cadmium), KN (kinetin).

fresh and dry mass per plant of wheat seedlings grown from grains soaked in a lower SA concentration (105 μ M) both rose considerably [70]. Maize's carbohydrate content was improved by the exogenous SA treatment as well [51]. In their pot experiment, Hussein *et al.* [72] sprayed salicylic acid on the leaves of wheat plants and watered them with water from the Mediterranean Sea. They found that this increased productivity resulted from improvements in all growth characteristics, including plant height, the number and area of green leaves, the diameter of the stem, and the dry weight of the stem, leaves, and plant as a whole. Additionally, the plants that were treated with SA had higher proline.

"Yield of lentil were increased significantly with seedpriming of SA @ 200ppm, as the growth promoting effect of SA which increased the level of cell division within the apical meristem of seedling root and caused higher plant growth and increased the dry matter production" [12]. SA can boost both the amount and quality of blooms. For instance, SA can affect blooming by encouraging flower buds to differentiate [73]. In addition to modulating seedling growth, SA also affects seedling dormancy and development. There is evidence from several research that salicylic acid influences plant growth and productivity. Abbas and Ibrahim, [74] reported that when SA was sprayed on *Niggella sativa* L. at different concentrations (50, 100, and 200 mg/l), it had a substantial impact on the examined attributes (growth, yield, and oil ratio). A study on *Datura stramonium* L. when sprayed with SA at 200ppm resulted that the maximum rate of all vegetative and investigated parameters (plant height, dry and vegetative content of the leaves, nitrogen and potassium, number of fruits, plant, and total yield kg/hectare) [75]. According to Sandoval-Yapiz [76], *Tagetes erecta* rooting was improved when salicylic acid levels were lower and a comparable promotion was also produced in the shoot system, increasing plant production. In rapeseed, the yield were enhanced significantly when the plants were sprayed with salicylic acid @ 200ppm [77]. By comparing the combined effects of SA, GA, Kinetin, NAA, ethrel, and chloro chloro chloride (CCC), Kumar *et al.* [78] discovered that SA and GA had a synergistic impact on blooming when compared to other hormone combinations. Salicylic acid may therefore be inferred to function as an endogenous regulator that may have an impact on plant development and production.

6. CONCLUSION

As a result of the foregoing explanation, it can be inferred that salicylic acid functions as a powerful plant growth regulator that is capable of modulating a variety of plant development responses. SA improves a plant's ability to grow and produce. Plants are significantly protected against a variety of biotic and abiotic stresses because to salicylic acid application's ability to trigger the SAR in plants. In addition to offering defense against pathogen and infection assaults, it also lessened the harmful effects caused by exposure to different abiotic stressors, such as heavy metals, temperature, water, and salinity stress, etc. Exogenous application of reduced salicylic acid concentrations improved plant growth during photosynthesis and a number of other physiological and biochemical traits. However, SA itself has the potential to put plants under a lot of stress at greater concentrations. Additionally, SA boosts the functions of the antioxidant enzyme system and can safeguard and improve the nitrate metabolism enzymes in high-stress environments.

7. FUTURE PERSPECTIVES

Salicylic acid functions physiologically in a variety of ways in plants and may be able to lessen the harmful impacts of numerous biotic and abiotic stressors. However, more work still has to be done to clarify the precise routes of its biosynthesis, whether significant or minor, as well as its mode of action and other unique and cooperative regulatory roles played by SA that have up to now eluded researchers. The study of how this plant hormone interacts and is controlled by cross-talk in harmony with other recognized phytohormones and plant growth regulators working at long range (auxins, cytokinins, gibberellins, ethylene, etc.), short range (NO, jasmonates, brassinosteroids, etc.), and very short range (ROS, H₂O₂) is also necessary. Another argument may be made that these short-range phytohormones are mostly synthesized nearby biotic infestations and then transferred systemically to function during widespread abiotic pressures. Clarifying the function of the aforementioned phytohormone in the tissue-specific differentiation and growth of plant parts throughout growth and development is also important. Mutant research and biochemical inhibitors of important enzymes and pathways may provide some light on these issues. This problem could be resolved by identifying tissue-specific concentrations when

seedlings are developing and fusing them with reporter genes or radioactive compounds. Exogenous application of this phytohormone may prove to be an effective strategy in the future for promoting plant growth and production as well as thwarting the negative effects brought on by various abiotic stressors. Future uses of this plant hormone as a management tool to help our agricultural crops adapt to the aforementioned constraints offer enormous promise for accelerating crop output potential in the near future.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. FAO. How to feed the world in 2050; 2009. Food and Agriculture Organization, Rome, Italy. Available:http://www.fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf.
2. Dimeyeva LA, Sultanova BM, Permitina VN, Islamgulova AF, Kerdyashkin AV. Specific features of the spatial distribution pattern of vegetation in the eastern part of the Zaisan Depression. *Arid Ecosystems*. 2015;5(4):249-60.
3. Sinha R, Fritschi FB, Zandalinas SI, Mittler R. The impact of stress combination on reproductive processes in crops. *Plant Sci*. 2021; 311:111007. DOI:10.1016/j.plantsci.2021.111007.
4. Verma K, Agrawal SB. Salicylic Acid: A Multifaceted Hormone. Springer; Singapore: Salicylic acid-mediated defence signalling in respect to its perception, alteration and transduction. 2017;97-122.
5. Peng Y, Yang J, Li X, Zhang Y. Salicylic acid: Biosynthesis and signaling. *Annu. Rev. Plant Biol*. 2021; 72:761-791. doi:10.1146/annurev-arplant-081320-092855.
6. Laishram B, Singh TB, Kalpana A, Wangkheirakpam M, Chongtham SK, Singh W. Effect of Salicylic Acid and Potassium Nitrate on Growth and Yield of Lentil (*Lens culinaris* L.) under Rainfed Condition. *International Journal of Current Microbiology and Applied Sciences*. 2020;9(11):2779-2791. Available:<https://doi.org/10.20546/ijcmas.2020.911.337>.
7. Raskin I, Skubatz H, Tang W, Mense BJD. Salicylic acid levels in thermogenic and non thermogenic plants. *Annals of Botany*. 1990; 66:373-376.
8. Korkmaz A, Uzunlu M, Demirkiran AR. Treatment with acetylsalicylic acid protects muskmelon seedlings against drought stress. *Acta Physiologiae Plantarum*. 2007; 29, 503-508.
9. Horvath E, Szalai S, Janda T. Induction of abiotic stress tolerance by salicylic acid signaling. *Journal of Plant Growth Regulators*. 2007; 26:290-300.
10. Senaratna T, Touchell D, Bunn E, Dixon K. Acetyl salicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *Plant Growth Regulation*. 2000;30:157-161.
11. Yildirim E, Turan M, Guvenc I. Effect of foliar salicylic acid applications on growth, chlorophyll and mineral content of cucumber (*Cucumis sativus* L.) grown under salt stress. *Journal of Plant Nutrition*. 2008; 31:593-612.
12. Laishram B, Singh TB, Devi OR, Khumukcham PS, Ngairangbam H. Yield, Economics, Nutrient uptake and Quality of Lentil (*Lens culinaris* L.) as Influence by Salicylic Acid and Potassium Nitrate under Rainfed Condition. *Environment and Ecology*. 2023;41(3A):1591-1596. Available:<https://doi.org/10.60151/envec/hdsa3286>.
13. Alvarez AL. Salicylic acid in machinery of hypersensitive cell death and disease resistance. *Plant Mol. Biol*. 2000;44:429-442.
14. Kawano T, Furuichi T, Muto S. Controlled salicylic acid levels and corresponding signaling mechanisms in plants. *Plant Biotechnol*. 2004; 21:319-335.
15. Mustafa NR, Kim HK, Choi YH, Erkelens C, Lefeber AW, Spijksma G, van der Heijden R, Verpoorte R. Biosynthesis of salicylic acid in fungus elicited *Catharanthus roseus* cells. *Phytochemistry*. 2009; 70:532-539.
16. Wildermuth MC, Dewdney J, Wu G, Ausubel FM. Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature*. 2001; 414:562-565.
17. Klessig DF, Tian M, Choi HW. Multiple targets of salicylic acid and its derivatives in plants and animals. *Front. Immunol*. 2016; 7:206. DOI:10.3389/fimmu.2016.00206.

18. Yang H, Fang R, Luo L, Yang W, Huang Q, Yang C, Hui W, Gong W, Wang J. Uncovering the mechanisms of salicylic acid-mediated abiotic stress tolerance in horticultural crops. *Front. Plant Sci.* 2023;14:1226041.
19. Vidhyasekaran P. Salicylic Acid Signaling in Plant Innate Immunity. *Plant hormone signaling systems in plant innate immunity.* Springer, Netherlands. 2015;2:27–122.
20. Joseph B, Jini D, Sujatha S. Insight into the role of exogenous salicylic acid on plants grown under salt environment. *Asian J Crop Sci.* 2010;2:226–235.
21. Chen HJ, Hou WC, Kuc J, Lin YH, Ca²⁺-dependent and Ca²⁺-independent excretion modes of salicylic acid in tobacco cell suspension culture. *J. Exp. Bot.* 2001;52:1219–1226.
22. Morris K, Mackerness SAH, Page T, John CF, Murphy AM, Carr JP, Buchanan-Wollaston V. Salicylic acid has a role in regulating gene expression during leaf senescence. *Plant J.* 2000;23:677–685.
23. Uzunova AN, Popova LP. Effect of salicylic acid on leaf anatomy and chloroplast ultrastructure of barley plants. *Photosynthetica.* 2000; 38:243–250.
24. Fariduddin Q, Hayat S, Ahmad A. Salicylic acid influences net photosynthetic rate, carboxylation efficiency, nitrate reductase activity and seed yield in *Brassica juncea*. *Photosynthetica.* 2003; 41, 281–284.
25. Medvedev SS, Markova IV. Participation of salicylic acid in gravitropism in plants. *Dokl. Akad. Nauk SSSR.* 1991; 316:1014–1016 (in Russian).
26. Srivastava MK, Dwivedi U.N. Delayed ripening of banana fruit by salicylic acid. *Plant Sci.* 2000; 158:87–96.
27. Jumali SS, Said IM, Ismail I, Zainal Z. Genes induced by high concentration of salicylic acid in *Mitragyna speciosa*. *Aust J Crop Sci.* 2011;5:296.
28. Gémes K, Poór P, Horváth E, Kolbert Z, Szopkó D, Szepesi Á, Tari I. Cross-talk between salicylic acid and NaCl-generated reactive oxygen species and nitric oxide in tomato during acclimation to high salinity. *Physiol Plant.* 2011;142:179–192.
29. Habibi G. Exogenous salicylic acid alleviates oxidative damage of barley plants under drought stress. *Acta Biol Szeged.* 2012; 56:57–63.
30. Alam MM, Hasanuzzaman M, Nahar K, Fujita M. Exogenous salicylic acid ameliorates short-term drought stress in mustard (*Brassica juncea* L.) seedlings by up-regulating the antioxidant defense and glyoxalase system. *Aust J Crop Sci.* 2013;7:1053–1063.
31. Miura K, Okamoto H, Okuma E, Shiba H, Kamada H, Hasegawa PM, Murata Y. SIZ1 deficiency causes reduced stomatal aperture and enhanced drought tolerance via controlling salicylic acid-induced accumulation of reactive oxygen species in *Arabidopsis*. *Plant J.* 2013; 73:91–104.
32. Janda T, Szalai G, Tari I, Paldi E. Hydroponic treatment with salicylic acid decreases the effects of chilling injury in maize (*Zea mays* L.) plants. *Planta.* 1999; 208:175–180.
33. Sibozza XI, Bertling I, Odindo AO. Salicylic acid and methyl jasmonate improve chilling tolerance in cold-stored lemon fruit (*Citrus limon*). *J Plant Physiol.* 2014; 171:1722–1731.
34. Shi Q, Bao Z, Zhu Z, Ying Q, Qian Q. Effects of different treatments of salicylic acid on heat tolerance, chlorophyll fluorescence, and antioxidant enzyme activity in seedlings of *Cucumis sativa* L. *Plant Growth Regul.* 2006; 48:127–135.
35. Khan MIR, Iqbal N, Masood A, Per TS, Khan NA. Salicylic acid alleviates adverse effects of heat stress on photosynthesis through changes in proline production and ethylene formation. *Plant Signal Behav.* 2013; 8:e26374.
36. Vlot A.C, Liu PP, Cameron RK, Park SW, Yang Y, Kumar D, ... and Klessig DF. Identification of likely orthologs of tobacco salicylic acid-binding protein 2 and their role in systemic acquired resistance in *Arabidopsis thaliana*. *Plant J.* 2008; 56(3), 445-456.
37. Dempsey DMA, Vlot AC, Wildermuth CM, Klessig FD. Salicylic acid biosynthesis and metabolism. *The Arabidopsis Book.* 2011;9: e0156.
38. Seskar M, Shulaev V, Raskin I. Endogenous methyl salicylate in pathogen inoculated tobacco plants. *Plant Physiol.* 1998; 116:387–392.
39. Khoshru B, Mitra D, Joshi K, Adhikari P, Rion MSI, Fadiji AE, Alizadeh M, Priyadarshini A, Senapati A, Sarikhani MR, et al. Decrypting the multi-functional biological activators and inducers of defense responses against biotic stresses in plants. *Heliyon.* 2023; 9, e13825.
40. Alam MU, Fujita M, Nahar K, Rahman A, Anee TI, Masud AAC, Amin AKMR,

- Hasanuzzaman M. Seed priming upregulates antioxidant defense and glyoxalase systems to conferring simulated drought tolerance in wheat seedlings. *Plant Stress*. 2022; 6, 100120.
41. Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA, Tyagi A, Islam ST, Mushtaq M, Yadav P, et al. Pathogenesis related proteins and peptides as promising tools for engineering plants with multiple stress tolerance. *Microbiol. Res*. 2018; 212–213, 29–37.
 42. Cui H, Qiu J, Zhou Y, Bhandari DD, Zhao C, Bautor J, Parker JE. Antagonism of Transcription Factor MYC2 by EDS1/PAD4 Complexes Bolsters Salicylic Acid Defense in Arabidopsis Effector-Triggered Immunity. *Mol. Plant*. 2018; 11:1053–1066.
 43. Fu C, Khan MN, Yan J, Hong X, Zhao F, Chen L, Ma H, Li Y, Li J, Wu H. Mechanisms of nanomaterials for improving plant salt tolerance. *Crop Environ*. 2023; 2:92–99.
 44. Chávez-Arias CC, Ramírez-Godoy A, Restrepo-Díaz H. Influence of drought, high temperatures, and/or defense against arthropod herbivory on the production of secondary metabolites in maize plants. A review. *Curr. Plant Biol*. 2022; 32: 100268.
 45. Zhang H, Liu Z, Luo R, Sun Y, Yang C, Li X, Gao A, Pu J. Genome-Wide Characterization, Identification and Expression Profile of MYB Transcription Factor Gene Family during Abiotic and Biotic Stresses in Mango (*Mangifera indica*). *Plants*. 2022;11:3141.
 46. Jia H, Wang X, Wei T, Wang M, Liu X, Hua L, Ren X, Guo J, Li J. Exogenous salicylic acid regulates cell wall polysaccharides synthesis and pectin methylation to reduce Cd accumulation of tomato. *Ecotoxicol. Environ. Saf*. 2021; 207:111550.
 47. Wang WM, Ma XF, Zhang Y, Luo MC, Wang GL, Bellizzi M, Xiong XY, Xiao SY. PAPP2C Interacts with the Atypical Disease Resistance Protein RPW8.2 and Negatively Regulates Salicylic Acid-Dependent Defense Responses in Arabidopsis. *Mol. Plant*. 2012; 5:1125–1137.
 48. Khan MSS, Ahmed S, Ikram AU, Hannan F, Yasin MU, Wang J, Zhao B, Islam F, Chen J. Phyto-melatonin: A key regulator of redox and phytohormones signaling against biotic/abiotic stresses. *Redox Biol*. 2023; 64, 102805.
 49. Hasanuzzaman M, Ahmed N, Saha T, Rahman M, Rahman K, Alam M.M, Rohman MM, Nahar K. Exogenous salicylic acid and kinetin modulate reactive oxygen species metabolism and glyoxalase system to confer waterlogging stress tolerance in soybean (*Glycine max* L.). *Plant Stress*. 2022; 3, 100057.
 50. Widhalm JR, Dudareva N. A Familiar Ring to It: Biosynthesis of Plant Benzoic Acids. *Mol. Plant*. 2015; 8:83–97.
 51. Davies PJ. The plant hormones: Their nature, occurrence and function. In: Davies PJ, editor. *Plant Hormones: Biosynthesis, Signal Transduction, Action*. 3rd ed. Dordrecht: Kluwer Academic Publishers. 2004;1-15.
 52. Anjali, Kumar S, Korra T, Thakur R, Arutselvan R, Kashyap AS, Nehela Y, Chaplygin V, Minkina T, Keswani C. Role of plant secondary metabolites in defence and transcriptional regulation in response to biotic stress. *Plant Stress*. 2023; 8:100154.
 53. Liu N, Xu Y, Li Q, Cao Y, Yang D, Liu S, Wang X, Mi Y, Liu Y, Ding C, et al. A lncRNA fine-tunes salicylic acid biosynthesis to balance plant immunity and growth. *Cell Host Microbe*. 2022; 30, 1124–1138.e1128.
 54. Aldhanhani ARH, Ahmed ZFR, Tzortzakis N, Singh Z. Maturity stage at harvest influences antioxidant phytochemicals and antibacterial activity of jujube fruit (*Ziziphus mauritiana* Lamk. and *Ziziphus spinachristi* L.). *Ann. Agric. Sci*. 2022; 67:196–203.
 55. Musaqaf N, Jørgensen HJL, Sigsgaard L. Plant resistance induced by hemipterans—Effects on insect herbivores and pathogens. *Crop Prot*. 2023;163: 106122.
 56. Wang X, Fu M, Qu X, Liu J, Bu J, Feng S, Zhao H, Jiao W, Sun F. (E)-2-Hexenal-based coating induced acquired resistance in apple and its antifungal effects against *Penicillium expansum*. *LWT* 2022; 163, 113536.
 57. Horvath DP, Clay SA, Swanton CJ, Anderson JV, Chao WS. Weed-induced crop yield loss: A new paradigm and new challenges. *Trends Plant Sci*. 2023;28:567–582.
 58. Hussain B, War AR, Pfeiffer DG. Jasmonic acid and salicylic acid induced defensive response in wine grapes against

- Drosophila suzukii* (Diptera: Drosophilidae). *Heliyon*. 2023; 9, e16505.
59. Lafuente MT, Romero P. Hormone profiling and heat-induced tolerance to cold stress in citrus fruit. *Post harvest Biol. Technol.* 2022; 194: 112088.
 60. Xu L, Zhao H, Wang J, Wang X, Jia X, Wang L, Xu Z, Li R, Jiang K, Chen Z, et al. AIM1-dependent high basal salicylic acid accumulation modulates stomatal aperture in rice. *New Phytol.* 2023; 238, 1420–1430.
 61. Rahman SU, Li Y, Hussain S, Hussain B, Khan WUD, Riaz L, Ashraf MN, Khaliq MA, Du Z, Cheng H. Role of phytohormones in heavy metal tolerance in plants: A review. *Ecol. Indic.* 2023; 146, 109844.
 62. Khalil R, Haroun S, Bassyoini F, Nagah A, Yusuf M. Salicylic acid in combination with kinetin or calcium ameliorates heavy metal stress in *Phaseolus vulgaris* plant. *J. Agric. Food Res.* 2021; 5, 100182.
 63. Li M, Wang M, Chen J, Wu J, Xia Z. Sulfur dioxide improves the thermotolerance of maize seedlings by regulating salicylic acid biosynthesis. *Ecotoxicol. Environ. Saf.* 2023; 254, 114746.
 64. Munsif F, Shah T, Arif M, Jehangir M, Afridi MZ, Ahmad I, Jan BL, Alansi S. Combined effect of salicylic acid and potassium mitigates drought stress through the modulation of physio-biochemical attributes and key antioxidants in wheat. *Saudi J. Biol. Sci.* 2022; 29:103294.
 65. Alotaibi M, El-Hendawy S, Mohammed N, Alsamin B, Al-Suhaibani N, Refay Y. Effects of Salicylic Acid and Macro and Micronutrients through Foliar and Soil Applications on the Agronomic Performance, Physiological Attributes, and Water Productivity of Wheat under Normal and Limited Irrigation in Dry Climatic Conditions. *Plants.* 2023;12:2389.
 66. Villadangos S, Munné-Bosch S. Acclimation to a combination of water deficit and nutrient deprivation through simultaneous increases in abscisic acid and bioactive jasmonates in the succulent plant *Sempervivum tectorum* L. *J. Plant Physiol.* 2023; 287:154040.
 67. Yang W, Zhou Z, Chu Z. Emerging Roles of Salicylic Acid in Plant Saline Stress Tolerance. *Int. J. Mol. Sci.* 2023; 24:3388. Available: <https://doi.org/10.3390/ijms24043388>.
 68. Desoky ESM, Selem E, El-Maati MFA, Hassn AASA, Belal HEE, Rady MM, AL-Harbi MS, Ali EF. Foliar supplementation of clove fruit extract and salicylic acid maintains the performance and antioxidant defense system of *Solanum tuberosum* L. under deficient irrigation regimes. *Horticulturae.* 2021;7:435.
 69. Shakirova, FM. Role of hormonal system in the manifestation of growth promoting and anti-stress action of salicylic acid. In: Hayat, S., Ahmad, A. (Eds.), *Salicylic Acid, A Plant Hormone*. Springer, Dordrecht, Netherlands' 2007.
 70. Hayat S, Fariduddin Q, Ali B, Ahmad A, Effect of salicylic acid on growth and enzyme activities of wheat seedlings. *Acta Agron. Hung.* 2005; 53, 433–437.
 71. Khodary SFA. Effect of salicylic acid on the growth, photosynthesis and carbohydrate metabolism in salt stressed maize plants. *Int. J. Agric. Biol.* 2004; 6:5–8.
 72. Hussein MM, Balbaa LK, Gaballah MS, Salicylic acid and salinity effects on growth of maize plants. *Res. J. Agric. Biol. Sci.* 2007; 3(4):321–328.
 73. Guo PR, Wu LL, Wang Y, Liu D, Li JA. Effects of Drought Stress on the Morphological Structure and Flower Organ Physiological Characteristics of *Camellia oleifera* Flower Buds. *Plants.* 2023; 12:2585.
 74. Abbas J, Ibrahim BA. Effect of salicylic acid in some vegetative and fruit traits of the black bean. *Iraqi Journal of Agricultural Sciences.* 2014;45(8):845-853.
 75. Al-Mohammadi AN, Al-Rawi AR. Effect of planting date, growth promoters, type of organic manure on growth and yield, and active ingredient of *Datura stramonium* L. *Tikrit Journal for Agricultural Sciences.* 2016;16(2):26- 50. The Republic of Iraq.
 76. Sandoval-Yapiz MR. 2004. Reguladores de crecimiento XXIII: efecto del ácido salicílico en la biomasa del cempazuchitl (*Tagetes erecta*). Tesis de Licenciatura. Instituto Tecnológico Agropecuario, Conkal, Yucatan, Mexico.
 77. Wangkheirakpam M, Singh TB, Laishram B, Leitam OC, ChanuRL, Kalpana A, Singh WJ. Effect of salicylic acid and boron application on growth and yield of no-tilled rape-seed (*Brassica campestris*) under rainfed condition. *The Pharma Innov.* 2020;9(11):353-358.

78. Kumar P, Lakshmi NJ, Mani VP. Interactive effects of salicylic acid and phytohormones on photosynthesis and grain yield of soybean (*Glycine max* L. Merrill). *Physiol. Mol. Biol. Plants.* 2000; 6:179–186.

© 2023 Ansari et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109071>