



# Breeding for Bacterial Wilt Disease Resistance in Brinjal

P. Pandiyaraj<sup>a++\*</sup>, T. H. Singh<sup>b#</sup>, Ajay Kumar Pandav<sup>c++</sup>,  
E. Sreegayathri<sup>d†</sup>, Arindam Das<sup>e++</sup>, S. Mudhalvan<sup>f++</sup>  
and J. Jeyaprabha<sup>f++</sup>

<sup>a</sup> Department of Horticulture, Kalasalingam School of Agriculture and Horticulture, Kalasalingam Academy of Research and Education, Virudhunagar, Tamil Nadu – 626 126, India.

<sup>b</sup> Division of Vegetable Crops, ICAR-Indian Institute of Horticultural Research, Bengaluru, Karnataka- 560 089, India.

<sup>c</sup> Lovely Professional University, Jalandhar, Punjab – 144 001, India.

<sup>d</sup> Department of Plant Pathology, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu – 641 003, India.

<sup>e</sup> Department of Vegetable and Spice Crops, Uttar Banga Krishi Viswavidyalaya, Pundibari, West Bengal – 736165, India.

<sup>f</sup> Department of Agriculture, Kalasalingam School of Agriculture and Horticulture, Kalasalingam Academy of Research and Education, Virudhunagar, Tamil Nadu – 626 126, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2024/v36i34407

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107353>

Review Article

Received: 01/08/2023

Accepted: 06/10/2023

Published: 06/02/2024

<sup>++</sup> Assistant Professor;

<sup>#</sup> Principal Scientist;

<sup>†</sup> PhD Scholar;

\*Corresponding author: E-mail: pandiyaraj.p7@gmail.com;

## ABSTRACT

Brinjal is a warm-season vegetable that belongs to the Solanaceae family. Among the biotic stresses of brinjal, bacterial wilt is a major disease caused by a soil-borne bacterium, *Ralstonia solanacearum*. Both cultural and chemical disease control methods are not effective due to the wide host range and prolonged spore survival periods in soil. In recent past, the use of resistant varieties and hybrids are recognized as the effective method to control bacterial wilt disease. Information on genetics of resistance to bacterial wilt disease is of prime importance for improvement of varieties and hybrids in brinjal. The bacterial wilt resistance response and resistance mechanisms are strictly based on the environmental conditions and cultivars. In traditional crop improvement methods, the development of varieties and hybrids is expensive and time consuming. The discovery of molecular marker systems increases the speed and precision for developing of varieties and hybrids with desired agronomic traits.

**Keywords:** Bacterial wilt; resistance source; resistance genetics; molecular markers.

## 1. INTRODUCTION

“Brinjal (*Solanum melongena* L.) is a warm-season vegetable that belongs to the solanaceae family. It is widely cultivated and freshly consumed in India and other parts of the world. Among the biotic stresses of brinjal, bacterial wilt is a major disease caused by a soil-borne bacterium, *Ralstonia solanacearum*, which reduces the yield from 11.67% to 96.67%, while the humid and congenial climatic condition recorded up to 100% loss”. Bainsla et al. [1] “The disease was first described by E.F. Smith in potato, tomato and brinjal in 1896 and afterward in tobacco in 1908. *Ralstonia solanacearum* is ranked as the second most destructive pathogen among the 10 most important bacterial species affecting both agricultural and horticultural crops” [2] “This bacterium is responsible for causing disease of more than 200 plant species from 53 different botanical families, including Solanaceous vegetable crops such as potato, tomato, brinjal, chilli and capsicum” [3].

Both cultural and chemical disease control methods such as soil fumigation, crop rotation, adjusting the date of planting and application of chemicals are not effective due to the wide host range and prolonged spore survival periods in soil. In recent past, the use of resistant varieties and hybrids are recognized as the safest, economical, and effective method to control bacterial wilt disease. Many varieties and hybrids were developed from the public and private sectors are resistant to bacterial wilt disease for many solanaceous vegetable crops.

Information on genetics of resistance to bacterial wilt disease is of prime importance for improvement of varieties and hybrids for resistant

to bacterial wilt disease in eggplant. The bacterial wilt resistance response and resistance mechanisms are strictly based on the environmental conditions of the location. The genetics of resistance to bacterial wilt can be described from the segregating populations such as F<sub>2</sub> and back crosses. Many workers have reported that various type of gene actions have been control the bacterial wilt disease resistance in Brinjal. Nevertheless, the genetics of bacterial wilt resistance in brinjal is yet unclear. In resistance breeding, the selection of parents resistant to bacterial wilt is difficult due to environmental factors, such as temperature and pH of the soil that will affect the expression of the disease. These problems can be overcome by identifying closely linked molecular markers to disease resistance loci and using the same in marker-assisted selection (MAS), which helps in rapid identification of the trait of interest in plants and development of durable, resistant cultivars.

## 2. *Ralstonia solanacearum*

The *Ralstonia solanacearum* is a gram-negative, non-spore forming rod shaped bacterium which belongs to the family  $\beta$ -proteobacteria. The optimum temperature for the growth of *Ralstonia solanacearum* was observed between 27–35°C and no growth was observed at above 40°C or below 4°C [4]. The growth of *Ralstonia solanacearum* is optimum in alkaline pH while in acidic pH growth is inhibited. The Triphenyl Tetrazolium Chloride (TTC) chemical was used to identify the virulent and avirulent colonies of *Ralstonia solanacearum* strains in the growing medium [5]. The virulent colonies have typical characteristic symptoms of pink or light red color with red center and whitish margin and avirulent

colonies are smaller in size, off-white and non-fluidal.

### 3. DISEASE SYMPTOMS

The characteristic symptoms of bacterial wilt affected brinjal shows sudden wilt at the time of flowering followed by yellowing of leaves, stunted plant growth, and vascular discoloration. The disease can be easily detected by ooze out test. The slimy ooze was observed from the infected plant by dipping a piece of cut end of stem in a test tube containing sterile distilled water [6]. The intensity of the disease depends on the environmental factors such as soil temperature, soil moisture, soil type, soil pH, host susceptibility and virulence of strains.

### 4. BACTERIAL CULTURE ISOLATION

The *Ralstonia solanacearum* infected plants were collected from the bacterial wilt infested sick plot and pathogen was confirmed by ooze out or streaming test. The bacterial ooze was observed by dipping a piece of the lower cut stem from just above the root zone in a test tube contain sterile water. After few minutes, fine streaks of milky ooze coming out from the margin of the cut portion and the ooze composed of masses of bacteria. The above test distinguishes bacterial wilt from the other wilts like Fusarium wilt and physiological wilt.

The ooze was placed on the sterile agar plates contain Triphenyl Tetrazolium Chloride (TTC), which were incubated for 48 to 72 hours at 30°C [5]. The virulent (irregular shape, white to cream colour, slimy with pink colour in the centre) isolate were selected from the single colony and sub cultured in TTC broth for mass production. After inoculation, the bacterial broth was incubated for 72 hours at 30°C and the bacterial population was adjusted to 0.7 OD at 600 nm wave length ( $10^8$  cfu/ml). The composition of

Triphenyl Tetrazolium Chloride (TTC) media is shown in Table 1.

**Table 1. Composition of Triphenyl Tetrazolium Chloride media (for 1 litre) at pH 7**

S. No.	Chemicals	Quantity
1.	Dextrose	10 g
2.	Peptone	10g
3.	Casein hydrolysate	1 g
4.	Agar	20 g
5.	Distilled Water	1 liter

### 5. ARTIFICIAL INOCULATION

Thirty days old plants were subjected to artificial inoculation with *Ralstonia solanacearum* culture. The virulent bacterial culture was inoculated by using axil puncturing or pin prick and soil drenching methods. "In axil puncturing or pin prick method, the sterile dissection needle was used for inoculation, the method followed as dipping the needle in the bacterial culture and inserting into the axil of the leaf along with a drop of bacterial inoculum and gently pressing to ensure that the inoculum is reached the vascular tissues. In soil drenching method, the plants roots were little injured with the help of sterile scalpel blade and bacterial culture was drenched 50 mL per plant" [7].

### 6. RESISTANCE TEST AND DISEASE SCORING

The wilt incidence was recorded and percentage disease incidence (PDI) was calculated for individual germplasm as per Bi-hao et al. [8] The individual germplasm was categorized with 0–5 scale (Table 2) based on the PDI value according to Hussain et al. [9]. The dead individuals were examined for ooze out test for, in which individuals were died for bacterial wilt or other diseases.

**Table 2. Disease scale for the percentage of wilt plants**

Disease response	Percentage of disease incidence
Highly resistant (HR)	Plants did not show any wilt symptom
Resistant (R)	1-20% plants wilted
Moderately resistant (MR)	21-40% plants wilted
Moderately susceptible (MS)	41-60% plants wilted
Susceptible (S)	61-80% plants wilted
Highly susceptible (HS)	More than 80% plants wilted

## 7. PERCENTAGE DISEASE INCIDENCE (PDI)

$PDI = \text{Number of wilted plants} / c * 100$

## 8. RESISTANCE SOURCES

The bacterial wilt disease resistance source of brinjal was found in *Solanum torvum* [10] Somatic hybrids of *S. melongena* and *Solanum aethiopicum* produced using protoplast electrofusion and it was found tolerant to bacterial wilt Collonnier et al., [11]. The rootstocks of *Solanum sisymbriifolium* showed resistance against bacterial wilt disease under sick plots in field conditions [12] Namisy et al. [13] reported that high levels of resistance were found in *S. sisymbriifolium*, *S. incanum*, *S. anguivi* and *S. torvum* against bacterial wilt disease. The resistance reactions in wild relatives of brinjal and the result shows that species *Solanum ferox* and *Solanum toxicarium* were immune to bacterial wilt disease.

## 9. GENETICS OF RESISTANCE

Information on genetics of resistance to bacterial wilt disease is of prime importance for improvement of varieties and hybrids for resistant to bacterial wilt disease in brinjal. The bacterial wilt resistance response and resistance mechanisms are strictly based on the environmental conditions of the location and cultivars. Zhu et al. [14] studied the genetics of resistance in segregated F<sub>2</sub> population of brinjal for bacterial wilt disease. The results revealed that single dominant gene was control the bacterial wilt resistance in brinjal. Gopalakrishnan et al. [15] studied the inheritance of resistance to bacterial wilt disease in brinjal Surya × Pusa Kranti cross. The results of the study suggested that monogenic and incomplete dominance of susceptibility over resistance control the bacterial wilt resistance. Tian et al. [16] studied the genetics of resistance to brinjal bacterial wilt disease in 49 hybrids developed through diallel mating design (4 resistant and 3 susceptible parents). The results revealed that the genetics of resistance in brinjal bacterial wilt was governed by few recessive genes and influenced by epistasis effect.

Bi-hao et al [8] studied the genetics of resistance to bacterial wilt disease in F<sub>2</sub>s and back cross generations of E-31 (highly resistance) × E-32 (highly susceptible) cross. The results revealed

that single dominant gene was governing the resistance in brinjal bacterial wilt disease. The bacterial wilt resistance genetics was studied in four F<sub>2</sub> populations of brinjal which were derived from line × tester design. The bacterial wilt disease incidence was observed and the segregation patterns of four F<sub>2</sub> populations were recorded as 3 (resistance):1 (susceptible) ratio. They identified that the single dominant gene was governing the resistance in brinjal for bacterial wilt disease [17].

## 10. MOLECULAR MARKERS FOR DISEASE RESISTANCE

In traditional crop improvement methods, the development of varieties and hybrids is expensive and time consuming. The discovery of molecular marker systems increases the speed and precision for developing of varieties and hybrids with desired agronomic traits. The markers can assist the selection of target alleles, minimize the linkage drag and reduce the number of generations in breeding program. The ideal molecular marker should be highly polymorphic and distributed throughout the genome, require little quantity of genomic DNA, co-dominant gives sufficient resolution of genetic differences, generates multiple linkage to diverse phenotypes and inexpensive [18]. Currently, the majority of DNA markers used for brinjal genetic mapping and resistance breeding are based on PCR, including RAPD, AFLP, SCAR, CAPS, SRAP, SSRs and SNPs.

Lebeau et al.[19] identified the bacterial wilt resistance gene ERs1 in recombinant inbred lines population derived from intra-specific cross between MM738 (Susceptible) × AG91- 25 (resistance) with 4 *Ralstonia solanacearum* strains of phylotype 1. The resistance gene ERs1 is resistant to only 3 strains of phylotype 1 (CMR134, PSS366 and GM11000) and resistance was broken down with virulent strain PSS4. The action of ERs1 seems to be minimally influenced by environmental factors and mostly dependent on the inoculums strain. Ge et al.[20] assessed 141 brinjal accessions to identify the SSR markers associated with nine fruit traits. The population structure analysis was performed with 105 SSR markers and the results revealed that two subgroups were present in the population and they identified 49 SSR marker associations with 8 fruit traits.

The association mapping for morphological traits of brinjal were done through genome wide

association analysis (GWA) with SNP markers. The 194 both phenotype and genotype associations were discovered, related to 30 of the 33 measured traits. The association of 79 SNP marker loci was mapped to 39 distinct chromosomal regions and distributed all 12 brinjal chromosomes [21,22]. identified the QTLs resistant to *Ralstonia solanacearum* strains in brinjal mapping population derived from intra-specific cross between MM738 (susceptible) × EG203 (resistant). The 123 doubled haploid lines were used with phylotypes I (PSS4) and III (R3598) strains of *Ralstonia solanacearum*. They identified 10 QTLs with resistant to PSS4 and 3 QTLs resistant to R3598 strains. All the QTLs were strongly influenced by the environment. The most stable QTLs bacterial wilt resistance was identified on chromosomes 3 and 6.

## 11. CONCLUSION

Information on the genetics of resistance to bacterial wilt disease is critical for improving brinjal types and hybrids. The present review has well emphasised on Breeding for bacterial wilt disease resistance in brinjal. The climatic circumstances and cultivars have a strong influence on the bacterial wilt resistance response and mechanisms. The discovery of molecular marker systems has increased the speed and precision with which varieties and hybrids with desired agronomic features can be developed.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Bainsla NK, Singh S, Singh PK, Kumar K, Singh AK, Gautam RK. Genetic behaviour of bacterial wilt resistance in brinjal (*Solanum melongena* L.) in tropics of Andaman and Nicobar Islands of India. American Journal of Plant Sciences. 2016;7(02):333.
2. Mansfield J, Genin S, Magori S, Citovsky V, Sriariyanum M, Ronald P, Dow MA, Verdier V, Beer SV, Machado MA, Toth IA. Top 10 plant pathogenic bacteria in molecular plant pathology. Molecular Plant Pathology. 2012;13(6): 614-29.
3. Álvarez B, Biosca EG, López MM. On the life of *Ralstonia solanacearum*, a destructive bacterial plant pathogen. Current research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology. 2010;267–279.
4. Kelman A. The bacterial wilt caused by *Pseudomonas aolanacearum*. A literature review and bibliography. North Carolina Agricultural Experiment Station Technical Bulletin. 1953;99:194.
5. Kelman A. The relationship of pathogenicity of *Pseudomonas solanacearum* to colony appearance in a tetrazolium medium. Phytopathology. 1954;44(12).
6. Ghosh PP, Mandal NC. Some disease management practices for bacterial wilt of potato. The Journal of Plant Protection Sciences. 2009;1(1):51-4.
7. Artal RB, Gopalakrishnan C, Thippeswamy B. An efficient inoculation method to screen tomato, brinjal and chilli entries for bacterial wilt resistance. Pest Management in Horticultural Ecosystems. 2012;18(1):70-3.
8. Bi-hao C, Jian-jun L, Yong W, Guo-ju C. Inheritance and identification of SCAR marker linked to bacterial wilt-resistance in eggplant. African Journal of Biotechnology. 2009;8(20).
9. Hussain MZ, Rahman MA, Bashar MA. Screening of brinjal accessions for bacterial wilt caused by *Ralstonia solanacearum*. Bangladesh J. Bot. 2005;34(1):53-8.
10. Gousset C, Collonnier C, Mulya K, Mariska I, Rotino GL, Besse P, Servaes A, Sihachakr D. *Solanum torvum*, as a useful source of resistance against bacterial and fungal diseases for improvement of eggplant (*S. melongena* L.). Plant Science. 2005;1;168(2):319-27.
11. Collonnier C, Fock I, Kashyap V, Rotino GL, Daunay MC, Lian Y, Mariska IK, Rajam MV, Servaes A, Ducreux G, Sihachakr D. Applications of biotechnology in eggplant. Plant Cell, Tissue and Organ Culture. 2001;65:91-107.
12. Rahman MA, Rashid MA, Hossain MM, Salam MA, Masum AS. Grafting compatibility of cultivated eggplant varieties with wild *Solanum* species. Pakistan Journal of Biological Sciences (Pakistan); 2002.

13. Namisy A, Chen JR, Prohens J, Metwally E, Elmahrouk M, Rakha M. Screening cultivated eggplant and wild relatives for resistance to bacterial wilt (*Ralstonia solanacearum*). Agriculture. 2019;9(7): 157.
14. Huawu Z, Yuangan Y, Zhimin L, Jianguo Y, Huiming C. On resistance to bacterial wilt in eggplant (*Solanum melongena*). Hunan Nong ye da xue xue bao= Journal of Hunan Agricultural University. 2004;30(3):288-9.
15. Gopalakrishnan TR, Singh PK, Sheela KB, Shankar MA, Kutty PJ, Peter KV. Development of bacterial wilt resistant varieties and basis of resistance in eggplant (*Solanum melongena* L.). Bacterial wilt Disease and the Ralstonia Solanacearum Species Complex. 2005:293-300.
16. Tian S, Wang Y, Luo Z, Li M, Chen L, Chen Y, Hong Y. Genetic analysis of resistance to bacteria wilt in eggplant (*Solanum melongena*). Southwest China Journal of Agricultural Sciences. 2007;20(4):642-5.
17. Ajjappalavara PS, Dharmatti PR, Salimath PM, Patil RV, Patil MS, Krishnaraj PU. Genetics of bacterial wilt resistance in brinjal. Karnataka Journal of Agricultural Sciences. 2010;21(3):424–427.
18. Agarwal M, Shrivastava N, Padh H. Advances in molecular marker techniques and their applications in plant sciences. Plant cell reports. 2008;27(4)617–631.
19. Lebeau A, Gouy M, Daunay MC, Wicker E, Chiroleu F, Prior P, Frary A, Dintinger J. Genetic mapping of a major dominant gene for resistance to *Ralstonia solanacearum* in eggplant. Theoretical and Applied Genetics. 2013 126:143-58.
20. Ge HY, Liu Y, Zhang J, Han HQ, Li HZ, Shao WT, Chen HY. Simple sequence repeat-based association analysis of fruit traits in eggplant (*Solanum melongena*). Genetics and Molecular Research. 2013;12(4):5651-63.
21. Portis E, Cericola F, Barchi L, Toppino L, Acciarri N, Pulcini L, Sala T, Lanteri S, Rotino GL. Association mapping for fruit, plant and leaf morphology traits in eggplant. PLoS One. 2015;10(8):e0135200.
22. Salgon S, Raynal M, Lebon S, Baptiste JM, Daunay MC, Dintinger J, Jourda C. Genotyping by sequencing highlights a polygenic resistance to *Ralstonia pseudosolanacearum* in eggplant (*Solanum melongena* L.). International Journal of Molecular Sciences. 2018;19(2): 357.

© 2024 Pandiyaraj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/107353>