



Effect of Nutrient Management on Cowpea [*Vigna unguiculata* (L.) Walp]

P.M. Sindha ^{a++*}, H.H. Patel ^{a#}, J.N. Patel ^{a++},
S.M. Bambhaniya ^{b#} and M.D. Vaishnav ^{a++}

^a Department of Agronomy, College of Agriculture, Navsari Agricultural University, Bharuch-392012, India.

^b Department of Soil Science and Agricultural Chemistry, College of Agriculture, Navsari Agricultural University, Bharuch-392012, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2024/v36i44476

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113204>

Original Research Article

Received: 20/12/2023

Accepted: 25/02/2024

Published: 05/03/2024

ABSTRACT

A field experiment was carried out at Research Farm, College of Agriculture, Bharuch, NAU, (Gujarat) during kharif 2022 to study the effect of nutrient management on growth, yield and economics of cowpea. The experiment was laid out in RBD with factorial concept with three replications. The result revealed that significantly higher plant height, number of nodules/plant, number of branches/plant, dry matter accumulation, number of pods/plant, number of seeds/pod, length of pod, seed and stover yield and quality were recorded with the soil application of 20-40-00 N-P-S kg/ha. It also gave the higher net return and B:C ratio. Significantly higher plant height, number of nodules/plant, number of branches/plant, dry matter accumulation, number of pods/plant, number of seeds/pod, length of pod, seed and stover yield, quality, net return as well as B:C ratio with a foliar application of 1% enriched banana pseudostem sap (Novel) at branching and flowering stage.

⁺⁺ M. Sc Scholer;

[#] Assistant Professor;

*Corresponding author: E-mail: 03121999pradipsindha@gmail.com;

Keywords: Cowpea; nutrient management; foliar application; soil application; yield.

1. INTRODUCTION

"Pulses play vital role in an Indian diet as vegetable protein being an essential supplement to cereal-based diet per capita. In India, pulses are grown nearly on 28.34 Mha with an annual production of 23.15 MT and average productivity of 817 kg/ha" [1]. "In Gujarat the pulses are grown nearly on 0.88 Mha (3.10 % of total pulse area under countries) with an annual production of 1.06 MT leading to average productivity of 1204 kg/ha" [1]. Cowpea [*Vigna unguiculata* (L.) Walp.] is one of the most important pulse crops in the world. Cowpea to the family *Leguminosae* with subfamily *Papilionaceae*. It is called 'Black eye pea', 'Southern pea', 'China pea', 'kharif pea', 'Marble pea', 'Chowli' and 'Lobiya'. Cowpea is one of the important *kharif* pulse crops grown for vegetable, grain, forage and green manuring. This crop is of great importance because of availability of short duration, high yielding and quick growing varieties. It has considerable promise as an alternative pulse crop in dry land farming. Being rich in protein and containing many other nutrients, it is known as 'vegetable meat'.

Cowpea, being a legume, fixes atmospheric nitrogen and improves the soil fertility. The formation of root nodules and nitrogen fixation starts at about 15-20 days after sowing. Thus, a starter dose of nitrogen may be required for boosting the initial growth stage. Nitrogen is the most important nutrient for the plant growth. The importance of phosphorus and sulphur application to cowpea increases nodulation, symbiotic nitrogen fixation, photosynthesis, early flower initiation as well as increasing the number of flowers. Urea is the nitrogenous fertilizer in amide form. Foliar spray of urea increases the yield due to different metabolic process enhanced in plant which leads to increased yield and yield components. The extract of marine macro algae viz., brown, red and green algae is known to have positive effect on growth and yield of crops. The brown algae are the most commonly used seaweeds in agriculture. Cattle urine is a good source of nitrogen, phosphate, potassium, calcium, magnesium, chlorite and sulphate. It contains 95% water, 2.5% urea, 2.5% others (mineral salts, hormones and enzyme). Novel organic liquid nutrient is a product of Navsari Agricultural University which was patented in 2012. It is prepared from banana

pseudostem sap. N:P:K (19:19:19) is water soluble fertilizer with an optimum combination of Nitrogen, Phosphorus and Potassium. It is readily soluble in water and is best for drip irrigation and foliar application of fertilizer.

2. MATERIALS AND METHODS

The field experiment was carried out at College Farm, Navsari Agricultural University, Bharuch (Gujarat) during *kharif* season of 2022. The soil was clayey in texture and slightly alkaline in reaction. The soil was low in available N (234 kg/ha), low in available P₂O₅ (26 kg/ha), low in available sulphur (7.74 mg/kg) and high in available K₂O (287 kg/ha). The field experiment was laid out in RBD with factorial concept with 18 treatment combinations consisting of two factors viz., soil application [F₁: 20-40-00 (N-P-S kg/ha), F₂: 10-20-10 (N-P-S kg/ha) and F₃: 20-40-20 (N-P-S kg/ha)] and foliar spray [S₁: Control (Water), S₂: 1% Urea, S₃: 1% Cow urine, S₄: 1% Enriched banana pseudostem sap (Novel), S₅: 1% 19:19:19 (N:P:K) and S₆: 1% Sea weed extract] with three replications. Cowpea variety GC-5 was sown in line at 45 x 10 cm spacing. The crop was fertilized as per treatment. Nitrogen, phosphorus and sulphur were applied in the form of urea, DAP and element sulphur, respectively. Seeds were treated with *Rhizobium* and PSB strain of Navsari isolate (10 ml/kg seed) in treatment as common practices. While, foliar application was applied at branching and flowering stages as per treatment. For all the growth and development studies during the crop growth period, five plants were selected randomly from net plot and tagged in each plot for recording plant height, number of branches/plant, dry matter accumulation/plant, number of root nodules/plant, number of pod/plant, number of seeds/pod and pod length. 100 seeds were randomly taken from the bulk produce of each net plot and were counted and weighed. The weight was expressed as 100-seed weight in grams. The data on seed and stover yield were recorded from net plot and converted on hectare basis. The net realization and B:C ratio were calculated. The data recorded for various parameters during the course of investigation were statistically analysed by a producer appropriate to the design of experiment as described by Panse and Sukhatme [2]. The significance of difference was tested by "F" test at 5 per cent level.

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant height (cm)

The results (Table 1) indicated that significantly ($P < 0.05$) highest plant height at harvest (54.92 cm) was recorded at the application rate of 20-40-20 (N-P-S kg/ha) (F_3) which was statistically at par with treatment 20-40-00 (N-P-S kg/ha) (F_1). The increase in plant height might be due to availability of nutrients in balanced form especially nitrogen and phosphorus. Nitrogen is an important part of the compounds that regulate plant growth and development. Increased supply of available phosphorus play an important role in growth of new tissue and division of cells. Sulphur is essential for many growth functions in plants including nitrogen metabolism, enzyme activity and protein synthesis which increase growth of plants. The result was in conformity with the finding of Jat et al. [3], Upadhyay and Singh [4], Balai et al. [5] and Nadeem et al. [6]. Significantly taller plant at harvest (57.26 cm) was recorded under foliar application of 1% enriched banana pseudostem sap (Novel) (S_4) which was statistically at par with foliar application of 1% sea weed extract (S_6). This might be due to foliar application of nutrients at a critical crop growth stage (branching or flowering stage) which helps in promoting the cell division and multiplication as well as cell elongation and development of leaves and other plant parts. Banana pseudostem sap contains auxins and cytokinins, which are plant growth hormones. These growth-promoting substances are effectively absorbed by the cowpea plants through foliar spraying which possibly promote plant height. Similar result has been reported by Singhal et al. [7], Patel [8] and Raut [9].

3.1.2 Number of root nodules/plant

Data furnished in Table 1 indicated that the soil application of different nutrients levels in cowpea was significantly ($P < 0.05$) higher number of nodules/plant (26.19) at 45 DAS was noted when crop fertilized with 20-40-20 (N-P-S kg/ha) (F_3) but, it was statistically at par with treatment 10-20-10 (N-P-S kg/ha) (F_2). Root nodules are specialized structures formed by certain plants in a symbiotic relationship with nitrogen-fixing bacteria, such as those of the genus *Rhizobium*. These nodules are responsible for biological nitrogen fixation, where atmospheric nitrogen is converted into a usable form by plants. While

nitrogen is the primary nutrient involved in the formation and functioning of root nodules. Sulphur is involved in the synthesis of certain amino acids and proteins, which are essential for nodule development and nitrogen fixation. The results were closely related with the finding of Singh et al. [10] and Dey et al. [11]. With respect to foliar nutrition, mean data presented in Table 1 that number of root nodules/plant at 45 DAS were not significantly affected due to different foliar nutrition, but maximum number of root nodules/plant (26.81) was observed with foliar application of 1% enriched banana pseudostem sap (Novel) (S_4).

3.1.3 Number of branches/plant

Data mentioned in Table 1 indicated that the significantly higher number of branches/plant at harvest (5.90) was registered when application of 20-40-20 (N-P-S kg/ha) (F_3) but, it was statistically at par with treatment 20-40-00 (N-P-S kg/ha) (F_1). The maximum number of branches was due to application of nitrogen, phosphorus and sulphur which helps in efficient utilization of nutrients, which resulted in attaining better crop canopy and simulation of root growth and increased metabolic activities. These results were in line with Jat et al. [3], Upadhyay and Singh [4], Balai et al. [5] and Nadeem et al. [6]. Significantly, higher number of branches/plant at harvest (6.19) was recorded at foliar application of 1% enriched banana pseudostem sap (Novel) (S_4). This was, however found to be statistically at par with foliar application of 1% seaweed extract (S_6). Foliar spray of enriched banana pseudostem sap (Novel) at branching and flowering stage increased branches/plant due to content cytokinins and gibberellic acids. While sea weed, extract contains cytokinins and auxin, which promote cell division and proliferation and stem elongation. These results were in conformity with those of Parmar et al. [12], Patel [8] and Raut [9].

3.1.4 Dry matter accumulation/plant

Among three different soil application, significantly higher dry matter accumulation/plant (27.14 g) at harvest was noted when crop was fertilized with 20-40-20 (N-P-S kg/ha) (F_3) but, it was found statistically at par with treatment 20-40-00 (N-P-S kg/ha) (F_1). The increase in dry matter accumulation/plant might be due to application of nitrogen, phosphorus and sulphur fertilizer in adequate amount leads to improve the root growth and development and thereby

higher uptake of nutrients provided better condition for cell division and cell enlargement resulting in increases in the photosynthetic efficiency and thus increased the production of photosynthates reflected in better growth and ultimately in higher dry matter accumulation. These findings corroborate the results reported by Jat et al. [3], Upadhyay and Singh [4], Balai et al. [5] and Nadeem et al. [6]. It is inferred from the data furnished in Table 1, which indicate that the significantly higher dry matter accumulation/plant at harvest (28.01 g) was recorded under foliar application of 1% enriched banana pseudostem sap (Novel) (S₄). It was however statistically at par with foliar application of 1% sea weed extract (S₆). Banana pseudostem sap or sea weed extract contains nutrients as well as growth-promoting substances that can be absorbed by the leaves and effectively utilized by the plants, it may contribute to increased biomass production and dry matter accumulation. Similar finding was reported by Parmar et al. [12].

3.2 Yield Attributes

3.2.1 Numbers of pods/plant

Among three different soil application, significantly highest number of pod/plant (20.19) was recorded when crop was fertilized with 20-40-20 (N-P-S kg/ha) (F₃) but, it was statistically at par with treatment 20-40-00 (N-P-S kg/ha) (F₁). The increase in number of pods/plant might be due to optimum supply of nitrogen, phosphorus and sulphur to the crop. Nitrogen and phosphorus levels resulted in higher number of branches having a greater number of pods which could be due to maximization of photosynthesis and respiration, storage of energy, cell division and elongation which ultimately leads to enhancement in seed production. This result also been reported by Jat et al. [3], Singh and Singh [13], Pargi et al. [14] and Sonloi et al. [15]. It is inferred from the data in Table 1 that the number of pods/plant (20.09) was recorded with foliar application of 1% enriched banana pseudostem sap (Novel) (S₄), which was statistically at par with foliar application of 1% sea weed extract (S₆). Foliar spray of nutrients at different crop growth stages resulted in better growth of plants associated with increased availability of nutrients might have resulted in greater translocation of photosynthates from source to sink site that favourably resulted into increased number of

pods/plant. Banana pseudostem sap contains gibberellic acid and cytokinins, while sea weed extract contains auxin and cytokinins which promotes the plant growth, which leads to increases photosynthetic activities and increasing number of pods/plant. These results were in conformity with those of Singhal et al. [7], Fernando and Karunaratna [16], Patel [8] and Raut [9].

3.2.2 Numbers of seeds/pod and pod length

It is apparent from data in Table 1 showed that number of seeds/pod and length at harvesting under different nutrients levels as soil application was found significant. Significantly higher numbers of pods/plant (11.65) and pod length (13.11 cm) were recorded when crop fertilized with 20-40-20 (N-P-S kg/ha) (F₃) but, found statistically at par with those of treatment 20-40-00 (N-P-S kg/ha) (F₁). It may be due to the fact that application of nitrogen, phosphorus and sulphur increased seeds/pod and pod length of cowpea. Nitrogen plays a vital role in photosynthesis, promotes vegetative growth and can contribute to increased pod formation and seeds/pod. These results are in conformity with those of Jat et al. [3], Singh and Singh [13], Pargi et al. [14] and Sonloi et al. [15]. The data outlined in Table 1 clearly indicates that treatment with foliar application of 1% enriched banana pseudostem sap (Novel) (S₄) recorded significantly higher seeds/pod (12.00) and pod length was not significantly affected due to different foliar nutrition. This might be due to enhanced photosynthetic activity and higher uptake of nutrients leading to increased plant dry matter production in the pod-setting phase which might have improved the pod development. The results strongly support findings by Fernando and Karunaratna [16], Patel [8] and Raut [9].

3.2.3 100 Seed weight (g)

A perusal of data narrated in Table 1 showed that 100 seed weight was found non-significant due to different nutrients levels as soil application, but maximum 100 seed weight (8.62 g) were recorded when crop was fertilized with 20-40-20 (N-P-S kg/ha). The 100 seed weight was influenced non significantly by foliar nutrition, but maximum 100 seed weight (8.80 g) was recorded with foliar application of 1% enriched banana pseudostem sap (Novel) (S₄).

Table 1. Effect of different nutrients management treatments on growth and yield attributes of cowpea

Treatment	Plant height (cm)	Number of root nodules/plant	Number of branches/plant	Dry matter accumulation/plant	Number of pods/plant	Number of seeds/pod	Length of pod (cm)	100 seed weight (g)
Soil application (F)								
F ₁ : 20-40-00 (N-P-S kg/ha)	52.01	23.14	5.77	25.67	19.54	11.09	12.54	8.30
F ₂ : 10-20-10 (N-P-S kg/ha)	49.48	24.64	5.35	23.95	18.30	10.86	12.04	8.08
F ₃ : 20-40-20 (N-P-S kg/ha)	54.92	26.19	5.90	27.14	20.19	11.65	13.11	8.62
S.Em. ±	1.44	0.66	0.15	0.79	0.51	0.20	0.28	0.15
CD at 5%	4.13	1.91	0.44	2.27	1.46	0.56	0.82	NS
Foliar spray application (S)								
S ₁ : Control (Water)	48.07	23.06	5.25	23.90	17.80	10.36	11.92	7.96
S ₂ : 1% Urea	50.84	23.72	5.44	24.59	18.80	11.11	12.35	8.11
S ₃ : 1% Cow urine	50.75	24.16	5.48	24.47	18.51	11.01	12.28	8.19
S ₄ : 1% Enriched banana pseudostem sap (Novel)	57.26	26.81	6.19	28.01	21.09	12.00	12.93	8.80
S ₅ : 1% 19:19:19 (N:P:K)	51.36	24.10	5.55	24.71	18.98	11.16	13.13	8.69
S ₆ : 1% Sea weed extract	54.52	26.10	6.13	27.85	20.89	11.56	12.76	8.25
S.Em. ±	2.03	0.94	0.22	1.12	0.72	0.28	0.40	0.21
CD at 5%	5.85	NS	0.62	3.21	2.07	0.79	NS	NS
Interaction (F x S)								
S.Em. ±	3.52	1.62	0.37	1.93	1.25	0.49	0.69	0.30
CD at 5%	NS	NS	NS	NS	NS	NS	NS	NS
CV%	11.71	11.43	11.45	13.11	11.16	7.39	9.58	7.67

Table 2. Effect of different nutrients management treatments on yield, quality and economics of cowpea

Treatment	Grain yield (kg/ha)	Stover yield (kg/ha)	Protein content (%)	Cost of cultivation (₹/ha)	Net returns (₹/ha)	B: C ratio
Soil application (F)						
F ₁ : 20-40-00 (N-P-S kg/ha)	1034	1801	22.33	35730	47657	2.33
F ₂ : 10-20-10 (N-P-S kg/ha)	1000	1725	22.79	36170	44340	2.23
F ₃ : 20-40-20 (N-P-S kg/ha)	1099	1895	23.21	38841	49625	2.28
S.Em. ±	27.43	46.92	0.28	-	-	-
CD at 5%	79	135	NS	-	-	-
Foliar spray application (S)						
S ₁ : Control (Water)	947	1709	22.13	34303	42653	2.24
S ₂ : 1% Urea	1024	1738	22.76	34363	47814	2.39
S ₃ : 1% Cow urine	1020	1740	22.59	34403	47556	2.38
S ₄ : 1% Enriched banana pseudostem sap (Novel)	1151	2005	23.24	35603	57233	2.61

Treatment	Grain yield (kg/ha)	Stover yield (kg/ha)	Protein content (%)	Cost of cultivation (₹/ha)	Net returns (₹/ha)	B: C ratio
S₅: 1% 19:19:19 (N:P:K)	1039	1808	22.93	36003	47797	2.33
S₆: 1% Sea weed extract	1083	1843	23.01	40303	46691	2.16
S.Em. ±	38.80	66.36	0.39	-	-	-
CD at 5%	111	191	NS	-	-	-
Interaction (F x S)						
S.Em. ±	67.19	114.93	0.67	-	-	-
CD at 5%	NS	NS	NS	-	-	-
CV%	11.15	11.02	5.14	-	-	-

Table 3. Effect of different nutrients management treatments on soil fertility status

Treatment	pH	EC (dS/m)	OC (%)	Available nutrient (kg/ha)			S (mg/kg)
				N	P ₂ O ₅	K ₂ O	
Soil application (F)							
F₁: 20-40-00 (N-P-S kg/ha)	7.78	0.301	0.301	245	29.2	295	7.42
F₂: 10-20-10 (N-P-S kg/ha)	7.82	0.305	0.294	230	28.0	291	7.83
F₃: 20-40-20 (N-P-S kg/ha)	7.68	0.296	0.310	247	30.5	303	8.14
S.Em. ±	0.08	0.003	0.003	4.71	0.54	5.04	0.16
CD at 5%	NS	NS	0.010	13	1.56	NS	0.46
Foliar spray application (S)							
S₁: Control (Water)	7.82	0.308	0.291	229	27.8	284	7.56
S₂: 1% Urea	7.77	0.293	0.299	245	28.8	301	7.76
S₃: 1% Cow urine	7.74	0.310	0.296	234	28.2	289	7.70
S₄: 1% Enriched banana pseudostem sap (Novel)	7.68	0.294	0.311	250	31.1	306	7.98
S₅: 1% 19:19:19 (N:P:K)	7.80	0.301	0.302	247	29.3	294	7.87
S₆: 1% Sea weed extract	7.75	0.297	0.308	239	30.1	304	7.94
S.Em. ±	0.11	0.005	0.005	6.66	0.77	7.12	0.23
CD at 5%	NS	NS	NS	NS	NS	NS	NS
Interaction (F x S)							
S.Em. ±	0.19	0.008	0.009	11.54	1.33	12.34	0.39
CD at 5%	NS	NS	NS	NS	NS	NS	NS
CV%	4.25	4.56	4.89	8.30	7.87	7.21	8.75




Fig. 1. Seed and stover yield of cowpea as influenced by different nutrient management treatments

3.2.4 Seed and stover yield (kg/ha)

The results in Table 2 indicate that different nutrient levels as soil application exert significant effect on seed and stover yield. Among three different nutrients levels as soil application, significantly higher seed and stover yield (1099 and 1895 kg/ha, respectively) were statistically at par with treatments of 20-40-00 (N-P-S kg/ha) (F₁) (1034 and 1801 kg/ha, respectively). However, significantly the lowest seed and stover yield (1000 and 1725 kg/ha, respectively) were noted with crop fertilized with 10-20-10 (N-P-S kg/ha) (F₂). The highest seed and stover yield could be due to the improvement in yield attributes viz., number of pods/plant, number of seeds/pod, pod length and 100 seed weight. The improvement in seed yield was mainly on account of increase in the growth parameters and yield attributes of cowpea which resulted in higher production of photosynthates were utilized by the plant for development of sink under adequate supply of nutrients due to increasing in rate of nitrogen, phosphorus and sulphur fertilizer. Similar response trend was also observed by Singh et al. [17], Jat et al. [3], Singh and Singh [13] and Pargi et al. [14]. As regards to foliar spray application, significantly higher seed and stover yield (1151 and 2005 kg/ha, Respectively) of cowpea was observed under the foliar application of 1% enriched banana pseudostem sap (Novel) (S₄). This was however statistically at par with foliar application of 1% sea weed extract (S₆) (1038 and 1843 kg/ha, respectively). Moreover, significantly the lowest seed and stover yield (947 and 1709 kg/ha, respectively) were noted with the Control (S₁). The overall improvement in all the growth and yield components may be due to adequate supply of nutrients with easy availability to plant at most critical growth period resulting in better growth and yield components. The better growth of crop ultimately diverted more energy under sink source relationship which helped in providing more yield. These results were in conformity with those of Singhal et al. [7], Fernando and Karunarathna [16], Patel [8] and Raut [9].

3.3 Quality Parameters

3.3.1 Protein content in seed (%)

Protein content in seed (%) was not significantly influenced due to different nutrients levels as soil application. However, crop fertilized with 20-40-20 (N-P-S kg/ha) (F₃) registered maximum protein content (23.21 %). It is inferred from the

data furnished in Table 2 that the protein content in seed (%) was not significantly influenced due to foliar application. maximum protein content (23.24%) was registered under foliar spray application of 1% enriched banana pseudostem sap (Novel) (S₄).

3.4 Economics

Data presented in Table 2, it could be seen that the maximum net realizations and benefit: cost ratio of ₹49625/ha and 2.28, respectively with crop fertilize with 20-40-20 (N-P-S kg/ha). These were closely followed by crop fertilize with 20-40-00 (N-P-S kg/ha) (47657 ₹/ha) with B: C ratio of 2.33. The data from Table 2 reflected that maximum net return (57233 ₹/ha) was secured with foliar spray application of 1% enriched banana pseudostem sap (Novel) (S₄) along with B:C ratio of 2.61 followed by foliar spray application of 1% urea (S₂) with net return (47814 ₹/ha) and B:C ratio of 2.39.

3.5 Soil Fertility Status

The mean data given in Table 3 showed that different nutrient levels of soil ammendment did not exert significant effect on pH, electrical conductivity and available K₂O. Among three different soil application, significantly higher organic carbon, available N, P₂O₅ and S in soil (0.310%, 247 kg/ha, 30.50 kg/ha and 8.14 mg/kg) was assessed when crop fertilized with 20-40-20 (N-P-S kg/ha). Data furnished in Table 3 inferred that pH, electrical conductivity, organic carbon, available N, P₂O₅, K₂O and S in soil were assessed not significantly influenced by different foliar spray application treatments.

4. CONCLUSION

Based on the results of one year experiment, it can be concluded that cowpea crop was fertilized with 20-40-00 N-P-S kg/ha for getting higher and profitable yield. Besides, foliar spray application of 1% enriched banana pseudostem sap (Novel) at branching and flowering stage was also found remunerative.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Agricultural statistics at a glance 2020. Directorate of Economics and

- Statistics, Department of Agriculture Cooperation and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Government of India; 2021.
2. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. ICAR, New Delhi. 1985;199-200.
 3. Jat SR, Patel BJ, Shivran AC, Kuri BR, Jat G. Effect of phosphorus and sulphur levels on growth and yield of cowpea under rainfed conditions. *Annals of Plant and Soil Research*. 2013;15 (2):114-117.
 4. Upadhyay RG, Singh A. Effect of nitrogen and zinc on nodulation, growth and yield of cowpea (*Vigna unguiculata*). *Legume Research-An International Journal*. 2016;39(1):149-151.
 5. Balai RC, Meena LR, Sharma SC. Effect of different levels of nitrogen and phosphorus on cowpea [*Vigna unguiculata* (L.) Walp] under rainfed conditions of Rajasthan. *Journal of Agriculture and Ecology*. 2017;3:19- 24.
 6. Nadeem MA, Singh V, Dubey RK, Pandey AK, Singh B, Kumar N, Pandey S. Influence of phosphorus and bio-fertilizers on growth and yield of cowpea [*Vigna unguiculata* (L.) Walp.] in acidic soil of NEH region of India. *Legume Research-An International Journal*. 2018;41(5):763-766.
 7. Singhal VK, Patel GG, Patel DH, Kumar U, Saini LK. Effect of foliar application of water soluble fertilizers on growth, yield and economics of vegetable cowpea production. *An International Quarterly Journal of Environmental Science*. 2015;7:79-83.
 8. Patel RR. Effect of planting techniques and nutrient management in soybean [*Glycine max* (L.)]. Thesis M.Sc. (Agri.), (Unpublished) College of Agriculture, Bharuch, Navsari Agricultural University, Navsari, Gujarat, India; 2022.
 9. Raut PS. Effect of planting techniques and nutrient management in blackgram [*Vigna mungo* (L.)]. Thesis M.Sc. (Agri.), (Unpublished) College of Agriculture, Bharuch, Navsari Agricultural University, Navsari, Gujarat, India; 2022.
 10. Singh SP, Kumar Y, Singh S. Effect of sources and levels of sulphur on yield, quality and uptake of nutrients in greengram (*Vigna radiata*). *Annals of Plant and Soil Research*. 2017;19(2):143-147.
 11. Dey SK, Singh V, Tiwari D. Effect of phosphorus and sulphur on the growth and yield of summer mungbean (*Vigna radiata* L.). *The Pharma Innovation Journal*. 2021;10 (8):1442-1445.
 12. Parmar SK, Virdia HM, Pankhaniya RM. Response of integrated nutrient management on growth and yield of chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal*. 2022;11(7): 2841-2843.
 13. Singh R, Singh AP. Effect of phosphorus, sulphur and biofertilizer on yield, quality and uptake of nutrients in cowpea (*Vigna unguiculata*). *Annals of Plant and Soil Research*. 2017;19 (2):175-179.
 14. Pargi KL, Leva RL, Vaghasiya HY and Patel HA. Integrated nutrient management in summer cowpea (*Vigna unguiculata* L.) under south Gujarat condition. *International Journal of Current Microbiology and Applied Sciences*. 2018;7(9):1513-1522.
 15. Sonloi P, Choudhary AS, Raj S, Sonwani A, Paswan A. Effects of integrated nutrient management on green pod yield of cowpea. *International Journal of Agriculture and Plant Science*. 2022;4(1):35-39.
 16. Fernando WSLV, Karunarathna B. Effect of foliar application of banana pseudostem sap on yield of cowpea (*Vigna unguiculata* L. Walp.). *International Letters of Natural Sciences*, 2020;79:9-15.
 17. Singh AK, Tripathi PN, Singh ROOM. Effect of Rhizobium inoculation, nitrogen and phosphorus levels on growth, yield and quality of kharif cowpea [*Vigna unguiculata* (L.) Walp.]. *Crop Research Hisar*. 2007;33(1/3):71-73.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113204>