



Magnetic Resonance Imaging (MRI) Insights into Brain Morphology and Connectivity Disruptions in Schizophrenia

Samiksha Joshi ^a, Mohit Kumar Pandey ^{b++},
Bhawna Solanki ^{b++*}, Pushendra Kumar Rajput ^c,
Mame Shalini Singh ^d, Hina Praveen ^e and Manthan Gurav ^a

^aJ J Hospital, J J Marg, Nagpada-Mumbai Central, Off Jijabhoy Road, Mumbai, Maharashtra, India.

^b Medical Imaging Technology, Department of Radio-diagnosis, Goa Medical College, Goa, India.

^c Department of Paramedical Sciences Subharti Medical College, Swami Vivekanand Subharti University Meerut, India.

^d NPS Department Mangalayatan University, Beswan Aligarh, India.

^e Gautham College Affiliated to RGUHS, Bangalore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/INDJ/2024/v21i3431

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114612>

Mini-review Article

Received: 16/01/2024
Accepted: 20/03/2024
Published: 27/03/2024

ABSTRACT

This review explores the transformative impact of Magnetic Resonance Imaging (MRI) on understanding schizophrenia. Structural MRI has unveiled significant alterations in brain regions like the hippocampus and prefrontal cortex, offering detailed insights into anatomical irregularities.

⁺⁺ Assistant Professor;

*Corresponding author: E-mail: bhavna.solanki4739@gmail.com;

Functional MRI studies have shed light on disruptions within the default mode network, providing valuable insights into cognitive deficits associated with the disorder. Additionally, Diffusion Tensor Imaging has highlighted white matter abnormalities, underscoring compromised interregional communication within the brain. Despite persistent challenges such as population heterogeneity and variations in imaging protocols, MRI techniques have significantly advanced our understanding of schizophrenia's neurobiological aspects.

Keywords: Artificial intelligence; brain morphology; cognitive deficits; connectivity disruptions; magnetic resonance imaging; schizophrenia; neurobiological aspects.

1. INTRODUCTION

Magnetic Resonance Imaging (MRI) has emerged as a powerful tool in the field of neuroscience, offering unprecedented insights into the structure and function of the human brain. In recent years, MRI has played a crucial role in advancing our understanding of psychiatric disorders, including schizophrenia. This write-up delves into the applications of MRI in unraveling the intricacies of brain morphology and connectivity disruptions associated with schizophrenia [1].

- a. Functional and structural insights:** Magnetic Resonance Imaging (MRI) offers both functional and structural insights into schizophrenia, revealing alterations in brain regions and networks associated with the disorder [2].
- b. Improved understanding of connectivity:** MRI techniques elucidate disruptions in connectivity networks, such as the default mode network, contributing to a better understanding of how connectivity abnormalities relate to cognitive deficits in schizophrenia [3].
- c. Disease progression monitoring:** MRI enables longitudinal monitoring of disease progression, allowing for the assessment of changes in brain morphology and connectivity over time in individuals with schizophrenia [4].
- d. Combined with other techniques:** Integrating MRI with other imaging and analytical techniques enhances the depth of understanding of schizophrenia, providing a comprehensive view of its neurobiological underpinnings and potential treatment targets [5].

2. UNDERSTANDING SCHIZOPHRENIA

Schizophrenia is a complex and debilitating mental disorder characterized by disruptions in

thought processes, emotions, and perception of reality. Despite decades of research, the underlying neurobiological mechanisms of schizophrenia remain elusive. MRI, with its non-invasive nature and high spatial resolution, provides an avenue for investigating structural and functional abnormalities associated with this disorder [6].

3. STRUCTURAL MRI AND BRAIN MORPHOLOGY

Structural MRI allows for the visualization of anatomical structures in the brain, enabling researchers to identify alterations in morphology associated with schizophrenia. Studies using structural MRI have consistently reported abnormalities in various brain regions of individuals with schizophrenia. These include changes in the size of the hippocampus, amygdala, and the prefrontal cortex, regions implicated in emotion regulation, memory processing, and executive functions [7].

Furthermore, advances in imaging techniques, such as voxel-based morphometry (VBM) and surface-based morphometry (SBM), have enhanced our ability to detect subtle alterations in gray matter density and cortical thickness, respectively. These techniques have been instrumental in identifying region-specific abnormalities that contribute to the neurobiology of schizophrenia [8].

4. FUNCTIONAL MRI AND CONNECTIVITY DISRUPTIONS

In addition to structural changes, functional MRI (fMRI) has played a pivotal role in elucidating connectivity disruptions in the brains of individuals with schizophrenia. Resting-state fMRI, which measures spontaneous fluctuations in blood oxygen level-dependent (BOLD) signals, has been particularly useful in assessing functional connectivity alterations [9].

Table 1. Magnetic Resonance Imaging (MRI) studies into brain morphology and connectivity disruptions in schizophrenia

Aspect	Findings/Insights
Brain Morphology	1. Gray Matter Reduction: Reduced gray matter volume in various brain regions, including the prefrontal cortex and hippocampus.
	2. Enlarged Ventricles: Increased lateral and third ventricle sizes, suggesting potential atrophy and fluid-filled spaces in the brain.
Connectivity Disruptions	1. White Matter Abnormalities: Disruptions in white matter tracts, affecting interregional connectivity.
	2. Default Mode Network (DMN): Altered connectivity within the DMN, impacting self-referential and introspective processes.
	3. Fronto-Temporal Disconnectivity: Reduced connectivity between frontal and temporal lobes, affecting cognitive and emotional processing.
	4. Hippocampal Connectivity: Aberrant connections involving the hippocampus, contributing to memory and emotion regulation deficits.
	5. Thalamo-Cortical Dysconnectivity: Disrupted communication between the thalamus and cortical regions, influencing sensory processing.

Resting-state fMRI studies in schizophrenia have consistently revealed disruptions in the default mode network (DMN), which is crucial for self-referential thinking and introspection. Altered connectivity patterns within the DMN have been linked to the characteristic cognitive deficits observed in individuals with schizophrenia, such as impaired working memory and executive function [10].

5. DIFFUSION TENSOR IMAGING (DTI) AND WHITE MATTER ABNORMALITIES

To understand the connectivity disruptions at a more fundamental level, researchers have turned to diffusion tensor imaging (DTI). DTI measures the diffusion of water molecules in white matter tracts, providing information about the structural integrity of these pathways. Studies using DTI in schizophrenia have consistently reported abnormalities in white matter tracts, including the corpus callosum and the arcuate fasciculus [11].

These white matter abnormalities are thought to contribute to the disorganized communication between different brain regions, which is a hallmark of schizophrenia. The disruptions in white matter connectivity may underlie the cognitive and perceptual disturbances observed in individuals with the disorder [12].

6. CHALLENGES AND FUTURE DIRECTIONS

While MRI has significantly advanced our understanding of schizophrenia, several

challenges and avenues for future research remain. Heterogeneity within the schizophrenia population, variations in imaging protocols, and the influence of medication are factors that complicate the interpretation of findings. Additionally, longitudinal studies are essential to discern whether observed brain abnormalities are a cause or consequence of the disorder [13].

In the future, combining multimodal imaging approaches, such as integrating structural MRI, functional MRI, and DTI data, may provide a more comprehensive understanding of the complex neurobiology of schizophrenia. Moreover, advancements in machine learning and artificial intelligence can aid in the development of biomarkers for early detection and personalized treatment strategies [14,15].

7. CONCLUSION

Magnetic Resonance Imaging has revolutionized the field of neuroscience, offering unprecedented insights into the structural and functional aspects of the human brain. In the context of schizophrenia, MRI has been instrumental in uncovering abnormalities in brain morphology and connectivity disruptions. Structural MRI has highlighted alterations in key brain regions, while functional MRI has revealed changes in connectivity patterns, particularly within the default mode network.

Diffusion Tensor Imaging has provided insights into white matter abnormalities, shedding light on the disruptions in communication between

different brain regions. Despite these advancements, challenges persist, and future research should focus on addressing the heterogeneity within the schizophrenia population and incorporating longitudinal study designs.

Ultimately, the integration of various imaging modalities, coupled with advancements in analytical techniques, holds the promise of unraveling the intricate neurobiological underpinnings of schizophrenia. This knowledge is crucial for developing targeted interventions and personalized treatment strategies, ultimately improving the lives of individuals affected by this complex mental disorder.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Voineskos AN, Hawco C, Neufeld NH, Turner JA, Ameis SH, Anticevic A, Buchanan RW, Cadenhead K, Dazzan P, Dickie EW, Gallucci J, Lahti AC, Malhotra AK, Öngür D, Lencz T, Sarpal DK, Oliver LD. Functional magnetic resonance imaging in schizophrenia: Current evidence, methodological advances, limitations and future directions. *World Psychiatry*. 2024;23(1):26-51.
2. Karlsgodt KH, Sun D, Cannon TD. Structural and Functional Brain Abnormalities in Schizophrenia. *Curr Dir Psychol Sci*. 201;19(4):226-231. DOI: 10.1177/0963721410377601
3. Voineskos AN, Hawco C, Neufeld NH, Turner JA, Ameis SH, Anticevic A, Buchanan RW, Cadenhead K, Dazzan P, Dickie EW, Gallucci J, Lahti AC, Malhotra AK, Öngür D, Lencz T, Sarpal DK, Oliver LD. Functional magnetic resonance imaging in schizophrenia: current evidence, methodological advances, limitations and future directions. *World Psychiatry*. 2024;23(1):26-51. DOI: 10.1002/wps.21159
4. Sun D, Stuart GW, Jenkinson M, Wood SJ, McGorry PD, Velakoulis D, et al. Brain surface contraction mapped in first-episode schizophrenia—A longitudinal magnetic resonance imaging study. *Molecular Psychiatry*. 2009;14:976–986.
5. Arulpragasam AR, van't Wout-Frank M, Barredo J et al. Low intensity focused ultrasound for non-invasive and reversible deep brain neuromodulation – A paradigm shift in psychiatric research. *Front Psychiatry*. 2022;13:825802.
6. Parkes L, Satterthwaite TD, Bassett DS. Towards precise resting-state fMRI biomarkers in psychiatry: synthesizing developments in transdiagnostic research, dimensional models of psychopathology, and normative neurodevelopment. *Curr Opin Neurobiol*. 2020;65:120-8.
7. Green MF, Horan WP, Lee J. Nonsocial and social cognition in schizophrenia: Current evidence and future directions. *World Psychiatry*. 2019;18:146-61.
8. Gallucci J, Tan T, Schifani C et al. Greater individual variability in functional brain activity during working memory performance in schizophrenia spectrum disorders (SSD). *Schizophr Res*. 2022; 248:21-31.
9. Hawco C, Dickie EW, Jacobs G et al. Moving beyond the mean: subgroups and dimensions of brain activity and cognitive performance across domains. *Neuroimage*. 2021;231:117823.
10. Oliver LD, Hawco C, Homan P et al. Social cognitive networks and social cognitive performance across individuals with schizophrenia spectrum disorders and healthy control participants. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2021;6: 1202-14.
11. Kaczurkin AN, Moore TM, Sotiras A et al. Approaches to defining common and dissociable neurobiological deficits associated with psychopathology in youth. *Biol Psychiatry*. 2020;88:51-62.
12. Tamminga CA, Clementz BA, Pearlson G et al. Biotyping in psychosis: using multiple computational approaches with one data set. *Neuro psychopharmacology*. 2021;46: 143-55.
13. Oliver LD, Hawco C, Viviano JD et al. From the group to the individual in

- schizophrenia spectrum disorders: neuroanatomy for precision psychiatry. Biol Psychiatry. 2020;88:28-39.
- Biomarkers of social cognitive impairments and therapeutic translation. Biol Psychiatry. 2022;91:699-708.
14. Gratton C, Kraus BT, Greene DJ et al. Defining individual-specific functional 15. Chen J, Rashid B, Yu Q et al. Variability in resting state network and functional network connectivity associated with schizophrenia genetic risk: a pilot study. Front Neurosci.2018;12:114.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114612>