



# The Effect of $Mg(NO_3)_2$ Compound Fertilizer and N, P, K Fertilizer on N-Soil, N-Plant Total Contents, and Yield of Tomatoes (*Solanum lycopersicum* L.) on Inceptisols of Jatinangor

Emma Trinurani Sofyan <sup>a++\*</sup>, Rina Devnita <sup>a++</sup>  
and Yudha Ramdika <sup>b</sup>

<sup>a</sup> Department of Soil Science and Land Resources, Faculty of Agriculture, University of Padjadjaran, Jl. Raya Bandung Sumedang Km 21 Jatinangor, Indonesia.

<sup>b</sup> Agrotechnology Study Program, Faculty of Agriculture, University of Padjadjaran, Jl. Raya Bandung Sumedang Km 21 Jatinangor, Indonesia.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jaeri/2024/v25i4613>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119243>

Original Research Article

Received: 20/04/2024

Accepted: 26/06/2024

Published: 28/06/2024

<sup>++</sup> Lecturer;

\*Corresponding author: E-mail: [emma.trinurani@unpad.ac.id](mailto:emma.trinurani@unpad.ac.id);

**Cite as:** Sofyan, Emma Trinurani, Rina Devnita, and Yudha Ramdika. 2024. "The Effect of  $Mg(NO_3)_2$  Compound Fertilizer and N, P, K Fertilizer on N-Soil, N-Plant Total Contents, and Yield of Tomatoes (*Solanum Lycopersicum* L.) on Inceptisols of Jatinangor". *Journal of Agriculture and Ecology Research International* 25 (4):49-55. <https://doi.org/10.9734/jaeri/2024/v25i4613>.

## ABSTRACT

The research aimed to determine the effect of applying compound fertilizer  $Mg(NO_3)_2$  combined with N, P, and K fertilizer on N-total content in the soil, N-plant content, and tomato yield; and to obtain the best dosage for tomato yields from Inceptisols from Jatinangor. The research was carried out at the Soil Fertility and Plant Nutrition Laboratory Experiment Field and soil and plant analysis was carried out at the Soil Fertility and Plant Nutrition Laboratory, Department of Soil Science and Land Resources, Faculty of Agriculture, University of Padjadjaran, Jatinangor, Sumedang Regency, West Java during from October to December 2023. The research used a Completely Randomized Block Design (CRBD), which consisted of 6 treatments that were repeated four times, namely as follows: A = control; B = 1 dose of N, P, K recommended ( $200 \text{ kg ha}^{-1}$  Urea,  $250 \text{ kg ha}^{-1}$  SP-36, and  $100 \text{ kg ha}^{-1}$  KCl); C = 1 dose of  $Mg(NO_3)_2$  compound fertilizer; D =  $\frac{3}{4}$  dose of  $Mg(NO_3)_2$  compound fertilizer +  $\frac{3}{4}$  recommended dose of N, P, K ( $150 \text{ kg ha}^{-1}$  Urea,  $187.5 \text{ kg ha}^{-1}$  SP-36, and  $75 \text{ kg ha}^{-1}$  KCl); E = 1 dose of  $Mg(NO_3)_2$  compound fertilizer +  $\frac{3}{4}$  recommended dose of N, P, K ( $150 \text{ kg ha}^{-1}$  Urea,  $187.5 \text{ kg ha}^{-1}$  SP-36, and  $75 \text{ kg ha}^{-1}$  KCl); and F =  $1 \frac{1}{2}$  doses of  $Mg(NO_3)_2$  compound fertilizer +  $\frac{3}{4}$  recommended doses of N, P, K ( $150 \text{ kg ha}^{-1}$  Urea,  $187.5 \text{ kg ha}^{-1}$  SP-36, and  $75 \text{ kg ha}^{-1}$  KCl). The research results showed that the application of  $Mg(NO_3)_2$  compound fertilizer and N, P, K fertilizer had a significant effect on the N-total content of the soil, N-plant content, yield components, and fruit yield of tomato plants. Treatment of 1 dose of  $Mg(NO_3)_2$  compound fertilizer +  $\frac{3}{4}$  recommended dose of N, P, K fertilizer resulted in the highest N-total soil content, N-plant content, and fruit weight per plot, namely respectively 0,34%; 2.01%; and  $6.13 \text{ kg plot}^{-1}$  while the lowest was produced in treatment A (control), namely 0.13%; 0.99% and  $1.40 \text{ kg plot}^{-1}$ .

**Keywords:** Compound Fertilizer  $Mg(NO_3)_2$ ; N; P; K Fertilizer; N-total soil; N-plant; Tomatoes; Inceptisols.

## 1. INTRODUCTION

Tomatoes are one of the horticultural commodities that are widely consumed by Indonesian people. Market demand for tomatoes continues to increase from year to year along with the increase in population. This has resulted in many farmers in Indonesia developing tomato cultivation. Data on tomato production in Indonesia in 2021 is 1,114,399 tons and in 2022 it will increase to 1,168,744 tons, and in 2023 it will decrease to 1,143,780 tons [1]. Household tomato consumption in 2022 was recorded at 687,980 tons, an increase of 1.48% from 2021, namely 677,970 tons [2]. If the level of tomato consumption continues to increase without being matched by high productivity, the need for tomatoes will not be met.

To meet the needs for food in general and tomatoes in particular, the agricultural sector is required to increase its production. Increasing crop production can be achieved through various efforts, including the use of superior varieties and expanding planting areas. The Gustavi F1 tomato variety is a hybrid tomato that can adapt to the lowlands. Apart from that, this variety also has a high level of resistance to Gemini Virus and Bacterial Wilt.

Utilizing dry land is one alternative that can be done to increase crop production. One type of soil that is dominant in dry land is Inceptisol. In Indonesia, this land has a wide distribution, namely 70.52 million hectares, which means 40% of the land area in Indonesia [3-7]. Furthermore, it was stated by Nursyams and Suprihati [8] that generally, Inceptisols have low productivity, because they have low organic matter content (<5%) and also low availability of nitrogen elements. This can affect the quality of the soil and plants cannot grow optimally [9].

To overcome the low nitrogen content in Inceptisols by fertilizing, [10] stated that to achieve balanced and optimal availability of essential nutrients in the soil, it is necessary to carry out balanced fertilization. Providing fertilizer at a dose that is appropriate to the nutritional conditions of the soil will produce high productivity, apart from saving on fertilizer use and also protecting the environment.

The fertilizer used in this research is a combination of magnesium nitrate ( $Mg(NO_3)_2$ ) compound fertilizer and N, P, and K single fertilizer.  $Mg(NO_3)_2$  compound fertilizer contains macro elements, namely nitrogen (N) at 10% and magnesium (Mg) at 15%. According to Damanhuri et al. [11], the elements N and Mg

are the elements most widely used in the plant growth phase. The element N plays an important role in the formation of leaf chlorophyll which plays a role in photosynthesis, while the element Mg acts as an enzyme activator involved in the photosynthesis process [12].

The research aimed to determine the effect of applying  $Mg(CO_3)_2$  compound fertilizer combined with N, P, and K single fertilizer on the N-total content in the soil, N-plant content, and tomato yield on Inceptisols from Jatinangor; and to obtain the best dosage for tomato yields on Inceptisols from Jatinangor.

## 2. METHODOLOGY

### 2.1 Place and Time

This research was carried out at the Soil Fertility and Plant Nutrition Laboratory Experiment Field and soil and plant analysis was carried out at the Soil Fertility and Plant Nutrition Laboratory, Department of Soil Science and Land Resources, Faculty of Agriculture, Padjadjaran University, Jatinangor, Sumedang Regency, West Java during from October to December 2023.

### 2.2 Materials and Tools

The materials used are planting media in the form of Inceptisols soil from Jatinangor, Gustafi F1 tomato seeds, a mixture of cocopeat and burnt charcoal as a seeding medium, cow manure,  $Mg(CO_3)_2$  compound fertilizer, fertilizer Urea, SP-36 and KCl.

The equipment used is laboratory tools for analyzing soil and plants, calipers, analytical scales, seedling trays, shovels, rulers, zip plastic, envelopes, and stationery.

### 2.3 Experimental Design

The research used a Randomized Group Design (RAK), which consisted of 6 treatments which were repeated four times, namely as follows:

- A = Control
- B = N, P, K recommended doses (200 kg ha<sup>-1</sup> Urea, 250 kg ha<sup>-1</sup> SP-36, and 100 kg ha<sup>-1</sup> KCl)
- C = 1 Dose of  $Mg(NO_3)_2$  compound fertilizer
- D =  $\frac{3}{4}$  Compound Fertilizer Dose  $Mg(NO_3)_2$  +  $\frac{3}{4}$  N, P, K Recommendation (150 kg ha<sup>-1</sup> Urea, 187.5 kg ha<sup>-1</sup> SP-36, and 75 kg ha<sup>-1</sup> KCl)

E = 1 Dose of  $Mg(NO_3)_2$  +  $\frac{3}{4}$  N, P, K Compound Fertilizer Recommendation (150 kg ha<sup>-1</sup> Urea, 187.5 kg ha<sup>-1</sup> SP-36, and 75 kg ha<sup>-1</sup> KCl)

F = 1  $\frac{1}{2}$  Dose of  $Mg(NO_3)_2$  +  $\frac{3}{4}$  N, P, K Compound Fertilizer Recommendation (150 kg ha<sup>-1</sup> Urea, 187.5 kg ha<sup>-1</sup> SP-36, and 75 kg ha<sup>-1</sup> KCl).

## 2.4 Research Activities

Stages of research activities, namely: land and seed preparation, seeding, planting, applying fertilizer, plant maintenance (watering, weeding, pruning, and pest control), harvesting, data collection, data analysis, and reporting.

## 2.5 Data Collection

The main data collected consisted of total soil N content, plant N content, number of fruit per plant, and fruit diameter. Weight per fruit, and fruit weight per plot. Supporting data is the results of initial soil analysis before treatment. To determine the status category for soil chemical properties, an assessment was carried out based on criteria from the Bogor Soil Research Center.

## 2.6 Data Analysis

To determine the effect of applying  $Mg(NO_3)_2$  compound fertilizer and N, P, K fertilizer on total soil N content, plant N content, and tomato plant yield, analysis of variance was used and continued with Duncan's multiple range test at 5% level [13]. Data analysis using SPSS software.

## 3. RESULTS AND DISCUSSION

### 3.1 Initial Soil Analysis

The soil used in the research was the soil of the Inceptisols order from Jatinangor which had a pH = 6.52 (slightly acidic), C-organic content = 1.57% (low category); N-total = 0.16% (low category), C/N ratio = 10 (low category), CEC = 25.35 cmol. kg<sup>-1</sup> (high category), base saturation = 50.39 % (medium category),  $P_2O_5$  (Bray) = 11.10 ppm (low category),  $K_2O$  HCl 25 % = 32.20 mg.100 g<sup>-1</sup> (medium category), texture Clay is dusty because it is dominated by dust (50%) than clay (46%), and sand (4%). Based on the results of the initial soil analysis, it shows that the Inceptisols on the research land have a low level of fertility so it is necessary to fertilize to increase

soil fertility so that tomato plant growth can be optimal.

### 3.2 Soil N-Total Content

The statistical test results showed that the application of a combination of compound fertilizer  $Mg(NO_3)_2$  and single fertilizer N, P, and K had a significant effect on the N-total content of the soil. The research results are presented in Table 1.

Based on the results of the 5% DMRT test, it shows that the total N content in treatments B, C, D, E, and F is significantly different compared to A (control). The highest total N content was produced in treatment E (0.34%), followed by treatments F (0.33%), B and D (0.27%), C (.24%), and the lowest in treatment A (. 0.13%). The results of the study showed that the application of  $Mg(CO_3)_2$  compound fertilizer combined with N, P, K (E and F) fertilizer resulted in a higher soil N-total content compared to treatments B, C, and D. This situation was caused by the application N fertilizer from both  $Mg(NO_3)_2$  and Urea fertilizers can increase soil N content. As stated by Firmansyah and Sumarni [14], applying nitrogen fertilizer to the soil, especially from Urea and ZA, can increase the total N content in the soil, but tends to reduce the pH value of the soil or in other words acidify the

soil. The low total N content of the soil at the start of the experiment caused the application of N fertilizer to have a very significant effect on the total N content of the soil. As the dose of N fertilizer increases, the total N content of the soil also increases. Providing high doses of N fertilizer can cause the total N available in the soil to be higher. Furthermore, it was reported by Widiana et al. [15] that the application of compound NPK fertilizer affected the total N content of the soil, the application of compound fertilizer at 75% of the recommended dose resulted in the highest N-total soil content, but after the dose was increased to 125% and 150% The recommended dose does not increase the total N content of the soil and even tends to reduce the total N content of the soil.

### 3.3 N-Plant Content

The statistical test results showed that the combination of compound fertilizer  $Mg(CO_3)_2$  and single fertilizer N, P, and K had a significant effect on the N-plant content. The research results are presented in Table 2.

Based on the results of the 5% DMRT test, it shows that the total N content in treatments B, C, D, E, and F is significantly different compared to A (control). The highest plant N content was

**Table 1. Effect of  $Mg(NO_3)_2$  Compound Fertilizer and N, P, K Fertilizer on Soil N-total Content in Inceptisols from Jatinangor**

Code	Treatments	N-Total (%)
A	Control	0.13 <sup>a</sup>
B	1 dose N, P, K recommended	0.27 <sup>bcd</sup>
C	1 dose compound fertilizer $Mg(NO_3)_2$	0.24 <sup>b</sup>
D	$\frac{3}{4}$ dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	0.27 <sup>bc</sup>
E	1 dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	0.34 <sup>d</sup>
F	1 $\frac{1}{2}$ dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	0.33 <sup>cd</sup>

Note: Numbers followed by the same letter do not show significant differences according to Duncan's Multiple Range Test at the 5% level

**Table 2. Effect of  $Mg(NO_3)_2$  Compound Fertilizer and N, P, K Fertilizer on N-Plant Content in Inceptisols from Jatinangor**

Code	Treatments	N-Plant (%)
A	Control	0.99 <sup>a</sup>
B	1 dose N, P, K recommended	1.88 <sup>bc</sup>
C	1 dose compound fertilizer $Mg(NO_3)_2$	1.82 <sup>b</sup>
D	$\frac{3}{4}$ dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	1.79 <sup>b</sup>
E	1 dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	2.01 <sup>c</sup>
F	1 $\frac{1}{2}$ dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	1.99 <sup>c</sup>

Note: Numbers followed by the same letter do not show significant differences according to Duncan's Multiple Range Test at the 5% level

produced in treatment E (2.01%), followed by treatments F (1.99%), B (1.88%), C (1.82%), D (1.79%), and the lowest was in treatment A (0.99%). The results showed that the application of  $Mg(CO_3)_2$  compound fertilizer combined with N, P, K (E and F) fertilizer resulted in higher plant N content compared to treatments B, C, and D. The high N content of the plants was related to closely with the total N content in the soil. This is in line with the opinion of [16] that the availability of nutrients in the soil influences plant nutrient levels. The results of the research reported by Subhan et al. [17] showed that N uptake in the roots stems, and leaves at 45 days after planting did not show significant differences between treatments, but at 75 and 90 days after planting there were significant differences, on N uptake by tomato plants treated with compound fertilizer. The N content in roots, stems and leaves reached the highest value at a dose of 1,000 kg ha<sup>-1</sup>. N uptake by tomato plants at 110 days after planting from all compound fertilizer dosage treatments tended to decrease N uptake by plants compared to N uptake at 90 days after planting. Furthermore, it was reported by Widiana et al. [15] that plant N uptake in the compound NPK fertilizer treatment was significantly different from the control. The compound NPK fertilizer treatment of 75% of the recommended dose had the highest plant N uptake, namely 40.76 mg plant<sup>-1</sup>, while the lowest plant N uptake was in the control treatment, namely 20.31 mg plant<sup>-1</sup>.

The results of the research also show that there is a tendency for F treatment to reduce the N content of plants, this is due to the increasing dose of  $Mg(NO_3)_2$  compound fertilizer given causing an imbalance of nutrients in the soil solution and competition for nutrient uptake by plants. As reported by Qu et al. [18], an

excessive supply of Mg fertilizer encourages roots to release phenolic acid and prevents plants from absorbing several nutrients in different vegetable plant genotypes, although it has no effect on Mg absorption by plants.

### 3.4 Yield of Plant

The statistical test results showed that the application of a combination of compound fertilizer  $Mg(NO_3)_2$  and N, P, and K fertilizers had a significant different on the number of fruit per plant, fruit diameter, weight per fruit, and fruit weight per plot. The research results are presented in Table 3.

Based on the results of the 5% DMRT test, the effect of  $Mg(NO_3)_2$  compound fertilizer combined with N, P, and K fertilizer on the number of fruit per plant, fruit diameter, weight per fruit, and fruit weight per plot generally showed that in treatment B, C, D, E and F are significantly different compared to treatment A (control). Fertilization treatment resulted in a greater number of fruit per plant, larger fruit diameter, greater weight per fruit, and higher fruit weight per plant compared to treatment A (control). Treatment E produced the highest fruit weight per plant, namely 6.13 kg, followed by treatment D (5.15 kg), treatment B (5.02 kg), treatment F (4.82 kg), treatment C (4.22 kg), and the lowest in treatment A (1.40 kg). This is related to the large number of leaves. The number of leaves influences the increase in the rate of absorption of sunlight by the leaves which causes an increase in photosynthate yield [19]. To stimulate flower formation, leaves mobilize nitrogen elements which are then distributed to flowers and fruit [20]. The lowest number of fruit and weight was in the control treatment, namely the average number of fruit was 137.25 and the fruit weight per plot was 1.4 kg.

**Table 3. Effect of  $Mg(NO_3)_2$  Compound Fertilizer and N, P, K Fertilizer on Yield Components and Yield of Tomato Plants on Inceptisols from Jatinangor**

Treatments	Number of Fruit	Fruit Diameter (cm)	Weight per fruit (g)	Fruit Weight per plot (kg)
A Control	13,73 <sup>a</sup>	2,67 <sup>a</sup>	12,90 <sup>a</sup>	1,40 <sup>a</sup>
B 1 dose N, P, K recommended	24,38 <sup>bc</sup>	3,57 <sup>b</sup>	19,55 <sup>b</sup>	5,02 <sup>bc</sup>
C 1 dose compound fertilizer $Mg(NO_3)_2$	19,50 <sup>ab</sup>	3,60 <sup>b</sup>	29,80 <sup>c</sup>	4,22 <sup>b</sup>
D $\frac{3}{4}$ dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	25,55 <sup>bc</sup>	3,83 <sup>b</sup>	37,45 <sup>d</sup>	5,15 <sup>bc</sup>
E 1 dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	26,80 <sup>c</sup>	3,84 <sup>b</sup>	36,75 <sup>d</sup>	6,13 <sup>c</sup>
F 1 $\frac{1}{2}$ dose compound fertilizer $Mg(NO_3)_2$ + $\frac{3}{4}$ dose NPK recommended	23,98 <sup>bc</sup>	3,61 <sup>b</sup>	35,45 <sup>cd</sup>	4,82 <sup>bc</sup>

Note: Numbers followed by the same letter are not significantly different according to the DMRT test at the 5% significance level

#### 4. CONCLUSION

Based on the results of the research and discussion, it can be concluded, namely:

1. Providing  $Mg(NO_3)_2$  compound fertilizer and N, P, K fertilizer has a significant effect on the N-total content of the soil, N-plant content, yield components, and fruit yield of tomato plants.
2. Treatment of 1 dose of  $Mg(NO_3)_2$  compound fertilizer +  $\frac{3}{4}$  dose of N, P, K fertilizer produced the highest total N-total content, N-plant content, and fruit weight per plot, namely respectively 0,34%; 2.01%; and 6.13 kg plot<sup>-1</sup> while the lowest was produced in treatment A (control), namely 0.13%; 0.99% and 1.40 kg plot<sup>-1</sup>.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Ministry of Agriculture, Directorate General of Horticulture. Horticultural Roof Book 2023. Department of Agriculture of the Republic of Indonesia; 2024.
2. Agricultural Statistics Agency, Agricultural Statistics 2022. Jakarta. Center for Agricultural Data and Information Systems, Ministry of Agriculture of the Republic of Indonesia; 2022.
3. Soil and Agrochemical Research Center. Technical Guidelines for Land Evaluation for Agricultural Commodities, Department of Agriculture; 2003.
4. Sofyan, Emma Trinurani, Benny Joy, Fatharani Raidasari. Effect of Combination of Soil Improvers and N, P, K Fertilizer on Organic C, Total N, Uptake N, and Yield of Lowland Rice (*Oryza Sativa* L.) in Inceptisols from Jatinangor, West Java, Indonesia". International Journal of Plant & Soil Science 2024;36 (3):142-49. Available:https://doi.org/10.9734/ijpss/2024/v36i34409.
5. Affendi, Syafrul, Jamilah, Isril Berd, Sunadi, Widodo Haryoko, Zulman Harja Utama, and Prima Novia. The Role of Zeolite and NPK Fertilizer on Maize (*Zea Mays* L.) Growth in Inceptisol, Southern Solok District". Asian Journal of Soil Science and Plant Nutrition. 2023;9(4):95-103. Available:https://doi.org/10.9734/ajsspn/2023/v9i4195.
6. De-shui T, Ji-yun JI, Shao-wen HU, Shu-tian LI, Ping HE. Effect of long-term application of K fertilizer and wheat straw to soil on crop yield and soil K under different planting systems. Agricultural Sciences in china. 2007;6(2):200-7.
7. 20Baligar VC, Bennett OL. NPK-fertilizer efficiency—a situation analysis for the tropics. Fertilizer research. 1986;10:147-64.
8. Nursyamsi D, Suprihati. Chemical and mineralogical properties of soil and their relation to fertilizer requirements for Rice (*Oryza sativa*), Corn (*Zea mays*) and Soybeans (*Glycine max*). Indonesian Journal of Agronomy. 2005;33 (3):40-47.
9. Siswanto B, Widowati W. The effect of seaweed agar industry waste on soil chemical properties and Corn Plant Growth in Inceptisol Soil, Pandaan District, Pasuruan. Buana Science. 2018;18(1):57-66.
10. Widyastutik Firdaus M, Aminah M, Panjaitan DV. Cost benefit analysis of balanced fertilization in the context of fulfilling optimal Nutrients: RIA Approach. Journal of Development Economics and Policy. 2022;11(1):35–55.
11. Damanhuri D, Widodo T, Fauzi A. Regulation of Nitrogen and Magnesium Balance to Increase Corn Growth and Production (*Zea mays* L.). Innovation Scientific Journal. 2022;22(1):10–15.
12. Wiyantoko B, Kurniawati P, Purbaningtias TE. Testing of total nitrogen, water content and lead metal contamination in solid NPK Inorganic Fertilizer. Journal of Science and Technology. 2017;6(1).
13. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. (Translation by E, Sjamsuddin and J.S. Baharsyah). UI. Press, Jakarta; 1995.
14. Firmansyah I, Sumarni N. Effect of N Fertilizer doses and varieties on Soil pH, Soil N-Total, N Uptake, and Shallot Bulb Yield (*Allium ascalonicum* L.) in Entisols-

- Brebes Soil, Central Java. Journal of Horticulture. 2013;23(4):358–364.
15. Widiana S, Yunarti A, Sofyan ET, Sara DS. Effect of Compound NPK Fertilizer on N-Total, N Uptake, and Yield of Shallot Bulbs (*Allium ascalonicum* L.) on Inceptisols from Jatinangor. Soilrens. 2020;18(1):50-56
  16. Hardjowigeno S. Soil Science. Publisher: Akademika Pressindo, Jakarta; 2015.
  17. Subhan N, Nurtika N. Gunadi. Response of Tomato Plants to the Use of NPK 15-15-15 Compound Fertilizer on Latosol Soil in the Dry Season. J. Horticulture. 2009;19(1):40-48.
  18. Qu SH, Li X, Zhang J, Gao R, Ma Ma J, Ma. Effect of magnesium imbalance on root growth and nutrient uptake in various genotypes of vegetable plants. PubMed Central. 2023;12(20) 3518. DOI: 10.3390/plants12203518
  19. Ross CW, Salisbury FB. Plant Physiology Volume 1. ITB, Bandung; 1995.
  20. Djarwatiningsih D, Suwandi S, Guniarti G, Wardani W. Response of Tomato Plants (*Lycopersicum esculentum*, Mill), Due to Application of Urea and Mamigro Foliar Fertilizer on Plant Growth and Yield. Agritrop: Journal of Agricultural Science. 2018;16(2):211-216

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/119243>