



Influence of Major Abiotic Factors on the Population Trend of *Tetranychusurticae* Koch on Okra at Samastipur, Bihar, India

Surendra Prasad ^{a*} and Swetalin Gouda ^a

^a Department of Entomology, PGCA, Rajendra Prasad Central Agricultural University, Pusa, Samastipur, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i114562>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125982>

Original Research Article

Received: 27/08/2024

Accepted: 30/10/2024

Published: 04/11/2024

ABSTRACT

Field experiments were conducted to study the population fluctuation of the phytophagous mite (*Tetranychus urticae* Koch) in okra and its relation with different abiotic factors under field conditions during summer 2023 at Rajendra Prasad Central Agricultural University, Pusa, Samastipur, Bihar. The population of *T. urticae* in okra commenced from the first fortnight of March and reached its peak of 34.60 mites/ 2.5cm² leaf areas on the first fortnight of June. There was a positive and significant correlation of the population of *T. urticae* with maximum temperature ($r = 0.779^{**}$) and minimum temperature ($r = 0.699^*$).

*Corresponding author: E-mail: spasad@rpcau.ac.in, spd06_bhu@rediffmail.com;

Keywords: Correlation; okra; seasonal incidence; tetranychidae.

1. INTRODUCTION

Okra, *Abelmoschus esculentus* (L.) Moench (family: Malvaceae), is one of the world's oldest, traditional cultivated vegetables. Its primary centre of origin in Africa probably Ethiopia and is grown in many Asian countries and other parts of the world. Within India, its cultivation occurs during both the summer and rainy seasons. It is commonly farmed as a summer crop in Northern India and is also prevalent as a winter crop in states like Gujarat, Andhra Pradesh, Karnataka, and Tamil Nadu. (Lal and Sinha, 2005). In India okra crop is cultivated in an area of 0.531 million hectares and its production is 6.47 million tons (Anonymous, 2020-21).

"The low yield of okra is attributed to the attack of different pests from sowing to harvesting. Recently, there has been a change in agricultural scenario and mites are becoming serious pests in most crops and okra is no exception. The spider mite, *Tetranychus urticae* Koch, poses a serious threat to okra crop particularly during spring, summer and post rainy seasons. This crop is infested mainly by six different mite pest species, viz., *T. urticae*, *T. macfarlanei*, *T. ludeni*, *Brevipalpus phoenicis* (Gij.), *Polyphagotarsonemus latus* (Bank) and *Aceria lycopersici* (Wolff.)", (Gupta, 1985; Prasad and Singh, 2011). "Out of these mite species, *T. urticae* is responsible for causing the loss of foliage of the crop plant resulting in a reduction of the economic yield of fruits ranging from 20-45% depending upon cropping season and agro-climatic conditions. *T. urticae* is well adapted to various environmental conditions, causing loss of quality and yield or death of plants by sucking out the contents of leaf cells" (Mondel and Ara, 2006, Kumaran et al., 2007).

"*T. urticae* causes direct damage in terms of loss of chlorophyll, stunting of growth, stippling, webbing, leaf yellowing, defoliation, leaf burning, reduction in size and quality of fruits, the appearance of various types of plant deformities, followed by death etc. which severely affect the yield and in extreme outbreaks, plant death. Indirect effects of mite feeding may include decreased photosynthesis and transpiration. Due to high reproductive potential and extremely short life cycle, combined with frequent acaricide applications this mite has developed resistance to almost all conventional pesticides in vogue" (Chiasson et al., 2004; Van Leeuwen et al.,

2005). "The mites become serious pests because they have several generations per season. Phytophagous nature, high reproductive potential and short life cycle contributed to rapid resistance development to many acaricides even after few applications" (Devine et al., 2001; Stumpf and Nauen, 2001). Since the degree of incidence of red spider mites changes with the season, it is desirable to have a thorough understanding of the seasonal incidence of the mite, which will lead to the development of suitable management programs. Hence, an attempt was made to correlate the effect of weather factors on the incidence and population dynamics of the spider mite in okra.

2. MATERIALS AND METHODS

Field trials were conducted from the first fortnight of March 2023 to second fortnight of July during the summer, 2023 at the Vegetable Research plot of Dr Rajendra Prasad Central Agricultural University, Pusa, Bihar. Okra cultivar of the variety "Kashi Kranti" was sown on 3rd February 2023. The plot selected had uniform topography with well-drained sandy loam soil. The latitude, longitude and altitude of the place are 85.67° E 25.98° N and 52.92 m respectively. The climate in this region is hot dry summers with mild winters. Other climatic factors such as rainfall, temperature and humidity in this region are highly favourable for the growth of vegetables and other crops in all seasons with the proliferation of mites infesting a variety of crops.

A bulk-plot was raised to conduct research on the population fluctuation of major mites associated with okra. The crop was cultivated by following appropriate agricultural practices and managed without the application of any insecticide. The population of mites was recorded at fortnight intervals. Five places were selected for each host and three leaves were plucked randomly from host plants for recording population. All the leaves were mixed, labeled, and brought to the laboratory of the Department of Entomology, Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar for counting the mites. Care was taken to avoid disturbance to the natural population of mites on the leaf. All the mites were counted under a stereo binocular microscope. The whole population was recorded on 2.5cm² areas at four spots per leaf and the mean population was recorded. The meteorological data were taken from the

meteorological section, Dr Rajendra Prasad Central Agricultural University, Pusa, Bihar. After that, these abiotic parameters were correlated with the corresponding pest population. For testing the significance of the simple correlation coefficient, the 't' test was applied. The multiple regression coefficients and multiple correlation coefficient (R) of the mite population in relation to abiotic factors were also computed. The significance of all the multiple correlation coefficients (R) was tested for their significance by applying the 'F' test. The value of the coefficient of determination (R²) was calculated to highlight the joint contribution of independent variables on mite population.

3. RESULTS AND DISCUSSION

The population of *T. urticae* was recorded on okra in different months from the 1st fortnight of March to the 2nd fortnight of July in 2023 is presented in Table 1 and Fig 1. The *T. urticae* population started increasing from the 2nd fortnight of March and reached its peak during the summer months. The maximum population of *T. urticae* was recorded in June 1st fortnight (34.57) during 2023 when the maximum temperature, minimum temperature, relative humidity maximum and minimum relative humidity, and total rainfall were 40.26°C, 23.88°C, 71.66%, 34.20%, 0.00 mm, respectively. The populations of *T. urticae* was gradually reduced from the June 2nd fortnight to the July 1st fortnight with a decrease in temperature. The minimum incidence of *T. urticae* was recorded in March 1st fortnight (0.37).

The present findings are somewhat consistent with that of Mandal et al. (2006) who observed that red spider mite populations on okra fluctuated significantly between 2000 and 2001, peaking during the 15th standard meteorological week and decreasing until the 24th SMW. The population increased gradually from March to July, Similarly Kumar et al. (2015) observed that the population of red spider mites on okra during the crop was higher in May and the temperature was also high during the period. Veerendra et al (2015) revealed that the population of *T. urticae* in grape vineyards, began in November 2012 and increased until April 2013.

The population of *T. urticae* had a positive and significant correlation with maximum temperature (r= 0.779**) and minimum temperature (r = 0.699*) in Table 2. There was a negative and non significant relation of the *T. urticae* population with maximum relative humidity (r = - 0.375) and minimum relative humidity (r = - 0.068). The rainfall had a positive and non-significant correlation (r =0.306) with the *T. urticae* population. The regression equation for the data, with a population(Y) as the dependent variable and weather factors as the independent variables, was as follows:

$$Y = -184.644 + X_1(1.129) + X_2(4.626) - X_3(1.333) - X_4(0.810) - X_5(0.107)$$

X₁ = Maximum temperature, X₂ = Minimum temperature, X₃ = Maximum relative humidity, X₄ = Minimum relative humidity, X₅ = Rainfall.




Fig. 1. Influence of weather parameters on the population of *Tetranychus urticae* on okra crop during summer 2023

Table 1. Population fluctuation of *Tetranychus urticae* on okra during summer 2023

Interval	*Mean population of <i>T. urticae</i> mites/2.5cm ² leaf area	Temperature (°C)		Relative humidity (%)		Total Rainfall (mm)
		Maximum	Minimum	Maximum	Minimum	
1 st Fortnight March	0.37	31.11	14.67	93.33	48.67	0.00
2 nd Fortnight March	2.53	30.18	16.73	89.06	50.56	18.00
1 st Fortnight April	8.23	35.67	16.20	76.80	27.46	0.00
2 nd Fortnight April	13.43	35.80	20.73	71.06	34.27	0.00
1 st Fortnight May	30.10	35.71	19.93	75.07	37.20	1.40
2 nd Fortnight May	33.63	35.20	22.52	82.56	52.18	16.80
1 st Fortnight June	34.57	40.26	23.88	71.66	34.20	0.00
2 nd Fortnight June	29.67	36.27	25.36	86.20	55.20	92.60
1 st Fortnight July	25.30	33.22	19.70	89.20	50.40	59.40
2 nd Fortnight July	6.20	32.45	20.20	85.30	51.20	24.00

* Mean of 5 plants per plot, 3 leaves of leach plant and one leaf 2.5 cm² areas of four spots.

Table 2. Correlation coefficients and regression equation of *Tetranychus urticae* with weather parameters on okra crop during summer 2023

Correlation Factor	Temperature (°C)		Relative humidity (%)		Rainfall
	Maximum	Minimum	Maximum 700hr	Minimum 1400 hr	
Red spider mite	0.779**	0.699*	-0.375 ^{NS}	-0.068 ^{NS}	0.306 ^{NS}
Regression factor	Regression equation			Multiple R ² Value	R ² Value
Red Spider Mite	Y= -184.644 +X ₁ (1.129) + X ₂ (4.626) -X ₃ (1.333) -X ₄ (0.810) -X ₅ (0.107)			0.8751	0.7456

**Correlation is significant at the 0.01 level *Correlation is significant at the 0.05 level; X₁ - Maximum temperature, X₂- Minimum Temperature, X₃-Relative humidity 700hrs, X₄- Relative humidity 1400 hrs, X₅- Rainfall; R₂- Coefficient of Determination

Based on the calculation, the coefficient value (R^2) was derived as 0.7456 which suggests that 74% of the fluctuations in the red spider mite population were attributed to the impact of abiotic factors. In other words, the R^2 value indicates that per cent of the variation in the red spider mite population was influenced by abiotic factors.

This finding is somehow inconsistent with the results of Mohanasundaram and Sharma (2011) revealed that the population of *T. urticae* had a non-significant positive correlation with temperature (maximum, minimum and average), maximum relative humidity and weekly total rainfall while the population had a significant positive correlation with minimum and average relative humidity.

The present findings study differ slightly from the findings of Singh et al. (2018), who revealed that the mite population fluctuation showed a highly significant negative correlation with minimum temperature and a significant negative correlation with maximum temperature and rainfall.

These findings obtained are inconsistent with the results of Ghosh (2019) who revealed that the mite population exhibited a significant positive correlation with various temperature measures (maximum, minimum, and average). Additionally, there was a significant positive correlation with both minimum and average relative humidity.

4. CONCLUSION

The population of *T. urticae* was the highest during the 1st fortnight of June 2023. There was a positive significant correlation of the *T. urticae* population with maximum temperature and minimum temperature.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (Chat GPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

ACKNOWLEDGEMENT

The authors are grateful thanks to the Head, Department of Entomology and Department of Horticulture, Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar for providing

necessary facilities during the course of experimentation.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Anonymous. (2021). *Indian horticulture database*. Ministry of Agriculture, Government of India.
- Chiasson, H., Bostanian, N. J., & Vincent, C. (2004). Acaricidal properties of a chenopodium-based botanical. *Journal of Economic Entomology*, 97(5), 1373–1377.
- Devine, G. J., Barber, M., & Denholm, P. S. (2001). Incidence and inheritance of resistance to acaricides in European strains of the two-spotted spider mite (*Tetranychus urticae*) (Acari: Tetranychidae). *Pest Management Science*, 57(1), 43–448.
- Ghosh, S. K. (2019). Climate impact on red spider mite (*Tetranychus* sp. Koch) infesting eggplant (*Solanum melongena* L.) and their management using plant extracts. *Journal of Entomological Research*, 43(3), 345–350.
- Gupta, S. K. (1985). *Handbook of plant mites of India*. Sri Aurobindo Press.
- Kumar, D., Raghuraman, M., Singh, R. N., Santeswari, & Singh, J. (2014). Effect of environmental factors on the population of spider mite, *Tetranychus urticae* Koch on okra in Varanasi region. *Effect of Environmental Factors*, 4(Special Issue), 231–235.
- Kumaran, N., Douressamy, S., & Ramaraju, K. (2007). Bio-efficacy of botanicals to two spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae) infesting okra (*Abelmoschus esculentus* L.). *Pestology*, 31(9), 43–49.
- Lal, O. P., & Sinha, S. R. (2005). Impact of imidacloprid seed treatment along with some insecticidal sprayings against insect pests of okra. *Indian Journal of Entomology*, 67(4), 328–333.
- Mandal, S. K., Sattar, A., & Banerjee, S. (2006). Impact of meteorological parameters on population build-up of red spider mite in okra, *Abelmoschus esculentus* L. under North Bihar condition. *Journal of Agricultural Physics*, 6(1), 35–38.

- Mohanasundaram, A., & Sharma, R. K. (2011). Abundance of pest complex of okra in relation to abiotic and biotic factors. *Annals of Plant Protection Sciences*, 19(2), 286–290.
- Mondel, M., & Ara, N. (2006). Biology and fecundity of the two spotted spider mite, *Tetranychus urticae* Koch. (Acari: Tetranychidae) under laboratory conditions. *Journal of Life and Earth Sciences*, 1, 43–47.
- Prasad, R., & Singh, J. (2011). Status of mite pest fauna prevailing in brinjal agro-ecosystem. *Uttar Pradesh Journal of Zoology*, 31, 15–23.
- Singh, A. K., Koul, K., Shankar, U., Singh, S. K., Mondal, A., & Singh, M. (2018). Seasonal incidence and management of red spider mite, *Tetranychus urticae* Koch on okra, *Abelmoschus esculentus* (L.) Moench. *Journal of Entomology and Zoology Studies*, 6, 650–656.
- Stumpf, N., & Nauen, R. (2001). Cross-resistance, inheritance, and biochemistry of mitochondrial electron transport inhibitor-acaricide resistance in *Tetranychus urticae* (Acari: Tetranychidae). *Journal of Economic Entomology*, 94(6), 1577–1583.
- Van Leeuwen, T., Dermauw, W., Van de Veire, M., & Tirry, L. (2005). Systemic use of spinosad to control the two spotted spider mite (Acari: Tetranychidae) on tomatoes grown in Rockwool. *Experimental and Applied Acarology*, 37(1), 93–105.
- Veerendra, A. C., Udikeri, S. S., & Karabhantanal, S. S. (2015). Dynamics of two spotted red spider mite, *Tetranychus urticae* Koch (Acari: Tetranychidae) in grape vineyards and its correlation with abiotic factors and a predator. *Journal of Entomology and Zoology Studies*, 3(6), 373–376.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125982>