



Influence of Date of Planting on Total Sugars and Titratable Acidity of Strawberry Varieties in Vertical Farming

L.Shruthi Reddy ^{a++*}, M.Rajasekhar ^{b#}, V.Suchitra ^{c†},
G.Jyothi ^{d‡} and B.Naveen Kumar ^{e^}

^a College of Horticulture, Rajendranagar, SKLTSHU, Telangana, India.

^b SKLTSHU, Mulugu, Siddipet, Telangana, India.

^c Fruit Research Station, Sangareddy, Telangana, India.

^d Horticulture Research Station, Kondamallepally, Nalgonda, Telangana, India.

^e Horticulture Polytechnic College, Peddapalli, Telangana, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i115129>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126464>

Original Research Article

Received: 06/09/2024

Accepted: 08/11/2024

Published: 12/11/2024

⁺⁺ Ph. D Scholar;

[#] Professor (Horticulture);

[†] Senior Scientist (Horticulture);

[‡] Scientist (Horticulture);

[^] Assistant Professor (Soil Science);

*Corresponding author: E-mail: lattupallishruthireddy@gmail.com;

Cite as: Reddy, L.Shruthi, M.Rajasekhar, V. Suchitra, G. Jyothi, and B.Naveen Kumar. 2024. "Influence of Date of Planting on Total Sugars and Titratable Acidity of Strawberry Varieties in Vertical Farming". *International Journal of Plant & Soil Science* 36 (11):145-51. <https://doi.org/10.9734/ijpss/2024/v36i115129>.

ABSTRACT

The research using vertical towers was carried out at Centre of Excellence (COE) for vegetables and flowers, Jeedimetla, Medchal Malkajgiri district of Telangana during Rabi 2022-23 and 2023-24 in Factorial Completely Randomized Design (FCRD) evaluating four varieties V1- Nabila (Short day), V2- Flaminia (Short day), V3- Murano (Everbearing), V4-Vivara (Everbearing) and four planting dates D1- planting on October 15th D2-Planting on October 30th, D3- planting on November 15th, D4- planting on November 30th. The vertical tower was divided into three tiers, namely the top tier, middle tier and lower tier and the observations were taken accordingly. The results revealed that, the top tier of the vertical tower has exhibited higher percentage of total sugars and lower percentage of titratable acidity than subsequent tiers of the vertical tower among all the varieties, dates of planting and interactions. Among varieties, Shortday varieties V1 (Nabila) and V2 (Flaminia) has revealed highest percentage of total sugars and lowest percentage of titratable acidity than everbearing varieties V3 (Murano) and V4 (Vivara) in all the tiers of the vertical tower. Among planting dates, D2 (Planting on October 30th) has registered with highest percentage of Total sugars and lowest percentage of titratable acidity in all the tiers of the vertical tower. The interaction of V1 (Nabila) and V2 (Flaminia) with D2 (planting on October 30th) indicated highest percentage of total sugars and the lowest percentage of titratable acidity in all the tiers of the vertical tower.

Keywords: Vertical farming; strawberry; dates of planting; tier.

1. INTRODUCTION

The Strawberry (*Fragaria x ananassa* Duch.) is a member of the Rosaceae family. The edible portion of strawberry fruit is 90% and is a rich source of minerals like calcium, phosphorus, potassium, iron, sodium and vitamins A, B, C, E and antioxidants. In the world, strawberry crop has extended to all arable regions of the globe from the Arctic to the tropics with the USA (34.0%) as the leading producer followed by Turkey (8.20%), Spain (5.60%), Egypt (5.40%) and Mexico (5.30%) (FAOSTAT, 2020).

Strawberry is mainly a crop of temperate climate, however the introduction of suitable varieties lead to the cultivation of strawberry in various geographical areas. Strawberry behaves as perennial in temperate climate and as an annual in subtropical and tropical climate. Planting time within the season also influences the growth, floral production, fruit size, quality, and yield of strawberry. The qualitative evaluation of strawberry is as important as quantitative analysis as Strawberry has become an important table fruit throughout the world. It is not only consumed as fresh fruit but also has great demand in fruit processing. The quality parameters of the strawberry are significantly affected by the choice of season and date of planting which varies across different locations.

In view of the above, an experiment was done to evaluate the date of planting on total sugars and titratable acidity on strawberry.

2. MATERIALS AND METHODS

The present study was carried out at Centre of Excellence (COE) for vegetables and flowers, Jeedimetla, Medchal Malkajgiri district of Telangana during Rabi 2022-23 and 2023-24 in Factorial Completely Randomized Design (FCRD) evaluating four varieties V1- Nabila (Short day), V2- Flaminia (Short day), V3- Murano (Everbearing), V4-Vivara (Everbearing) under naturally ventilated polyhouse at four planting dates viz. D1- planting on October 15th D2-Planting on October 30th, D3- planting on November 15th, D4- planting on November 30th in the vertical towers. Each strawberry seedling was transplanted into individual pouches of the vertical tower. The vertical tower was divided into three tiers, namely the top tier, middle tier and lower tier in which each tier had eight pouches accommodating one plant in each pouch such that each tier has a total of eight plants and each vertical tower has twenty four plants.

Procedure for evaluation of total sugars (%):

Total sugars were determined by Lane and Eynon's (AOAC, 1965) method. The clarified lead free solution (50 ml) was taken into a 250 ml volumetric flask and to it 10 ml of HCl was added, mixed well and allowed to stand at room temperature for 24 hours. The solution after 24 hours was neutralized with NaOH using a drop of phenolphthalein as an indicator and volume was made up. The solution was taken into a

burette and titration was carried out against standard Fehling's solution mixture of A and B (1:1) using methylene blue as an indicator and taking brick red colour as an endpoint. The titer value is noted, and total sugars were calculated by following the formula.

$$\text{Total sugars (\%)} = \frac{\text{Factor} \times \text{Volume made up}}{\text{Titer value} \times \text{Weight of sample}} \times 100$$

Procedure for evaluation of titratable acidity (%): Ten grams of fruit pulp was taken and ground well and then transferred to a volumetric flask. The volume was made up to 100 ml in a volumetric flask. The contents were filtered through Whatman No.1 filter paper and an aliquot of 10 ml was taken into the conical flask to which 2-3 drops of phenolphthalein indicator was added and titrated against 10 N NaOH till a pink colour, as endpoint and titer value is noted. The titratable acidity was estimated in terms of per cent citric acid (Ranganna, 1986) and calculated by using the following formula.

$$\text{Titratable acidity (\%)} = \frac{\text{Titer value} \times \text{Normality of NaOH} \times 0.0064}{\text{The volume of aliquot taken}} \times 100$$

Statistical analysis: The experimental data was arranged (Two factor completely randomized design) and analyzed by using analysis of variance (ANOVA). The overall significance difference (α - 0.05) among treatments was tested using critical difference (C.D. at 5%).

3. RESULTS AND DISCUSSION

Total sugars (%): The pooled data regarding total sugars of strawberry varieties at planted at various dates in top, middle and lower tier of the vertical tower was presented in Table 1. All three tiers of the vertical tower have exhibited a significant difference among varieties, planting dates, and interactions between varieties and dates of planting in total sugars.

In the top tier, the variety V2 (Flaminia) has revealed the highest percentage of total sugars (5.51) followed by V1 (Nabila) (5.41), while, the variety V4 (Vivara) revealed the lowest percentage of total sugars (4.79). Among various dates of planting, D2 (Planting on October 30th) have recorded the highest percentage of total sugars (5.54) followed by D3

(Planting on November 15th) (5.30) and the lowest percentage of total sugars (4.87) was observed in D4 (Planting on November 30th) on par with D1 (Planting on October 15th) (4.95). In the interaction between varieties and dates of planting, the combination of V2D2 recorded the highest percentage of total sugars (6.02) and the combination of V3D1 resulted in the lowest percentage of total sugars (4.54) which was on par with V4D1 (4.69), V3D4 (4.58) and V4D4 (4.57).

In the middle tier, the variety V2 (Flaminia) has recorded the highest percentage of total sugars (5.35) followed by V1 (Nabila) (5.16), while the variety V4 (Vivara) revealed the lowest percentage of total sugars (4.61). Among various dates of planting, D2 (Planting on October 30th) have resulted in the highest percentage of total sugars (5.49) followed by D3 (Planting on November 15th) (5.14), while the lowest percentage of total sugars (4.53) was noticed in D4 (Planting on November 30th). In the interaction between varieties and dates of planting, the combination of V1D2 revealed the highest percentage of total sugars (5.77) which was on par with V2D2 (5.73) and the combination of V3D4 resulted in the lowest percentage of total sugars (4.24) which was on par with V4D4 (4.31), V1D4 (4.43), V3D1 (4.35) and V4D1 (4.28).

In the lower tier, the variety V2 (Flaminia) has exhibited the highest percentage of total sugars (5.10) followed by V1 (Nabila) (4.90), while, the variety V4 (Vivara) revealed the lowest percentage of total sugars (4.29). Among various dates of planting, D2 (Planting on October 30th) have recorded the highest percentage of total sugars (5.29) followed by D3 (Planting on November 15th) (4.94) and while, the lowest percentage of total sugars (4.40) was observed in D1 (Planting on October 15th) which was on par with D4 (Planting on November 30th) (4.41). In the interaction between varieties and dates of planting, the combination of V2D2 recorded the highest percentage of total sugars (5.62) and the combination of V4D1 resulted in the lowest percentage of total sugars (3.82).

In this experiment, it was observed that short day varieties V1 (Nabila) and V2 (Flaminia) registered with a higher percentage of sugars in all the tiers of the vertical tower than everbearing varieties V3 (Murano) and V4 (Vivara). Everbearing strawberry cultivars often show lower sugar content compared to short-day

Table 1. Effect of different dates of planting on total sugars of strawberry varieties in different tiers (Pooled)

Treatment	Total sugars (%)														
	Top tier					Middle tier					Lower tier				
	D1	D2	D3	D4	Mean	D1	D2	D3	D4	Mean	D1	D2	D3	D4	Mean
V1	5.11	5.65	5.72	5.17	5.41	5.15	5.77	5.30	4.43	5.16	4.67	5.45	5.13	4.36	4.90
V2	5.47	6.02	5.40	5.16	5.51	5.20	5.73	5.33	5.15	5.35	4.83	5.62	5.25	4.70	5.10
V3	4.54	5.39	5.28	4.58	4.95	4.35	5.27	5.25	4.24	4.78	4.30	5.27	5.02	4.46	4.76
V4	4.69	5.11	4.79	4.57	4.79	4.28	5.22	4.65	4.31	4.61	3.82	4.82	4.37	4.14	4.29
Mean	4.95	5.54	5.30	4.87		4.74	5.49	5.14	4.53		4.40	5.29	4.94	4.41	
	SEm±				CD (5%)	SEm±				CD (5%)	SEm±				CD (5%)
Varieties (V)	0.03				0.09	0.04				0.11	0.02				0.06
Dates of planting (D)	0.03				0.09	0.04				0.11	0.02				0.06
VxD	0.06				0.18	0.08				0.23	0.04				0.12

Table 2. Effect of different dates of planting on titratable acidity of strawberry varieties in different tiers (Pooled)

Treatment	Titratable acidity (%)														
	Top tier					Middle tier					Lower tier				
	D1	D2	D3	D4	Mean	D1	D2	D3	D4	Mean	D1	D2	D3	D4	Mean
V1	0.43	0.35	0.37	0.52	0.42	0.53	0.44	0.47	0.55	0.50	0.58	0.50	0.49	0.57	0.53
V2	0.36	0.27	0.38	0.47	0.37	0.50	0.40	0.44	0.55	0.47	0.54	0.45	0.48	0.51	0.49
V3	0.54	0.41	0.46	0.59	0.50	0.64	0.54	0.63	0.66	0.62	0.64	0.56	0.64	0.66	0.62
V4	0.56	0.50	0.58	0.65	0.57	0.53	0.54	0.54	0.67	0.57	0.60	0.56	0.56	0.72	0.61
Mean	0.47	0.38	0.45	0.56		0.55	0.48	0.52	0.61		0.59	0.52	0.54	0.61	
	SEm±				CD (5%)	SEm±				CD (5%)	SEm±				CD (5%)
Varieties (V)	0.01				0.03	0.01				0.02	0.01				0.03
Dates of planting (D)	0.01				0.03	0.01				0.02	0.01				0.03
VxD	0.02				NS	0.01				0.04	0.02				0.06

V1: Nabila, V2: Flaminia, V3: Murano, V4: Vivara

D1: Planting on October 15th, D2: Planting on October 30th, D3: Planting on November 15th, D4: Planting on November 30th

cultivars due to the differences in their physiological responses to environmental factors such as photoperiod and temperature. Everbearing strawberries have a facultative long-day flowering response, which means they accelerate flowering and fruiting under longer photoperiods and high temperatures which lead to higher respiration and lowers the allocation of food material, whereas short day cultivars require shorter photoperiods and lower temperatures to initiate flowering, this difference in flowering response affects the plant's carbohydrate metabolism and allocation (Rivero et al., 2022).

It was noted that climatic conditions, especially warmer climate and longer day length impacts negatively on sugar metabolism within fruit tissues (Chandler et al., 2003). In this experiment as mentioned above, among planting dates D2 (Planting on October 30th) and D3 (Planting on November 15th) revealed higher percentage of total sugars than other planting dates. This might be attributed to the better accumulation of sugars in berries due to the favorable climatic conditions preferably lower temperatures prevailing at that time of fruit development. Rahman (2014), Jahangeera et al. (2010) Sadiq and Kaur (2017), Anwar et al. (2016) also reported the same in strawberry fruits planted at different times.

Titrateable Acidity (%): The pooled data regarding titrateable acidity of strawberry varieties planted at various dates in top, middle and lower tier of the vertical tower was presented in Table 2.

The top tier of the vertical tower exhibited a significant difference among varieties and various dates of planting. Among varieties lowest titrateable acidity (0.37) was revealed by V2 (Flaminia) followed by V1 (Nabila) (0.42). While variety V4 (Vivara) has recorded the highest percentage of titrateable acidity (0.57). Among, dates of planting, D2 (Planting on October 30th) has revealed the lowest percentage of titrateable acidity (0.31) followed by D3 (Planting on November 15th) (0.45), while D4 (Planting on November 30th) recorded the highest percentage of titrateable acidity (0.56). There was no significant difference observed among the interactions between varieties and dates of planting in the percentage of titrateable acidity in the top tier of the vertical tower.

In the middle tier, there was a significant difference observed among varieties and various

dates of planting. The lowest titrateable acidity (0.47) was revealed in V2 (Flaminia) followed by V1 (Nabila) (0.50), while the variety V3 (Murano) has shown the highest percentage of titrateable acidity (0.62). Among various dates of planting, D2 (Planting on October 30th) has recorded the lowest percentage of titrateable acidity (0.48) followed by D3 (Planting on November 15th) (0.52), while D4 (Planting on November 30th) revealed the highest percentage of titrateable acidity (0.61). There was also a significant difference observed among the interactions between varieties and dates of planting in titrateable acidity in the middle tier of the vertical tower, the combination of V2D2 has revealed lowest percentage of titrateable acidity (0.40) which was on par with V1D2 (0.44). The interaction of V4D4 recorded a highest percentage of titrateable acidity (0.67) which was on par with V3D3 (0.63).

In the lowest tier, there was a significant difference among varieties and various dates of planting. The lowest titrateable acidity (0.49) was recorded by V2 (Flaminia) followed by V1 (Nabila) (0.53), while the variety V3 (Murano) has resulted with the highest percentage of titrateable acidity (0.62). Among various dates of planting, D2 (Planting on October 30th) has revealed the lowest percentage of titrateable acidity (0.52) which was on par with D3 (Planting on November 15th) (0.54), while, highest percentage of titrateable acidity (0.61) was noticed in D4 (Planting on November 30th) which was on par with D1 (Planting on October 15th) (0.59). There was also a significant difference observed among the interactions between varieties and dates of planting in titrateable acidity in the lower tier of the vertical tower, the combination of V2D2 has revealed the lowest percentage of titrateable acidity (0.45) which was on par with V1D2 (0.50), V1D3 (0.49), V2D3 (0.48) and V2D4 (0.51). The interaction of V4D4 recorded the highest percentage of titrateable acidity (0.72) which was on par with V3D4 (0.66).

The differences in titrateable acidity among varieties may be due to variations in adaptability to changing environments. In this experiment, short day varieties V1 (Nabila) and V2 (Vivara) have recorded with the lowest titrateable acidity than everbearing varieties V3 (Murano) and V4 (Vivara). The continuous production cycle of everbearing strawberries may contribute to more consistent accumulation of organic acids throughout the growing season (Miri et al., 2020). The variations in titrateable acidity among

varieties were also noticed in the experiments conducted by Jami et al. (2015) and Kumar et al. (2022).

In this experiment, among planting dates, planting in the late October and early November revealed the lowest titratable acidity in strawberry fruits than early October and late November planting which was similar to findings of Gogoi et al. (2022), Kaur (2010), Mawkhiew and Pereira (2015), Singh et al. (2006) and Sadiq and Kaur (2017) in strawberry. This was likely due to the differences in the environmental conditions such as temperature and the sunlight during plant growth and fruit development period which can alter the metabolic processes involved in the acid accumulation. In fruits, cooler nights and warmer days generally synthesize more acidity (Wani et al., 2017) which was supported by the findings of Bhamini et al. (2017) who noticed that fruit acidity is increased with higher temperatures in strawberry.

4. CONCLUSION

Based on this research, It can be concluded as follows:

- The top tier of the vertical tower has exhibited higher percentage of Total sugars and lower percentage of titratable acidity than subsequent tiers of the vertical tower among all the varieties, potting media and interactions.
- Among varieties, Shortday varieties V1 (Nabila) and V2 (Flaminia) has revealed highest percentage of total sugars and lowest percentage of titratable acidity than everbearing varieties V3 (Murano) and V4 (Vivara) in all the tiers of the vertical tower.
- Among planting dates, D2 (Planting on October 30th) has registered with highest percentage of Total sugars and lowest percentage of titratable acidity in all the tiers of the vertical tower.
- The interaction of V1 (Nabila) and V2 (Flaminia) with D2 (planting on October 30th) indicated highest percentage of total sugars and lowest percentage of titratable acidity. in all the tiers of the vertical tower.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Anwar, R., Ahma, M., Hussain, Z., Azam, M. and Moaaz, M. A. 2016. Transplanting time influences plant growth and fruit quality of Strawberries grown under subtropical climate. *Proceedings of Pakistan Society for Horticultural Science 2nd International Conference on Horticultural Sciences*, 18-20.
- Bhamini, K., Ruby, R., Abu, Md., Nayyer, Feza, A. and Afzal, A. 2017. Influence of Planting Dates and Temperature on Plant Growth, Flowering and Fruiting of Strawberry in Agro Climatic Condition of Bihar, India. *International Journal of Current Microbiology and Applied Sciences*, 6(8):3184-3191.
- FAO. 2020. Food and agriculture organization. *Statistical database*.
- Gogoi, B., Dutta, S. and Das, R.P. 2022. Response of Strawberry Cultivars to Different Planting Dates and Growing Conditions in Jorhat, Assam. *Indian journal of agricultural research*, 5860.
- Jahangeera, B., Kher, R., Bakshi, P. and Wah, V. K. 2010. Effect of planting time and mulching material on quality of strawberry. *Journal of Research SKUAST-J* 9: 54-62.
- Jami, Y.Y, Sarkar, A. and Maiti, C.S.2015.Evaluation of strawberry cultivars in the foothills of Nagaland. *Journal Crop and Weed*, 11:198-200.
- Kaur, A. 2010. Cultivation of strawberry under protected conditions in sub tropical regions of Punjab. Ph.D. *Thesis, GNDU*.
- Kumar,P., Rajesh, K.B., Hansra., Neeru, D. and Amit, K. 2022. Potting substrate effect on yield and quality of strawberry (*Fragaria x ananassa*) in terrace gardening. *Indian Journal of Agricultural Sciences*, 92(5):667-669.
- Mawkhiew, A. and Pereira, L. S.2015. Effect of planting time on yield and quality of Strawberry in West Garo Hills, Meghalaya. *Journal of Agriculture Technology*.2(1&2):44-48.
- Miri,S.M., Fatemeh, H. and Darab, H. 2020. Morpho-physiological performance of seven short-day and day-neutral strawberry cultivars in the soilless culture.

- Journal of Photochemistry and Photobiology B: Biology*, 10(1):127-139.
- Rahman, M.M. 2014. Interactive influence of planting date and cultivar on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch.). *Journal of Horticulture and Forestry*, 6(3):31-37.
- Rivero, R.I., Remberg, S.F., Ola, M., Heide., and Anita, S. 2022. Effect of temperature and photoperiod preconditioning on flowering and yield performance of three everbearing strawberry cultivars. *Acta Horticulturae*, 8(6):504-504.
- Sadiq, A. and Kaur, A. 2017. Effect of planting time on yield and quality of strawberry cv. chandler in subtropical region of punjab. *Asian Journal of Science and Technology*, 08(10):6080-6083.
- Singh, R., Sharma, R., and Rajiv, G. 2006. Interactive effects of planting time and mulching on 'Chandler' strawberry. *Scientia Horticulturae*, 111(4): 344-351
- Wani, R.A., Baba, J.A., Hakeem, S.A., Umar, Ashaq, I., Pandit, H., Mir, M.A., Zubair, M., Bashir, S., Nissa, S.U., Dar, N.A. and Parray, G.A. 2017. Influence of mulching material on albinism disorder in strawberry under cold arid conditions. *International Journal of Current Microbiology and Applied Sciences*, 6(7): 4287-4290.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/126464>