



# Weed Dynamics and Soil Health as Influenced by Tillage Method and Establishment Method in Small Millets

Hasanath T. <sup>a</sup>, Bindhu J. S. <sup>b\*</sup>, Shalini Pillai P. <sup>a</sup>, John J. <sup>b</sup>,  
Meera A. V. <sup>b</sup> and Archana B. <sup>a</sup>

<sup>a</sup> Department of Agronomy, College of Agriculture, Vellayani, India.

<sup>b</sup> AICRP-IFS, IFSRS, Karamana, Thiruvananthapuram, Kerala Agricultural University, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i113048>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126565>

Original Research Article

Received: 08/09/2024

Accepted: 10/11/2024

Published: 14/11/2024

## ABSTRACT

Summer rice fallows are the potential areas for crop diversification with millets. A field experiment was conducted at Integrated Farming System Research Station (IFSRS), Karamana during summer season of 2023-24 with an objective to find out the effect of tillage method and establishment method on weed dynamics and soil health. The field experiment was laid out in split-split plot design with 12 treatment combinations and three replications. The main plot treatments were tillage method (M<sub>1</sub>-minimum tillage and M<sub>2</sub> - conventional tillage), sub plot treatments were crops (C<sub>1</sub>-little millet, C<sub>2</sub>-foxtail millet and C<sub>3</sub>-proso millet) and the sub-sub plot treatments were establishment methods (P<sub>1</sub>- solid row planting and P<sub>2</sub>- broadcasting). The study indicated that grasses were the predominant weed flora in summer fallows and the lowest weed density and

\*Corresponding author: E-mail: [bindu.js@kau.in](mailto:bindu.js@kau.in);

**Cite as:** T., Hasanath, Bindhu J. S., Shalini Pillai P., John J., Meera A. V., and Archana B. 2024. "Weed Dynamics and Soil Health As Influenced by Tillage Method and Establishment Method in Small Millets". *Journal of Experimental Agriculture International* 46 (11):244-54. <https://doi.org/10.9734/jeai/2024/v46i113048>.

weed dry weight were observed in conventional tillage method with solid row planting. Among the establishment method, the highest N, P and K removal by weeds was recorded in broadcasting method (134.52 kg ha<sup>-1</sup>, 4.38 kg ha<sup>-1</sup> and 10.29 kg ha<sup>-1</sup>, respectively). The results revealed that the highest organic carbon (2.09%) was recorded in minimum tillage. The highest available N and P were recorded in minimum tillage and solid row planting. From the study it could be concluded that minimum tillage with solid row planting enhanced the soil health while conventional tillage with solid row planting is a viable option for weed management of small millets in summer rice fallows.

*Keywords: Diversification; minimum tillage; small millets; soil health; weed flora.*

## 1. INTRODUCTION

Small millets are the new staple crops in the new era of climate change. Millets are sustainable food source for combating hunger in changing world climate (Kumar et al. 2018). foxtail millet, little millet and proso millet were the prominent small millets that are said to be the most calming, least allergenic, and easily digested grains available.

Millets are not very good competitors with weeds, especially in the early growth stages. Depending on climate, edaphic, and biotic variables, millets can experience yield losses from weeds ranging from 5 to 94 percent (Sneha and Sheeja 2022). The critical period for millets typically occurs between 15 and 42 days after sowing (Krauss et al. 2022).

The number of weeds was greatly impacted by various tillage techniques. When compared to minimum and reduced tillage, conventional tillage dramatically decreased the population of weeds, regardless of the type of weed. Weed populations were significantly reduced as a result of the deeper positioning of weed seeds caused by conventional tillage, which prevented weeds from emerging from the soil (Vijaymahantesh et al. 2013). Tshering et al. 2022 reported that direct-seeded millet recorded a lower number of productive tillers (2.65 tillers/hill) which could be due to higher weed pressure during the initial growth stages. In transplanted millet, there was no overcrowding of seedlings, and weed pressure was considerably less.

Uncontrolled weed infestation significantly reduces the crop yield between and 15 to 83% in sorghum, 16 to 94% in pearl millet, and 55 to 61% in finger millet depending on crop cultivars, nature, and intensity of weed infestation, management practices, and environmental condition (Mishra et al. 2014).

Conservation agriculture proved to be an excellent alternative to conventional agriculture in the long term of sustainable crop production and soil organic carbon sequestration (Choudhury et al. 2014). Conservation agriculture practices involving zero tillage, crop residue management and suitable crop rotation can serve as a better alternative to conventional agriculture for maintaining soil quality.

Understanding the role of tillage in weed management for millet cultivation is critical for balancing effective weed control with sustainable soil management practices. As farmers face challenges from labour constraints, changing climates, and a need for more sustainable farming systems, the role of tillage in managing weeds for millet crops is evolving and requires careful consideration of its benefits and limitations.

## 2. MATERIALS AND METHODS

The field study was conducted at Integrated Farming System Research Station (IFSRS), Karamana, Thiruvananthapuram, Kerala, India during summer season of 2023-24. The experimental site was positioned at an elevation of 5 meters above mean sea level on soil pH 5.7, bulk density 1.46 Mg m<sup>-3</sup> and organic carbon 1.6%. The available nitrogen, phosphorus and potassium recorded was 188.16 kg ha<sup>-1</sup>, 23.5 kg ha<sup>-1</sup> and 158.83 kg ha<sup>-1</sup>, respectively. A total rainfall of 674.88 mm was recorded during the study. The maximum and minimum temperature recorded during the study was 33.49<sup>o</sup> and 25.07<sup>o</sup>C, respectively.

The experiment was laid out in split-split plot design with 12 treatment combinations and three replications. The main plot treatment was tillage (M<sub>1</sub>- minimum tillage and M<sub>2</sub>- Conventional tillage), the sub plot treatment was crop (C<sub>1</sub> – little millet, C<sub>2</sub> – foxtail millet and C<sub>3</sub>- proso millet) and the sub-sub plot treatment was

establishment method (P<sub>1</sub>: solid row planting and P<sub>2</sub>: Broadcasting).

In minimum tillage, the field will be ploughed once and seed bed will be taken retaining the crop residues of the previous crop of rice. In conventional tillage, the field will be ploughed thrice followed by secondary tillage without the crop residues. Solid row planting was done with inter row spacing of 25cm and plants were thinned at 10 DAS to maintain an intra-row spacing of 10 cm. ATL 1, DHFt-109-3 and TNAU 202 were the varieties used for the study for little millet, foxtail millet and proso millet, respectively. Farm yard manure and fertilizer application was done as per IIMR recommendations. ATL 1, DHFt-109-3 and TNAU 202 were the varieties used for the study for little millet, foxtail millet and proso millet, respectively. The crop was raised as a rainfed crop and one irrigation was given at 3 DAS and weeding operations were carried out manually at 30 DAS and 45 DAS.

### 3. RESULTS AND DISCUSSION

#### 3.1 weed Flora

The floristic composition of experimental site is given in Fig. 1 Sixteen weed species were observed in experimental field. The predominant weed flora were grasses. The predominant species among grasses were *Echinochloa colona*, *Digitaria ciliaris*, *Isachne miliacea* and *Cynodon dactylon*. Among, broad- leaf weeds *Oldenlandia umbellate*, *Alternanthera sessilis* and *Euphorbia hirta*. *Cyperus iria* was predominant under sedges (Table 1).

#### 3.2 weed Density and Weed Dry Weight

Weed density was significantly influenced by tillage method at 15 DAS, 30DAS and 45 DAS (Tables 2a and 2b, Fig. 2). Weed density was the lowest (154.44, 56.33 and 12.44 no. m<sup>-2</sup>, respectively) at conventional tillage method (M<sub>2</sub>) and the highest at minimum tillage method (238.33, 75.94 and 23.72 no. m<sup>-2</sup>, respectively). At 15 DAS, crop had significant influence on weed density. Lower weed density was observed in little millet (184.75 no. m<sup>-2</sup>) and it was on par with foxtail millet (196.67 no. m<sup>-2</sup>). Between the establishment method solid row planting recorded the lowest weed density (188.06, 60.67 and 14.72 no. m<sup>-2</sup>) followed by broadcasting method. The possible reasons of higher weed incidence in broadcast seeding relative to drill seeding may be because of uneven/non-uniform crop stand in broadcast seeding compared to line-seeding manually or by drill (Saha et al. 2021).

The results revealed that tillage method and establishment method had significant influence on weed dry weight (Tables 3a and 3b, Fig. 3). At 15 DAS, 30DAS and 45 DAS, the highest weed dry weight was recorded in minimum tillage (53.12 g m<sup>-2</sup>, 68.50 g m<sup>-2</sup> and 76.62 g m<sup>-2</sup>, respectively). Between the establishment method, the highest weed dry weight was recorded in broadcasting method (46.21 g m<sup>-2</sup>, 59.33 g m<sup>-2</sup> and 67.28 g m<sup>-2</sup>) at 15, 30 and 45 DAS. At 45 DAS, crop had significant effect on weed dry weight. The highest weed dry weight (64.25 g m<sup>-2</sup>) was observed in proso millet and it was on par with foxtail millet.





Fig. 1. Floristic composition of weed in the experimental field

**Table 1. Weed flora observed in the experimental site**

Scientific name	Common name	Family
<b>Grasses</b>		
<i>Cynodon dactylon</i>	Bermuda grass	Poaceae
<i>Digitaria ciliaris</i>	Finger grass/ Henry's crabgrass	Poaceae
<i>Echinochloa colona</i>	Jungle rice	Poaceae
<i>Isachne miliacea</i>	Blood grass	Poaceae
<i>Setaria barbata</i>	Bristly foxtail grass/ corn grass	
<b>Broad leaved weeds</b>		
<i>Alternanthera sessilis</i>	Sessile joyweed, carpet weed	Amaranthaceae
<i>Boerhaavia diffusa</i>	Punarnava / Spreading hogweed	Nyctaginaceae
<i>Cleome rutidosperma</i>	Purple cleome	Capparaceae
<i>Commelina benghalensis</i>	Benghal day flower	Commelinaceae
<i>Euphorbia hirta</i>	Asthma plant	Euphorbiaceae
<i>Limncharis flava</i>	Yellow velvet leaf	Alismataceae
<i>Ludwigia perennis</i>	Perennial water primerose	Onagraceae
<i>Oldenlandia umbellata</i>	Chay root	
<i>Phyllanthus niruri</i>	Seed- under- leaf/ Gale of the swind	Euphorbiaceae
<b>Sedges</b>		
<i>Cyperus iria</i>	Rice flat sedge/ Umbrella sedge	Cyperaceae
<i>Fimbristylis miliacea</i>	Fimbry/ Lesser fimbriatylis	Cyperaceae

**Table 2a. Effects of treatments on weed density (no. m<sup>-2</sup>)**


Treatments	15DAS	30 DAS	45 DAS
<b>Tillage method (M)</b>			
M <sub>1</sub> - Minimum tillage	238.33	75.94	23.72
M <sub>2</sub> - Conventional tillage	154.44	56.33	12.44
SEm (±)	2.29	0.62	0.82
P= 0.05	13.910	3.757	5.008
<b>Crop (C)</b>			
C <sub>1</sub> -Little millet	184.75	63.33	16.08
C <sub>2</sub> - Foxtail millet	196.67	70.00	19.33
C <sub>3</sub> - Proso millet	207.75	65.08	18.83
SEm (±)	4.50	2.31	1.34
P= 0.05	14.675	NS	NS
<b>Establishment method (P)</b>			
P <sub>1</sub> - Solid row planting	188.06	60.67	14.72
P <sub>2</sub> - Broadcasting	204.72	71.61	21.44
SEm (±)	2.20	2.24	0.76
P= 0.05	6.790	6.899	2.347



**Fig. 2. Effect of treatment on weed density (no.m<sup>-2</sup>)**

**Table 2b. Interaction effects of treatments on weed density (no. m<sup>-2</sup>)**

Treatments	15 DAS	30 DAS	45 DAS
<b>M×C</b>			
M <sub>1</sub> C <sub>1</sub>	248.17	69.67	23.67
M <sub>1</sub> C <sub>2</sub>	218.33	86.00	24.33
M <sub>1</sub> C <sub>3</sub>	248.50	72.17	23.17
M <sub>2</sub> C <sub>1</sub>	121.33	57.00	8.50
M <sub>2</sub> C <sub>2</sub>	175.00	54.00	14.33
M <sub>2</sub> C <sub>3</sub>	167.00	58.00	14.50
SEm±	6.36	3.27	1.90
P= 0.05	20.753	10.660	NS
<b>M×P</b>			
M <sub>1</sub> P <sub>1</sub>	230.33	70.33	18.78
M <sub>1</sub> P <sub>2</sub>	246.33	81.56	28.67
M <sub>2</sub> P <sub>1</sub>	145.78	51.00	10.67
M <sub>2</sub> P <sub>2</sub>	163.11	61.67	14.22
SEm±	3.12	3.17	1.08
P= 0.05	NS	NS	3.319
<b>C×P</b>			
C <sub>1</sub> P <sub>1</sub>	173.17	57.17	12.17
C <sub>1</sub> P <sub>2</sub>	196.33	69.50	20.00
C <sub>2</sub> P <sub>1</sub>	188.83	64.17	16.33
C <sub>2</sub> P <sub>2</sub>	204.50	75.83	22.33
C <sub>3</sub> P <sub>1</sub>	202.17	60.67	15.67
C <sub>3</sub> P <sub>2</sub>	213.33	69.50	22.00
SEm±	3.82	3.88	1.32
P= 0.05	NS	NS	NS
<b>M×C×P</b>			
M <sub>1</sub> C <sub>1</sub> P <sub>1</sub>	236.67	64.33	17.67
M <sub>1</sub> C <sub>1</sub> P <sub>2</sub>	259.67	75.00	29.67
M <sub>1</sub> C <sub>2</sub> P <sub>1</sub>	212.33	79.00	20.00
M <sub>1</sub> C <sub>2</sub> P <sub>2</sub>	224.33	93.00	28.67
M <sub>1</sub> C <sub>3</sub> P <sub>1</sub>	242.00	67.67	18.67
M <sub>1</sub> C <sub>3</sub> P <sub>2</sub>	255.00	76.67	27.67
M <sub>2</sub> C <sub>1</sub> P <sub>1</sub>	109.67	50.00	6.67
M <sub>2</sub> C <sub>1</sub> P <sub>2</sub>	133.00	64.00	10.33
M <sub>2</sub> C <sub>2</sub> P <sub>1</sub>	165.33	49.33	12.67
M <sub>2</sub> C <sub>2</sub> P <sub>2</sub>	184.67	58.67	16.00
M <sub>2</sub> C <sub>3</sub> P <sub>1</sub>	162.33	53.67	12.67
M <sub>2</sub> C <sub>3</sub> P <sub>2</sub>	171.67	62.33	16.33
SEm±	5.40	5.49	1.87
P= 0.05	NS	NS	NS



**Fig. 3. Effect of treatment on weed dry weight (g m<sup>-2</sup>)**

**Table 3a. Effect of treatment on weed dry weight (g m<sup>-2</sup>)**

Treatments	15DAS	30 DAS	45 DAS
<b>Tillage method (M)</b>			
M <sub>1</sub> - Minimum tillage	53.12	68.50	76.62
M <sub>2</sub> - Conventional tillage	33.61	39.97	45.40
SEm (±)	0.08	1.79	2.513
P= 0.05	0.450	10.872	15.291
<b>Crop (C)</b>			
C <sub>1</sub> -Little millet	45.43	52.96	57.25
C <sub>2</sub> - Foxtail millet	43.68	55.58	61.54
C <sub>3</sub> - Proso millet	41.00	54.17	64.25
SEm (±)	2.33	1.31	1.59
P= 0.05	NS	NS	5.190
<b>Establishment method (P)</b>			
P <sub>1</sub> - Solid row planting	40.52	49.14	54.74
P <sub>2</sub> - Broadcasting	46.21	59.33	67.28
SEm (±)	0.80	1.43	1.28
P= 0.05	2.467	4.416	3.933

**Table 3b. Effect of treatment on weed dry weight (g m<sup>-2</sup>)**

Treatments	15 DAS	30 DAS	45 DAS
<b>M×C</b>			
M <sub>1</sub> C <sub>1</sub>	57.35	68.67	68.32
M <sub>1</sub> C <sub>2</sub>	56.02	69.17	81.03
M <sub>1</sub> C <sub>3</sub>	46.00	67.67	80.51
M <sub>2</sub> C <sub>1</sub>	33.50	37.25	46.18
M <sub>2</sub> C <sub>2</sub>	31.33	42.00	42.05
M <sub>2</sub> C <sub>3</sub>	36.00	40.67	47.98
SEm±	3.29	1.85	2.25
P= 0.05	NS	NS	7.34
<b>M×P</b>			
M <sub>1</sub> P <sub>1</sub>	51.38	60.67	69.77
M <sub>1</sub> P <sub>2</sub>	54.87	76.33	83.47
M <sub>2</sub> P <sub>1</sub>	29.67	37.61	39.71
M <sub>2</sub> P <sub>2</sub>	37.56	42.33	51.10
SEm±	1.13	2.03	1.81
P= 0.05	NS	6.245	NS
<b>C×P</b>			
C <sub>1</sub> P <sub>1</sub>	41.28	45.75	48.43
C <sub>1</sub> P <sub>2</sub>	49.57	60.17	66.07
C <sub>2</sub> P <sub>1</sub>	39.45	51.67	56.82
C <sub>2</sub> P <sub>2</sub>	47.90	59.50	66.26
C <sub>3</sub> P <sub>1</sub>	40.83	50.00	58.97
C <sub>3</sub> P <sub>2</sub>	41.17	58.33	69.53
SEm±	1.39	2.48	2.21
P= 0.05	4.274	NS	NS
<b>M×C×P</b>			
M <sub>1</sub> C <sub>1</sub> P <sub>1</sub>	50.90	60.67	57.00
M <sub>1</sub> C <sub>1</sub> P <sub>2</sub>	63.80	76.67	79.63
M <sub>1</sub> C <sub>2</sub> P <sub>1</sub>	51.57	61.67	77.01
M <sub>1</sub> C <sub>2</sub> P <sub>2</sub>	60.47	76.67	85.05
M <sub>1</sub> C <sub>3</sub> P <sub>1</sub>	51.67	59.67	75.30
M <sub>1</sub> C <sub>3</sub> P <sub>2</sub>	40.33	75.67	85.72
M <sub>2</sub> C <sub>1</sub> P <sub>1</sub>	31.67	30.83	39.85
M <sub>2</sub> C <sub>1</sub> P <sub>2</sub>	35.33	43.67	52.50
M <sub>2</sub> C <sub>2</sub> P <sub>1</sub>	27.33	41.67	36.63
M <sub>2</sub> C <sub>2</sub> P <sub>2</sub>	35.33	42.33	47.47
M <sub>2</sub> C <sub>3</sub> P <sub>1</sub>	30.00	40.33	42.63
M <sub>2</sub> C <sub>3</sub> P <sub>2</sub>	42.00	41.00	53.33
SEm±	1.96	3.51	3.13
P= 0.05	6.044	NS	NS

**Table 4a. Effect of treatments on nutrient removal by weeds (kg ha<sup>-1</sup>)**

Treatments	Nitrogen removal (kg ha <sup>-1</sup> )	Phosphorus removal (kg ha <sup>-1</sup> )	Potassium removal (kg ha <sup>-1</sup> )
<b>Tillage method (M)</b>			
M <sub>1</sub> - Minimum tillage	16.54	5.25	12.34
M <sub>2</sub> - Conventional tillage	7.63	2.70	6.20
SEm (±)	0.60	0.12	0.31
P= 0.05	3.69	0.726	1.910
<b>Crop (C)</b>			
C <sub>1</sub> -Little millet	11.75	3.63	8.65
C <sub>2</sub> - Foxtail millet	11.95	3.98	9.29
C <sub>3</sub> - Proso millet	12.55	4.31	9.88
SEm (±)	0.56	0.10	0.28
P= 0.05	NS	0.318	0.926
<b>Establishment method (P)</b>			
P <sub>1</sub> - Solid row planting	10.71	3.57	8.25
P <sub>2</sub> - Broadcasting	13.45	4.38	10.29
SEm (±)	0.38	0.14	0.21
P= 0.05	1.17	0.433	0.643

Field ploughing move weed seeds bidirectionally, and change the soil properties and thereby altering weed seed germination (Singh et al. 2015). Reduction in tillage operations as in minimum tillage increases the weed infestation. Tillage changes vertical distribution of weed seeds in soil profile and soil physical properties, and affects emergence and seed survival of weeds through changes in soil conditions and determines weed seedling emergence and species composition (Mishra et al. 2019). Abid et al. 2007 reported that weed density and dry weight were lower under conventional tillage followed by hand weeding and minimum tillage followed by herbicide treatments in summer fallows cultivated with green gram.

### 3.3 N, P and K Removal by Weeds

The results revealed that tillage method and establishment method had significant influence on N, P and K removal by weeds (Tables 4a and 4b). Among the tillage method, the highest nitrogen and phosphorus and potassium removal was recorded in minimum tillage (16.54 kg ha<sup>-1</sup>, 5.25 kg ha<sup>-1</sup> and 12.34 kg ha<sup>-1</sup>, respectively). Between the establishment method, the highest N, P and K removal was recorded in broadcasting method (13.45 kg ha<sup>-1</sup>, 4.38 kg ha<sup>-1</sup> and 10.29 kg ha<sup>-1</sup>, respectively). This may be due to the highest dry weight of weeds in the treatments. Crop had significant influence on P and K removal by weed. The highest P removal (4.31 kg ha<sup>-1</sup>) was recorded in proso millet and higher K removal (9.88 kg ha<sup>-1</sup>) was also recorded in proso millet and it was on

par with foxtail millet. The predominant grassy weeds caused more K removal by the weeds.

At 30 DAS and 45 DAS, nitrogen content of weed was significantly influenced by tillage method. The highest nitrogen content was observed in minimum tillage method (1.86 and 2.16 %, respectively) followed by conventional tillage method (1.37 and 1.69 %, respectively). The crop residues in the minimum tillage resulted more nutrient in soil resulted in more nutrient uptake. At 30 DAS, phosphorus content of weed was significantly influenced by tillage method. The highest P content (0.35 %) was observed in minimum tillage method followed by conventional tillage method (0.31 %). Tillage method had significant influence on potassium content of weed at 30 DAS and 45 DAS. The highest value was observed in minimum tillage method (1.48 and 1.61%, respectively) followed by conventional tillage method (1.26 and 1.36%, respectively). This is due to the predominance of grass weeds.

### 3.4 Plant Uptake of Nutrients

The tillage method had significant influence on K content. The highest K content was observed in conventional tillage method (3.06%) followed by minimum tillage method (2.96%). Among tillage method, the highest nutrient uptake (N, P and K) was observed in conventional tillage method (153.98, 28.57 and 110.83 kg ha<sup>-1</sup>, respectively). It can be attributed due to better soil parameters due to conventional tillage. The lowest nutrient uptake was in minimum tillage method (130.75, 23.86 and 96.09 kg ha<sup>-1</sup>, respectively).

**Table 4b. Interaction effect of treatments on nutrient removal by weeds (kg ha<sup>-1</sup>)**

Treatments	Nitrogen removal (kg ha <sup>-1</sup> )	Phosphorus removal (kg ha <sup>-1</sup> )	Potassium removal (kg ha <sup>-1</sup> )
<b>M×C</b>			
M <sub>1</sub> C <sub>1</sub>	15.50	4.59	11.06
M <sub>1</sub> C <sub>2</sub>	17.01	5.49	12.79
M <sub>1</sub> C <sub>3</sub>	17.10	5.65	13.15
M <sub>2</sub> C <sub>1</sub>	8.00	2.67	6.24
M <sub>2</sub> C <sub>2</sub>	6.89	2.46	5.78
M <sub>2</sub> C <sub>3</sub>	7.99	2.97	6.59
SEm±	0.79	0.14	0.40
P= 0.05	NS	0.449	NS
<b>M×P</b>			
M <sub>1</sub> P <sub>1</sub>	14.76	4.90	11.05
M <sub>1</sub> P <sub>2</sub>	18.32	5.59	13.63
M <sub>2</sub> P <sub>1</sub>	6.67	2.24	5.45
M <sub>2</sub> P <sub>2</sub>	8.58	3.16	6.95
SEm±	0.54	0.20	0.30
P= 0.05	NS	NS	NS
<b>C×P</b>			
C <sub>1</sub> P <sub>1</sub>	9.98	3.26	7.44
C <sub>1</sub> P <sub>2</sub>	13.52	4.00	9.86
C <sub>2</sub> P <sub>1</sub>	10.79	3.64	8.48
C <sub>2</sub> P <sub>2</sub>	13.11	4.31	10.09
C <sub>3</sub> P <sub>1</sub>	11.37	3.81	8.83
C <sub>3</sub> P <sub>2</sub>	13.73	4.81	10.91
SEm±	0.66	0.24	0.36
P= 0.05	NS	NS	NS
<b>M×C×P</b>			
M <sub>1</sub> C <sub>1</sub> P <sub>1</sub>	12.90	4.32	9.37
M <sub>1</sub> C <sub>1</sub> P <sub>2</sub>	18.10	4.87	12.76
M <sub>1</sub> C <sub>2</sub> P <sub>1</sub>	15.65	5.27	11.97
M <sub>1</sub> C <sub>2</sub> P <sub>2</sub>	18.37	5.71	13.62
M <sub>1</sub> C <sub>3</sub> P <sub>1</sub>	15.72	5.11	11.81
M <sub>1</sub> C <sub>3</sub> P <sub>2</sub>	18.48	6.19	14.50
M <sub>2</sub> C <sub>1</sub> P <sub>1</sub>	7.07	2.21	5.512
M <sub>2</sub> C <sub>1</sub> P <sub>2</sub>	8.93	3.13	6.96
M <sub>2</sub> C <sub>2</sub> P <sub>1</sub>	5.93	2.00	4.99
M <sub>2</sub> C <sub>2</sub> P <sub>2</sub>	7.86	2.91	6.57
M <sub>2</sub> C <sub>3</sub> P <sub>1</sub>	7.01	2.50	5.85
M <sub>2</sub> C <sub>3</sub> P <sub>2</sub>	8.97	3.43	7.32
SEm±	0.93	0.34	0.51
P= 0.05	NS	NS	NS

Among the crop, highest nutrient uptake (N, P and K) was observed in little millet (175.56, 30.68 and 118.82 kg ha<sup>-1</sup>, respectively). The lower nitrogen uptake was observed in proso millet (124.58 kg ha<sup>-1</sup>) and it was at par with foxtail millet (126.95 kg ha<sup>-1</sup>). The lower phosphorus uptake was observed in foxtail millet (23.94 kg ha<sup>-1</sup>) and it was on par with proso millet (24.011 kg ha<sup>-1</sup>). The lowest K uptake was observed in foxtail millet (92.53 kg ha<sup>-1</sup>). Between the establishment methods the highest nutrient uptake (N, P and K) was observed in broadcasting method of sowing (146.55, 26.98, and 106.28 kg ha<sup>-1</sup>) followed by solid row planting (138.17, 25.44 and 106.28 kg ha<sup>-1</sup>).

Among the interaction effects, tillage method and crop interaction were found to be significant on N and P uptake. The highest value was recorded in M<sub>2</sub>C<sub>1</sub> interaction (195.26 and 33.86 kg ha<sup>-1</sup>). P uptake was found to be significant on M×P interaction. The highest value was recorded in M<sub>2</sub>P<sub>2</sub> interaction (29.31249 kg ha<sup>-1</sup>). C×P interaction had significant influence on N and K uptake. The highest value was recorded in C<sub>1</sub>P<sub>2</sub> interaction (186.33 and 127.93 kg ha<sup>-1</sup>). M×C×P interaction was found to be significant on P uptake. The higher value was recorded in M<sub>2</sub>C<sub>1</sub>P<sub>2</sub> (34.63 kg ha<sup>-1</sup>) and it was on par with M<sub>2</sub>C<sub>1</sub>P<sub>1</sub> interaction (33.08 kg ha<sup>-1</sup>). Alam et al. (2023) reported that significantly the highest



value of total N uptake, total P uptake and total K uptake were observed in conventional tillage.

### 3.5 Soil Nutrient Status

The soil bulk density decreased with degree of soil ploughing during tillage practices. From the experimental results it was observed that conventional tillage recorded the lowest bulk density of (1.42 Mg m<sup>-3</sup>). Orzech et al. 2021 reported that the changes in the bulk density and moisture content of soil varied across crop species, the developmental stages of plants, and soil horizons.

Soil organic carbon is an indicator of soil health and microbial activity. The minimum tillage method significantly affected the soil organic carbon content (Fig. 4). The highest organic carbon content was observed in minimum tillage. This indicated that, compared to conventional tillage there was significant gain in soil organic carbon, that is 37% more organic carbon than conventional tillage method. Krauss et al. (2022) reported that the increase in soil organic carbon stocks between reduced tillage (RT) and Conventional tillage (CT) ranged from +3% to +44% depending on site and were significantly higher in RT in different locations of the experiment.

The results revealed that the highest organic carbon (2.09 %), available N and P were

recorded in minimum tillage and solid row planting. Foxtail millet recorded the highest available N and P, while the highest available K was recorded in proso millet. Seth et al (2020) obtained that organic carbon and available N, P and K was higher in zero tillage than conventional tillage. Tillage options did not influence the available potassium in soil after harvest of wheat and rice.

Between the tillage method, the highest available nitrogen content was recorded minimum tillage (301.85 kg ha<sup>-1</sup>). Among the crop, the highest available nitrogen was observed in foxtail millet (295.54 kg ha<sup>-1</sup>). The lowest nitrogen was recorded in little millet (277.50 kg ha<sup>-1</sup>). Among the establishment method, the highest available nitrogen (288.74 kg ha<sup>-1</sup>) was observed in solid row planting. Minimum tillage recorded the highest available phosphorus content (47.00 kg ha<sup>-1</sup>). Among the crop, higher available phosphorus content observed in foxtail millet (40.58 kg ha<sup>-1</sup>) and it was at par with proso millet. The lowest available phosphorus (36.17 kg ha<sup>-1</sup>) observed in little millet (Dubey et al. 2023).

The analysed data revealed that crop had significant influence on available potassium content. The higher available potassium was recorded in proso millet (163.17 kg ha<sup>-1</sup>) and it was on par with foxtail millet. The lowest available potassium was recorded in little millet (155.83 kg ha<sup>-1</sup>).




Fig. 4. Effect of tillage method on organic carbon content in the post- harvest soil, per cent

SEm (±): 0.060

P= 0.05=0.390

#### 4. CONCLUSION

Tillage and establishment method had a substantial impact on weed density, nutrient uptake, and soil properties and soil health. Conventional tillage, in combination with solid row planting, was most effective in minimizing weed density and dry weight, promoting optimal nutrient uptake by crops, and reducing nutrient losses to weeds. Minimum tillage and solid row planting contributed to higher organic carbon levels and nutrient availability in the soil. These findings suggest that adopting an appropriate tillage and establishment method can enhance crop yield and soil health, making it a viable strategy for sustainable agricultural practices.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Abid, V. (2017). *Performance of green gram (Vigna radiata (L.) Wilczek) cultivars under different tillage methods* (M.Sc. (Ag) Thesis). Kerala Agricultural University, Thrissur.
- Choudhury, S.G., Srivastava, S., Singh, R., Chaudhari, S.K., Sharma, D.K., Singh, S.K., & Sarkar, D. (2014). Tillage and residue management effects on soil aggregation, organic carbon dynamics, and yield attribute in rice–wheat cropping system under reclaimed sodic soil. *Soil and Tillage Research*, 136, 76-83. <https://doi.org/10.1016/j.still.2013.10.007>
- Dubey, R.P., Chethan, C.R., Choudhary, V.K., & Mishra, J.S. (2023). A review on weed management in millets. *Indian Journal of Weed Science*, 55(2), 141-148.
- Krauss, M., Wiesmeier, M., Don, A., Cuperus, F., Gattinger, A., Gruber, S., Haagsma, W.K., Peigné, J., Palazzoli, M.C., Schulz, F., & van der Heijden, M.G. (2022). Reduced tillage in organic farming affects soil organic carbon stocks in temperate Europe. *Soil and Tillage Research*, 216, 105262. <https://doi.org/10.1016/j.still.2021.105262>
- Kumar, A., Tomer, V., Kaur, A., Kumar, V., & Gupta, K. (2018). Millets: A solution to agrarian and nutritional challenges. *Agriculture & Food Security*, 7, 31-37. <https://doi.org/10.1186/s40066-018-0176-x>
- Mishra, J.S., Kumar, R., Kumar, R., Rao, K.K., & Bhatt, B.P. (2019). Weed density and species composition in rice-based cropping systems as affected by tillage and crop rotation. *Indian Journal of Weed Science*, 51(2), 116-122.
- Mishra, J.S., Rao, S.S., & Patil, J.V. (2014). Influence of sorghum cultivars and weed management practices on nutrient uptake by crop and weeds in semi-arid tropical India. *Indian Journal of Plant Physiology*, 19(4), 351-355.
- Orzech, K., Wanic, M., & Załuski, D. (2021). The effects of soil compaction and different tillage systems on the bulk density and moisture content of soil and the yields of winter oilseed rape and cereals. *Agriculture*, 11(7), 666. <https://doi.org/10.3390/agriculture1107066>
- Saha, S., Munda, S., Singh, S., Kumar, V., Jangde, H.K., Mahapatra, A., & Chauhan, B.S. (2021). Crop establishment and weed control options for sustaining dry direct seeded rice production in eastern India. *Agronomy*, 11(2), 389. <https://doi.org/10.3390/agronomy11020389>
- Seth, M., Manuja, S., & Singh, S. (2020). Effect of tillage and site-specific nutrient management on yield, nutrient uptake and status of soil in wheat in rice-wheat cropping system. *Journal of Crop and Weed*, 16(3), 32-37.
- Singh, V.P., Barman, K.K., Raghwendra Singh, P.K., & Sharma, A.R. (2015). *Weed management in conservation agriculture system*. ICAR - Directorate of Weed Research, Jabalpur, India.
- Sneha, S.R., & Sheeja, K.R. (2022). Weed management in millets – A holistic approach. *Agricultural Reviews*, 1-8.
- Tshering, L., Dendupr, C., Dorjir, S., Rair, K., Ngawangr, & Lakeys. (2022). Effect of planting methods on grain yield and crop management in finger millet. *Bhutanese Journal of Agriculture*, 5(1), 135-151.
- Vijaymahantesh, Nanjappa, H.V., & Ramachandrappa, B.K. (2013). Effect of tillage and nutrient management practices

on weed dynamics and yield of finger millet under rainfed pigeon pea-finger millet system in Alfisols of southern India. *African Journal of Agriculture Research*, 8(21), 2470-2475.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/126565>