



# **Chemical Properties, Microbial Biomass and Enzyme Dynamics of Post-Harvest Soil on Transplanted Rice with Effect of Organic Amendments**

**R. Ajaykumar<sup>1\*</sup> and N. Thavaprakash<sup>2</sup>**

<sup>1</sup>*Department of Agronomy, Vanavarayar Institute of Agriculture, Pollachi – 642 103, India.*

<sup>2</sup>*Department of Agronomy, Tamil Nadu Agricultural University, Coimbatore – 641 003, India.*

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/CSJI/2020/v29i930205

### Editor(s):

(1) Prof. Pradip K. Bhowmik, University of Nevada, USA.

### Reviewers:

(1) Selma Hamimed, Carthage University, Tunisia.

(2) Rifkah Akmalina, Universitas Pamulang, Indonesia.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/64901>

**Original Research Article**

**Received 25 October 2020**

**Accepted 30 December 2020**

**Published 31 December 2020**

## **ABSTRACT**

A field experiment was conducted at the farm of Tamil Nadu Agricultural University, Coimbatore during kharif season to study the effect of soil chemical properties, microbial biomass and soil enzyme dynamics on transplanted rice with organic amendments. Rice CO(R) 48 was used as a test variety. The experiment was laid out in randomized block design (RBD) with three replications and nine treatments. The treatment consists Recommended Dose of Fertilizers 100% NPK (150 : 50 : 50 kg ha<sup>-1</sup>) through inorganic fertilizers, Based on N equivalent basis, required quantities of organic manures (Dhaincha, vermicompost and Farmyard Manure @ 50% and 100%) were incorporated into the soil one week before transplanting of rice. The P and K requirement was supplied separately through inorganic sources as per treatment schedule. In the present investigation, an attempt was made to examine the influence of different levels and sources of fertilization on dynamics of soil chemical properties, microbial biomass and enzyme activities under anaerobic rice cultivation. Soil samples were taken before the start of experiment and harvest of

\*Corresponding author: E-mail: [ajaykumar.tnau@gmail.com](mailto:ajaykumar.tnau@gmail.com);

rice for analysis of soil chemical characteristics, microbial biomass and enzyme activity. The results revealed that maximum microbial population (bacteria, fungal & actinomycetes), soil enzyme dynamics (urease, dehydrogenase & soil phosphatase activity), soil available macro nutrients (N, P, K) & micro nutrients (Fe, Zn, Mn, Cu) were significantly influenced with application of 100% N through dhaincha + balance P & K through inorganic fertilizers followed by application of 100% NPK through inorganic fertilizers + 6.25 t dhaincha. Least enzyme activity, microbial population & soil available nutrients of rice were registered in absolute control.

*Keywords: Microbial biomass; enzyme dynamics; soil available nutrients; organic manure; rice.*

## 1. INTRODUCTION

The natural ability of soil to support optimum growth and yield of crops depends not only on its physical and chemical properties, but also on the intensity of biological processes. Soil microbes mediate the biochemical transformations of organic matter that underpin essential ecosystem functions, including decomposition, mineralization of plant available nutrients, and nutrient retention. Organic production relies on these microbially- derived ecosystem functions and thus, may be a model system for ecological intensification of agriculture [1]. Soil fertility at great extent is controlled by different biochemical activities of the microflora, especially in the immediate surroundings of the roots, the rhizosphere, which under the influence of roots, carry a particularly dense population of microorganisms. Microorganisms can alter the degree of nutrient supply to higher plants through decomposition of organic compounds, immobilization of available nutrients, mineralization of nutrients or by promoting solubilization of fixed or insoluble mineral forms. The quantity and quality of soil organic matter and carbon and nitrogen inputs are the overriding controls on soil microbial biomass and activity. Thus, distinct organic amendments (e.g. manure, leguminous cover crops, and composted materials) can stimulate microbial biomass differently through increased labile organic matter [2].

Soil enzymes are a group of enzymes present in soil and are continuously playing important role in maintaining soil ecology, physical and chemical properties, fertility and health. These enzymes also play a key role in biochemical functions in the overall process of organic matter decomposition in soil system. They are important in catalyzing several vital reactions necessary for the life processes of micro-organisms in soil and stabilization of soil structure, decomposition of organic wastes, organic matter formation and nutrient cycling and hence, play an important role

in agriculture. The enzymatic activities of soil catalyzes the biochemical activities performed by bacteria and thereby, indicates the potential of soil to permit the basic biochemical processes necessary for maintaining soil fertility. Organic management increases overall enzyme activity [3], but activities of specific enzymes may change depending on the composition of the amendments and the relative availability of nutrients, as well as other factors, such as soil type and its unique characteristics [4]. Dehydrogenase activity estimates overall microbial activity due to its presence in all the microorganisms [5]. Phosphatase activity is related to phosphorus cycle in soil. Whereas, higher phosphatase activity increases significantly in typical purple soil irrigated by biogas slurry in China [6]. Therefore, while studying the effect of different levels and sources of fertilization on soil fertility, attention must be focused to the activity of the micro flora and enzymatic activity towards soil biological quality.

In rice (*Oryza sativa*) among the various practices, optimizing the use of manures and fertilizers is one of the important strategies for increasing productivity of rice. The contribution of organic manures is to be judged not only in terms of nutrient contribution, but also by their role in building up nutrient reserves in soil and increasing organic matter level of soil which ultimately improves physical, chemical and biological properties of soil. Using organic sources like farm yard manure, vermicompost, green manure combination with inorganic fertilizer deserves priority for sustained production and better on farm resource recycling and utilization. Organic manure provides nutrients, growth promoting substances, as well as they are microbial enriched products that play a vital role in organic rice production. In the present investigation, an attempt was made to examine the influence of different levels and sources of fertilization on dynamics of soil chemical properties, microbial biomass and

enzyme activities under anaerobic rice cultivation.

## 2. MATERIALS AND METHODS

A field experiment was conducted during *kharif* season at Wetland farm of TNAU, Coimbatore. The initial analysis of the soil of the experimental site (Table 1) revealed that the soil was slightly alkaline (pH = 7.85) with low in soluble salts (EC = 0.42dSm<sup>-1</sup>), medium in organic carbon content (0.58 per cent), low in available N (216 kg ha<sup>-1</sup>), medium in P<sub>2</sub>O<sub>5</sub> (16.2 kg ha<sup>-1</sup>) and high in K<sub>2</sub>O (426 kg ha<sup>-1</sup>). The irrigation water was found to be neutral in reaction (pH = 7.6) with medium level of the soluble salts (EC = 1.18 dSm<sup>-1</sup>).

The maximum and minimum temperature ranged from 29.0 to 32.7°C and 19.2 to 23.7°C, respectively. With regard to relative humidity, there was a fluctuation from 77.5 to 94 per cent (07 22 hours) and from 49.6 to 77.3 per cent (14 22 hours). There was a total rainfall of 110 mm was received in 16 rainy days. The evaporation and bright sunshine hour's day<sup>-1</sup> ranged from 2.8 to 6.6 mm and 3.1 to 7.4 hours, respectively

The study was conducted with nine treatments which are T<sub>1</sub> –100% N through dhaincha + balance P & K through inorganic fertilizers, T<sub>2</sub> – 50% N through dhaincha + balance N, P & K through inorganic fertilizers, T<sub>3</sub> –100% N through vermicompost + balance P & K through inorganic fertilizers, T<sub>4</sub> –50% N through vermicompost + balance N, P & K through inorganic fertilizers, T<sub>5</sub> –100 % NPK (150 : 50 : 50 kg ha<sup>-1</sup>) through inorganic fertilizers, T<sub>6</sub> –100% NPK through inorganic fertilizers + 12.5 t farmyard manure, T<sub>7</sub> – 100% NPK through inorganic fertilizers + 6.25 t dhaincha, T<sub>8</sub> – 100% NPK through inorganic fertilizers + 5 t vermicompost, T<sub>9</sub> – Control. The experiment was laid out in RBD with three replications. The test crop used is medium duration rice variety CO(R) 48.

### 2.1 Green Manure Incorporation

Dhaincha (*Sesbania aculeate* Poir.), a leguminous green manure crop, local variety was used as green manure and raised separately for incorporation as per treatment schedule. *S. aculeata* is an indigenous and a very common green manure crop traditionally grown by Indian farmers in lowland rice. It is a root nodulating, quick growing and succulent green manure, capable of producing 20 to 25 t ha<sup>-1</sup> biomass accumulation with 80 to 120 kg N ha<sup>-1</sup> in 55 days.

It is tolerant to flooding and salinity. Seeds of dhaincha were obtained from Department of Farm Management, Tamil Nadu Agricultural University. The field was thoroughly ploughed with tractor and the seeds were sown @ 25 kg ha<sup>-1</sup>. Dhaincha was incorporated two weeks before transplanting of rice at the age of 48 and 47 days during *samba* season as per treatment schedule. The nutrient value of *Sesbania aculeata* was furnished in Table 1.

### 2.2 Organic Manures

Well decomposed farm yard manure and vermicompost were obtained from Department of Farm Management, Tamil Nadu Agricultural University. Manures required for the experiment were worked out on N equivalent basis are presented in Table 1.

### 2.3 Inorganic Fertilizer Application

In treatments, where chemical fertilizers are to be applied urea (46% N), single super phosphate (16% P<sub>2</sub>O<sub>5</sub>) and muriate of potash (60% K<sub>2</sub>O) were used. Nitrogen was applied in four equal splits at basal, active tillering, panicle initiation and flowering stages. The entire dose of P and 50 per cent of K were applied basally before sowing. Remaining 50 per cent of K was applied at panicle initiation stage.

### 2.4 Soil Chemical Analysis

Soil samples were taken before the start of experiment and harvest of rice for analysis. Pre-sowing composite soil sample was analysed for mechanical and chemical properties. The soil samples were collected from each plot at 0-15 cm depth, dried under shade, powdered, sieved through 2 mm sieve and analysed for pH, EC, organic carbon, macro, secondary and micro-nutrients following the standard procedures as shown in Table 2.

### 2.5 Assessment of Microbial Population

The microbial population in the soil before the start of experiment of the crop was determined. The standard serial dilution plating technique of Pramer and Schemidt [13] was adopted for the estimation of microbial population and expressed as colony forming unit (cfu) g<sup>-1</sup> of soil. The different types of microorganisms were enumerated using differential media favouring the growth of bacteria, fungi and actinobacteria as shown in Table 3.

## 2.6 Assessment of Enzyme Activity

The enzyme activity was determined at initial and postharvest stages of rice. The substrates and methods followed for enzyme assays were presented in Table 4.

## 3. RESULTS AND DISCUSSION

### 3.1 Soil Chemical Analysis

Results on soil chemical analysis were presented in Table 5. Soil available nitrogen, phosphorus and potassium status was also influenced owing to the INM practice, recommended NPK fertilizers and different organic manures application.

### 3.2 Soil Available Nitrogen

The maximum soil available N ( $255 \text{ kg ha}^{-1}$ ) was observed with application 100% NPK through inorganic fertilizers + 6.25 t dhaincha which was comparable with 100% NPK through inorganic fertilizers + 5 t vermicompost and 100% NPK through inorganic fertilizers + 12.5t FYM. Among the other treatments, higher soil available N ( $233 \text{ kg ha}^{-1}$ ) was attained with 100% N through dhaincha + balance P and K through inorganic fertilizers (Table 5). This might be due to lower amount of residual nutrient in inorganic fertilizer applied field. Inorganic fertilizers cause immediate release of nutrients, which will be utilized by the crop or might have lost the environment through leaching or identification process. Higher N availability in the organic manures such as vermicompost, FYM might be due to higher N content and continuous and slow release of nutrients from organic manure and increased biomass and accumulated soil organic matter. Similar findings were also reported by Amanullah [20].

### 3.3 Soil Available Phosphorus

The soil available P status was also improved by the addition of organic and inorganic sources of nutrient management practices (Table 5). Higher soil available P ( $18.5 \text{ kg ha}^{-1}$ ) was attained with 100% NPK through inorganic fertilizers + 6.25t dhaincha which was comparable with 100% NPK through inorganic fertilizers + 5t vermicompost (18.2) and 100% NPK through inorganic fertilizers + 12.5t FYM (17.8). Among the other treatments, 100% N through dhaincha + balance P and K through inorganic fertilizers recorded the

maximum soil available P ( $16.1 \text{ kg ha}^{-1}$ ) compared to control. This might be due to the fact that during the mineralization of enriched organics, a number of organic acids, especially the hydroxyl ions (product of microbial metabolism) are produced, which might have released P through chelation or by removal of metal ions from the insoluble metal phosphates as observed by Mohandas and Appavu [21]. The influence of organic manure in increasing the label P through complexing of cations like  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  responsible for P fixation has been reported by Balaguravaiah [22]. Rock phosphate enriched manures maintained higher levels of P in soil solution for a longer period than the inorganic fertilizer. Higher soil available P could be attributed to decomposition of organic manures in the production of organic acids which in turn stabilize native insoluble P and led to available for longer period. Further, the higher quantity of crop residues might also have contributed P to the soil [23].

### 3.4 Soil Available Potassium

The impact of organic and inorganic sources of nutrient management practices on residual soil K was significant (Table 5). Application of 100 % NPK through inorganic fertilizers + 6.25t dhaincha registered higher soil available K ( $432 \text{ kg ha}^{-1}$ ) and was comparable with 100 % NPK through inorganic fertilizers + 5t vermicompost and 100 % NPK through inorganic fertilizers + 12.5 t FYM. Among the other treatments, significantly higher soil available K ( $406 \text{ kg ha}^{-1}$ ) was attained with 100 % N through dhaincha + balance P and K through inorganic fertilizers. The enhanced K availability irrespective of the season coupled with higher K uptake due to organic manure incorporation could be attributed to higher DMP and K absorption, evidencing the priming effect of K contribution by organic manure [24]. Among the organic manures, *Sesbania aculeata* played a vital role in improving the uptake of NPK. This might be due to the fact that quick release of N from the added green manure with increased availability of P through the mechanism of reduction, chelation and favourable changes in soil pH and K through the priming effect and besides the direct contribution of K by green manure [25]. Higher K uptake in rice might be due to the increase in available K, which might have contributed to mineralization of organic manures or solubilization of nutrients from native sources during decomposition [26].

### 3.5 Soil Available Micronutrients

The available soil micronutrients like Iron (Fe), Manganese (Mn), Zinc (Zn) and Copper (Cu) status were presented in Table 6. Application of 100% NPK through inorganic fertilizers + 6.25t dhaincha observed that maximum uptake of Fe (2582 g ha<sup>-1</sup>), Mn (3639 g ha<sup>-1</sup>), Zn (380 g ha<sup>-1</sup>) and Cu (89.4 g ha<sup>-1</sup>) and it was followed by 100 per cent RDN through green manure and the lowest soil available micronutrients reported with absolute control. Among the other treatments, 100% NPK through inorganic fertilizers + 5t vermicompost and 100 % NPK through inorganic fertilizers + 12.5 t FYM which were comparable with each other. Addition of organic materials might have enhanced the microbial activity in the soil and the consequent release of complex organic substances (Chelating agents) could have prevented micronutrients from precipitation,

fixation, oxidation and leaching and also addition of these nutrients through organic sources. Suvarna Latha and Sankara Rao [27] stated that organic manure increased the available Fe, Mn, Zn and Cu gradually from tillering to harvest stage.

### 3.6 Soil Microbial Population

The INM practice, organic manures and recommended NPK fertilizers had influenced bacterial, fungal and actinobacterial population at postharvest stage of the crop (Table 7). 100 per cent N through dhaincha + balance P and K through inorganic fertilizers was recorded higher bacterial population (40.4 x 10<sup>6</sup> CFU g<sup>-1</sup> of dry soil), fungal population (17.5 x 10<sup>3</sup> CFU g<sup>-1</sup> of dry soil), actinobacterial population (13.9 x 10<sup>4</sup> CFU g<sup>-1</sup> of dry soil) which was followed by 100 % NPK through inorganic fertilizers + 6.25 t dhaincha.

**Table 1. Nutrient content of organic manures on dry weight basis**

S. no.	Manures	Nutrient content (%)		
		N	P	K
1.	Farmyard manure	0.60	0.40	0.62
2.	Vermicompost	1.81	0.64	1.20
3.	<i>Sesbania aculeata</i> L.	2.67	0.68	1.26

**Table 2. Soil chemical characteristics of the pre experimental field**

Particulars	Value	Method	Reference
<b>II. Chemical characteristics</b>			
pH	7.85	Using glass electrode in the "ELICO" pH meter	[7]
Electrical conductivity (dS m <sup>-1</sup> )	0.42	Using "ELICO" conductivity bridge	
Organic carbon (%)	0.58	Chromic acid wet digestion method	[8]
Available N (kg ha <sup>-1</sup> )	216	Alkaline permanganate method	[9]
Available P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	16.2	Olsen method using colorimeter	[10]
Available K <sub>2</sub> O (kg ha <sup>-1</sup> )	426	Flame photometric method	[11]
DTPA-Fe (mg kg <sup>-1</sup> )	27.8	Atomic Absorption Spectrophotometry	
DTPA-Cu (mg kg <sup>-1</sup> )	3.2		
DTPA-Mn (mg kg <sup>-1</sup> )	8.6		[12]
DTPA-Zn (mg kg <sup>-1</sup> )	3.8		
<b>II. Irrigation water characteristics</b>			
PH	7.6	Potentiometry	[7]
Electrical conductivity (dS m <sup>-1</sup> )	1.18	Conductometry	

**Table 3. Soil biological properties of the pre experimental field**

Particulars	Value	Method	Reference
<b>I. Biological properties</b>			
Total bacteria (cfu x 10 <sup>6</sup> g <sup>-1</sup> of soil)	18	Serial dilution method using Nutrient glucose Agar medium	[14]
Total fungi (cfu x 10 <sup>3</sup> g <sup>-1</sup> of soil)	10	Serial dilution method using Martin's Rose Bengal Agar medium	[15]
Total actinobacteria (cfu x 10 <sup>4</sup> g <sup>-1</sup> of soil)	4	Serial dilution method using Kenknight's Agar medium	[16]

**Table 4. Standard methods followed for soil enzyme analysis of the pre experimental field**

Enzyme	Value	Substrate	Method	Reference
Dehydrogenase (µg of TPF released g <sup>-1</sup> of soil 24 h <sup>-1</sup> )	8.22	2,3,5- Triphenyl Tetrazolium chloride	Spectrophotometer at 485 nm	[17]
Urease (µg NH <sub>4</sub> <sup>+</sup> g <sup>-1</sup> of soil 24 h <sup>-1</sup> )	27.68	10 per cent urea solution	Spectrophotometer at 630 nm	[18]
Phosphatase (µg of p-nitrophenol released g <sup>-1</sup> of soil h <sup>-1</sup> )	12.02	p-nitrophenol phosphate	Spectrophotometer at 420 nm	[19]

**Table 5. Effect of integrated nutrient management on soil available macro nutrients (kg ha<sup>-1</sup>) at harvest stage of rice**

Treatments	Nitrogen	Phosphorous	Potassium
T <sub>1</sub>	233	16.1	406
T <sub>2</sub>	211	14.1	381
T <sub>3</sub>	212	14.4	381
T <sub>4</sub>	211	13.9	380
T <sub>5</sub>	210	13.5	380
T <sub>6</sub>	254	17.8	431
T <sub>7</sub>	255	18.5	432
T <sub>8</sub>	254	18.2	431
T <sub>9</sub>	189	11.7	345
SEd	9	0.7	11
CD (P=0.05)	19	1.4	24

**Table 6. Effect of integrated nutrient management on soil available micro nutrients (g ha<sup>-1</sup>) at harvest stage of rice**

Treatments	Zn	Cu	Fe	Mn
T <sub>1</sub>	345	80.2	2338	3297
T <sub>2</sub>	280	64.6	1776	2895
T <sub>3</sub>	299	71.6	2007	2972
T <sub>4</sub>	273	61.4	1990	2506
T <sub>5</sub>	264	59.8	1637	2311
T <sub>6</sub>	313	72.4	2116	2985
T <sub>7</sub>	380	89.4	2582	3639
T <sub>8</sub>	317	73.8	2152	2972
T <sub>9</sub>	199	52.4	1500	2118
SEd	22	6.1	180	252
CD (p=0.05)	46	12.4	378	528

**Table 7. Effect of integrated nutrient management on soil microbial population at harvest stage of rice**

Treatments	Bacterial population (x 10 <sup>6</sup> CFU g <sup>-1</sup> )	Fungal population (x 10 <sup>3</sup> CFU g <sup>-1</sup> )	Actinobacterial population (x 10 <sup>4</sup> CFU g <sup>-1</sup> )
T <sub>1</sub>	40.4	17.5	13.9
T <sub>2</sub>	37.1	14.3	11.3
T <sub>3</sub>	37.4	13.9	12.5
T <sub>4</sub>	36.5	15.6	11.2
T <sub>5</sub>	29.4	13.4	10.8
T <sub>6</sub>	37.8	15.9	12.9
T <sub>7</sub>	39.5	16.5	13.6
T <sub>8</sub>	38.8	16.1	13.1
T <sub>9</sub>	23.2	9.5	6.2
SEd	3.4	1.3	1.1
CD (p=0.05)	7.2	2.8	2.3

Among the other treatments, highest microbial population were registered with application of 100% NPK through inorganic fertilizers + 5 t vermicompost and it was on par with 100% NPK through inorganic fertilizers + 12.5 t FYM. The lowest microbial population (fungi, bacteria and actinobacterial) was recorded with absolute control. In the present study, the activity of soil microbial population was increased due to the application of organic manures and green manure with the advancement of crop growth. It could be due to enhanced organic carbon content of the soil as a result of organic manure application as compared to inorganic fertilizers [27]. Lower soil microbial load found in recommended NPK fertilizers might be due to inhibitory nature of chemical fertilizers on the growth and development of microbes [28].

### 3.7 Soil Enzyme Activity

The soil enzyme activity was influenced by the INM practice, organic manures and

recommended NPK fertilizers application in (Table 8).

1. Urease ( $\mu\text{g NH}_4^+ \text{g}^{-1}$  of soil 24 h<sup>-1</sup>)
2. Dehydrogenase ( $\mu\text{g}$  of TPF released g<sup>-1</sup> of soil 24 h<sup>-1</sup>)
3. Phosphatase ( $\mu\text{g}$  of p-nitrophenol released g<sup>-1</sup> of soil 24 h<sup>-1</sup>)

Higher urease activity ( $46.8 \mu\text{g NH}_4^+ \text{g}^{-1}$  soil 24 h<sup>-1</sup>), soil dehydrogenase activity ( $36.8 \mu\text{g}$  of TPF released g<sup>-1</sup> of soil 24 h<sup>-1</sup>), soil phosphatase activity ( $38.8 \mu\text{g}$  of p - nitrophenol released g<sup>-1</sup> of soil 24 h<sup>-1</sup>) were observed with 100 per cent N through dhaincha + balance P and K through inorganic fertilizers followed by 100 % NPK through inorganic fertilizers + 6.25 t dhaincha. The lowest soil enzyme activities (urease activity @  $29.2 \mu\text{g NH}_4^+ \text{g}^{-1}$  soil 24 h<sup>-1</sup>, soil dehydrogenase activity @  $20 \mu\text{g}$  of TPF released g<sup>-1</sup> of soil 24 h<sup>-1</sup>, soil phosphatase activity @  $22.4 \mu\text{g}$  of p - nitrophenol released g<sup>-1</sup> of soil 24 h<sup>-1</sup>) were registered with absolute control. The

**Table 8. Effect of integrated nutrient management on soil enzyme activity at harvest stage of rice**

Treatments	Urease	Dehydrogenase	Phosphatase
T <sub>1</sub>	46.3	36.3	38.3
T <sub>2</sub>	39.1	28.1	32.0
T <sub>3</sub>	40.5	30.5	32.5
T <sub>4</sub>	37.5	26.9	30.1
T <sub>5</sub>	36.0	24.8	26.9
T <sub>6</sub>	41.7	32.9	32.4
T <sub>7</sub>	45.4	35.6	36.9
T <sub>8</sub>	43.2	34.7	35.7
T <sub>9</sub>	29.2	20.0	22.4
SEd	3.6	2.81	3.1
CD (P=0.05)	7.9	5.98	6.42

organic manures particularly *sesbania aculatae* improves the soil urease and dehydrogenase activities and which might have improved improves soil phosphatase activities too. High organic carbon content in soil applied with dhaincha stimulated the soil microorganisms by serving as source of carbon, energy and other nutrients essential for their growth and multiplication and the increased the soil activities [4]. Higher phosphatase activity was observed with enriched organic manures treated with rock phosphate [5].

## 5. CONCLUSION

The maximum enzyme activity in soil under organic nutrient management practice due to incorporation of organic manures that increase decomposition process thereby increase the microbial activity & chemical properties. Based on the results, It can be concluded that higher microbial population (bacteria, fungal & actinomycetes), soil enzyme dynamics (urease, dehydrogenase & soil phosphatase activity), soil available macro nutrients (N, P, K) & micro nutrients (Fe, Zn, Mn, Cu) were significantly influenced with application of 100% N through dhaincha + balance P & K through inorganic fertilizers followed by application of 100% NPK through inorganic fertilizers + 6.25 t dhaincha. These integrated nutrient management practices seem to be better option for western zone of Tamil Nadu rice farmers.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Jackson LE, Bowles TM, Hodson AK and Lazcano C 2012. Soil microbial- root and microbial rhizosphere processes to increase nitrogen availability and retention in agroecosystems. *Curr. Opin. Environ. Sustain.* 2012;4:517-522.
- Kallenbach C and Grandy AS 2011. Control over soil microbial biomass responses to carbon amendments in agricultural systems: A meta-analysis. *Agric. Ecosyst. Environ.* 2011;144:241-252.
- Moeskops BD, Buchan S, Sleutel L, Herawaty E, Husen R, Saraswati D et al. Soil microbial communities and activities under intensive organic and conventional vegetable farming in West Java, Indonesia. *Appl. Soil Ecol.* 2010;45:112-120
- Stursova M, Baldrian P. Effects of soil properties and management on the activity of soil organic matter transforming enzymes and the quantification of soil-bound and free activity. *Plant and Soil*, 2010;338:99-110.
- Taylor JP, Wilson B, Mills MS, Burns RG. Comparison of microbial number and enzymatic activities in surface soils and subsoil using various techniques. *Soil Biol. Biochem.* 2002;34:387-401.
- Shiling C, Weiwei Y, Zhi Z, Surong L. Soil properties and enzyme activities as affected by biogas slurry irrigation in three gorges reservoir areas of China. *J. Environ. Biol.* 2015;36:513-520.
- Jackson ML 1973. *Soil chemical analysis.* Prentice Hall of India Pvt. Ltd., New Delhi.
- Walkley A, Black CA. An examination of the degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.* 1934;40:233-243.
- Subbiah BV, Asija GL. A rapid procedure for estimation of available nitrogen in soils. *Curr. Sci.* 1956;25:259-260.
- Olsen SR, Cole CV, Watanabe FS, Dean AL. Estimation of available phosphorus in soils by extraction with Sodium bicarbonate Circular no: 939, USDA; 1954.
- Stanford G, English L. Use of flame photometer in rapid soil test for K and Ca. *Agron J.* 1949;41: 446.
- Lindsay WL; Norvell WA. Development of DTPA soil test for zinc, iron, manganese and copper. *Soil Sci. Soc Am J.* 1978;42:421-428.
- Pramer D, Schemidt EL. *Experimental soil microbiology.* Burges Publishing Co. Minneapolis, Minnesota, 1965; 107.
- Collings CH, Lyne MP. *Microbiological methods.* 5<sup>th</sup> edition, Butter Worth, London; 1968.
- Martin JP. Use of acid, rose Bengal and streptomycin in the plate method for estimating soil fungi. *Soil Sci.* 1950;69: 215-233.
- Kenknight G, Muncie JH. Isolation of phytopathogenic actinomycetes. *Phytopathology.* 1939;29: 1000-1001.
- Casida LE Jr, Klein DA, Santoro T. Soil dehydrogenase activity. *Soil Sci.* 1964;98:371-376.
- Tabatabai MA, Bremner JM. Use of P-nitrophenyl phosphate for assay of soil

- phosphatase activity. Soil Biol. Biochem. 1969 ;1: 301-307.
19. Halstead RL. Phosphatase activity of soils as influenced by lime and other treatments. Canadian. J Soil Sci. 1964;44: 137-144.
  20. Amanullah MM, Yassin MM, Somasundaram E, Vaiyapuri K, Sathyamoorthi K, Pazhanivelan S. N availability in fresh and composted poultry manure. Res. J Agric Biol Sci. 2006b;2(6): 406-409.
  21. Mohandas S, Appavu K. Direct and residual effect of combined application of basic slag with green leaf manures on soil available nutrients and yield of rice. Madras Agric J. 2000;87(1-3):53-56.
  22. Balaguravaiah D, Adinarayana G, Prathap S, Yellamanda Reddy T. Influence of long-term use of inorganic and organic manures on soil fertility and sustainable productivity of rainfed groundnut in *alfisols*. J. Indian Soc. Soil Sci. 2005; 53(4): 606-611.
  23. Pazhanivela S, Amanullah MM, Vaiyapuri K, Sharmila Rahale C, Sathyamoorthi K and Alagesan A. Effect of rock phosphate incubated with FYM on nutrient uptake and yield of lowland rice. Res. J. Agric. Biol. Sci. 2006;2(6): 365-368.
  24. Shobarani N, Prasad GSV, Prasad ASR, Sailaja B, Muthuraman P, Numeera S, et al. Rice Almanac India. Technical Bulletin No 5, DRR, Rajendranagar, Hyderabad. 2010;6-7.
  25. Geethalakshmi V. Studies on the direct and residual effect of non-conventional green leaf manures, with N for rice-rice cropping system. Ph.D. Thesis, Tamil Nadu Agric. Univ., Coimbatore; 1996.
  26. Suvarna Latha AJL, Sankara Rao VR. Integrated use of fertilizers and poultry manure on nutrient availability and yield of rice. J. Indian Soc. Coastal Agric. Res. 2001;19(1&2) :153-157.
  27. Krishnakumar S, Saravanan A, Rajesh V, Mayil Samy P. Impact of organic farming on biological properties of rice (*Oryza sativa*) grown soil. J. Ecobiol. 2007;20(3): 275-280.
  28. Singh YV, Singh KK, Sharma SK. Influence of crop nutrition and rice varieties under two systems of cultivation on grain quality, yield and water use. Rice Science. 2012;19(4):1-10.

© 2020 Ajaykumar and Thavaprakash; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle4.com/review-history/64901>