



Comparative Analysis of Health Risk Associated with Occupational Exposure to Formaldehyde in Public and Private Mortuaries in Rivers State, Nigeria

Obed-Whyte, Roland^{1*}, K. E. Douglas¹ and Nte, Alice¹

¹Centre for Occupational Health, Safety and Environment, University of Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author OWR designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors KED and NA managed the analyses of the study. Author NA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2019/v15i230115

Editor(s):

(1) Dr. Janvier Gasana, Professor, Department of Environmental & Occupational Health, EO Epidemiology, and EO Medicine, Robert Stempel College of Public Health & Social Work, Florida International University, USA.

Reviewers:

(1) Venkata Sanyasi Seshendra Kumar Karri, GITAM University, India.

(2) Mai Sabry Saleh, National Research Centre, Egypt.

(3) Ibama, Asiton-a Solomon, Federal University of Technology, Nigeria.

(4) Basem Sabry, National Research Centre, Egypt.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48794>

Received 28 February 2019

Accepted 10 May 2019

Published 24 May 2019

Original Research Article

ABSTRACT

Background: Formaldehyde (FA) is a well-known chemical widely used in mortuaries in Nigeria for the preservation of human cadavers, yet little is known of the potential health risk associated with occupational exposure to formaldehyde in mortuaries. This study evaluated the potential health risk associated with occupational exposure to formaldehyde in mortuaries in Rivers State, Nigeria.

Methodology: The study was carried out in 7 public and 8 private mortuaries and the concentrations of formaldehyde to which the morticians are exposed were measured during the embalming process. Modeling of health related risk was carried out in accordance with methods recommended by the United States Environmental Protection Agency (US EPA).

Results: The results showed that the lethal concentrations of formaldehyde in the mortuaries far exceeded the "No Significant Risk Levels" (LC₅₀ = 3.3 ppm for public mortuaries; and 3.46 ppm for

*Corresponding author: E-mail: rollyandwhyte@yahoo.com;

private mortuaries). Analysis showed that 77.2% of workers in the public mortuaries have high daily formaldehyde exposure index, while 88.24% of the workers in the private mortuaries have high daily formaldehyde exposure index. The difference between the formaldehyde daily exposure index and daily potential dose in public and private mortuaries was not statistically significant ($p > 0.05$). Computed hazard quotients for both public and private were 1.25 and 3.0 respectively (> 1). Computed cancer related risk values for public and private mortuaries were 1.5×10^{-3} and 1.9×10^{-3} respectively.

Conclusion: The study showed that embalmers in both the public and private mortuaries in Rivers State occupationally exposed to formaldehyde have significant risk of developing carcinogenic and non-carcinogenic related health problems. It is therefore, recommended that operators of mortuaries and Rivers State Government should provide FA monitoring device and continuous health education for workers.

Keywords: Formaldehyde; mortuaries; carcinogenic; non-carcinogenic.

1. INTRODUCTION

The health care system offers various services, including mortuary services, to the society [1]. Mortuaries receive corpses, embalmed/preserved and finally deliver them to their respective owners [2]. These different activities involved in mortuary services expose the workers to occupational hazards with their associated health risks. These hazards include exposure to hazardous chemicals (such as formaldehyde, paraformaldehyde glutaraldehyde and methanol) and infectious diseases [2]. Physical, chemical and radiation risks have been identified as some of the occupational health and safety (OHS) risks associated with the operations of mortuaries [2-3]. Also, chemical, physical and biological hazards have been said to pose serious risk to mortuary workers [1].

Hazardous substances such as FA used in the mortuaries can enter the body by inhalation or through the skin contact [4]. Exposure to FA during embalment is one of the occupational hazards in mortuary services as continuous inhalation of FA poses adverse risk to the health of the morticians, or aggravates their existing health problems [4]. The Occupational Safety and Health Administration (OSHA), National institute for occupational safety and health (NIOSH) and other regulatory bodies and the World Health Organization (WHO) have put formaldehyde exposure limits for workers at short times and at longer durations [5]. Scientific evidences, both in experimental animals and humans have shown that exceeding these exposure limits have some adverse health implications.

The risk associated with FA inhalation can be short-term or long-term risk or both. Acute or

short-term health effects of FA exposure include eye and throat irritation and respiratory symptoms; while chronic or long-term health effects include chest tightness, cancers, swelling or spasms in the throat (glottis) and severe coughing [6]. Continual and prolonged exposure to formaldehyde has been associated with lung and nasal passage cancers and myeloid leukemia in humans [7-9]. Short-term and long-term exposure to formaldehyde is highly irritating to the upper respiratory tract and can cause respiratory symptoms, throat, nose and eye and irritations [8,10]. It has been reported that men, such as mortuary workers, who are exposed to high levels of formaldehyde, are at much greater risk of dying from Amyotrophic Lateral Sclerosis (ALS), otherwise called Lou Gehrig's disease [11]. He stated further that morticians who are continuously exposed to high levels to formaldehyde are almost 4.5 times more likely to die from ALS than those who are not exposed to formaldehyde in their workplaces. Another study [1] also reported that workers in mortuaries, particularly embalmers, are exposed to high concentration of formaldehyde above 0.75ppm threshold limit resulting in eye irritation and coughing.

In Nigeria, mortuary services are provided by both public sector (through government own hospitals) and the private sector. Formaldehyde exposure has been identified as one of the common hazards that constitute risk to the health of workers in mortuaries in Port Harcourt [12]. High concentrations of FA that far exceeded stipulated OSHA limit have been reported in public and private mortuaries in Rivers State, Nigeria [13]. It was further stated that the high levels of FA obtained in mortuaries pose short-term and long-term risk to workers [13]. The aim of this study was to carry out a comparative

analysis of the health risks associated with occupational exposure to FA in public and private mortuaries in Rivers State, Nigeria. This study attempted to determine the short-term and long-term risk levels associated with exposure to FA in public and private mortuaries.

2. METHODOLOGY

The study was carried out in 7 public mortuaries and 8 private mortuaries given a total of fifteen mortuaries as indicated. Concentrations of FA gas in the embalment sections of the mortuaries were measured using a Globe Instrument, model PGas-20 CH₂O gas detector [13]. The age, body weight, working time per day, and employment duration of the morticians were obtained and used for the health risk analysis. The study employed both semi-quantitative and quantitative risk assessment approach to determine health risk exposure to FA in mortuaries. The semi-quantitative risk assessment was based on the hazards rating and ranking of FA [14-15]. The quantitative risk assessment approach uses mainly mathematical relationships between variables based on the stipulated methods and guidelines by [16-17].

Data analysis was carried out using Microsoft Excel. Mean and standard deviation were computed and data were presented in either tables or graphs. The levels of significance in the formaldehyde daily exposure index (DEI) and daily potential dose (DPD) between public and private mortuaries were determined using analysis of variance (ANOVA) in Microsoft Excel

2.1 Determination of Lethal Concentration

Lethal concentration is the amount of formaldehyde concentration that proves fatal to the exposed mortuary workers. The values of formaldehyde concentrations and percentage of time it was equal to or exceeds the threshold limit were estimated using a linear regression technique presented in Equation (1).

$$LC_i = \alpha + \beta T_i \quad (1)$$

Where: LC_i is the formaldehyde concentration for a particular percentage of time in part per million (ppm), T_i is the percentage of time (%), α and β are coefficients of regressions. The formaldehyde lethal concentrations for both the public and private mortuaries were computed by

ranking the formaldehyde concentrations using the Weibull ranking approach. The corresponding lethal concentrations equal to or exceeded the threshold limit was determined and estimated from the plots of ranked observed concentrations versus the percentage of time exceeded or equal to threshold value. The lethal concentrations model was derived from the linear plots shown in Figs. 3 and 4 as follows:

$$LT_x = \alpha + \beta \ln(LC_x) \quad (2)$$

Where: LT_x is the percentage of time exceeded that proof lethal, LC_x is the lethal concentration (ppm), α and β are constants.

Precisely, LC_x is the lethal concentration of the formaldehyde over which a mortuary worker is exposed for some period of time.

From Equation (2), the lethal concentration is estimated as follows:

$$LC_x = Exp\left(\frac{LT_x - \alpha}{\beta}\right) \quad (3)$$

2.2 Semi-Quantitative Health Risk Assessment

The health risks associated with the exposure of morticians in both public and private mortuaries to formaldehyde were further assessed using a semi-quantitative approach [14,18-20]. The exposure rate and risk rate were computed using Equations (4) and (5) respectively. The formaldehyde exposure level (FEL) in the mortuaries was calculated using average concentrations of formaldehyde and the average duration each worker is exposed as well as the frequency of exposure as given in Equation (4) [14].

$$FEL = \frac{EF \times ED_{avg} \times C_{avg}}{W_{havg}} \quad (4)$$

Where:

- FEL = Formaldehyde Exposure level (ppm)
- EF = Exposure frequency per week
- ED_{avg} = average duration of each exposure (hours)
- C_{avg} = average concentration (ppm)
- W_{havg} = average working hours per week

The exposure rating (ER) was determined by comparing the formaldehyde exposure level (FEL) with the permissible exposure limit (PEL) as shown in Table 1.

Table 1. Exposure rating of formaldehyde

| FEL/PEL | Exposure rating (ER) |
|--------------|----------------------|
| < 0.1 | 1 |
| 0.1 to < 0.5 | 2 |
| 0.5 to < 1.0 | 3 |
| 1.0 to < 2.0 | 4 |
| ≥ 2.0 | 5 |

Source: [14-15]

The exposure rating (shown in Table 1) are represented in an ordinal scale of 1 to 5 categorized in the order of severity of exposure, so that 1 indicates very low exposure, 2 indicates low exposure, 3 indicates moderate, 4 indicates high exposure and 5 indicates very high exposure [19]. The exposure indices were rated and the risk calculated using Equation (5) [15, 19-20].

$$\text{Risk Rating} = \sqrt{\text{HR} \times \text{ER}} \quad (5)$$

Where HR is the formaldehyde hazard rating and ER is the Exposure rating.

Formaldehyde hazard rating (HR) is given as 4 in [14-15].

The risk for each mortuary category was ranked to determined levels of significance based on risk level shown in Table 1 [15] and risk ranking shown in Table 2 [19].

Table 2. Risk ranking level of formaldehyde

| Risk rating | Risk ranking |
|-------------|--------------|
| 1 | Very low |
| 2 | Low |
| 3 | Moderate |
| 4 | High |
| 5 | Very high |

Source: [19]

2.3 Quantitative Health Risk Assessment

2.3.1 Determination of daily exposure index (DEI)

The Formaldehyde daily exposure index for each exposed worker in both the public and private mortuaries was computed using the average formaldehyde concentrations and the OSHA

occupational exposure limit (OEL) of 0.75ppm as given in Equation 6 [17]. The formaldehyde daily exposure index was computed using Equation (4) modified from [14] and [15]. The Daily exposure index was scaled such that DEI less than 0.1 is considered as very low, DEI between 0.1 and 0.5 is considered as low, DEI between 0.5-1.0 is considered as moderate, DEI between 1.0 and 1.5 is considered as high, and DEI between 1.5 and 2.0 is considered as very high.

$$DEI = \frac{C \times ET(\text{hr})}{OEL \times 24(\text{hr})} \quad (6)$$

Where:

- C = concentration (mg/m³)
- ET = exposure time (hr)
- OEL = OSHA occupational exposure limit

2.3.2 Determination of Daily Potential Dose (DPD)

The Formaldehyde daily potential dose for each exposed worker in both the public and private mortuaries was computed using Equation 7.

$$DPD = \frac{C \times IR \times ET(\text{hr})}{24(\text{hr})} \quad (7)$$

Where:

- DPD = daily potential dose (mg/d)
- C = average formaldehyde concentration (mg/m³)
- IR = the inhalation rate (16m³/day)
- ET = daily exposure time (hour)

An inhalation rate (IR) of 16m³/day was adopted in this study [21].

Equations 4 and 5 assume that there is no exposure when embalment is not carried out.

2.4 Modeling the Non-Carcinogenic and Carcinogenic Risk

Formaldehyde health risk assessment was carried out for non-cancer and cancer related risk. The modeling approach used in this study was adopted from the recommended method by the United States Environmental Protection Agency [21].

2.4.1 Modeling Non-Cancer Related Risk (NCRR)

Non-cancer related risk assessment is carried out to evaluate the short-term or acute health

effects of formaldehyde exposure on mortuary workers. The average daily dose (ADD) and formaldehyde (Hazard) quotient (HQ) were used to evaluate the short-term non-carcinogenic effects of formaldehyde on the exposed morticians. The average daily dose (ADD) was used to evaluate different health effects other than cancer. It was computed by averaging the daily potential dose (DPD) over the body weights and the averaging time as shown Equation (8) [17].

$$ADD = \frac{\text{DailyPotentialDose(DPD)}}{\text{Body Weight}} = \frac{\text{DPD(mg)}}{\text{BW(kg)}} \quad (8)$$

Dose rate averaged over a pathway-specific period of exposure expressed as a daily dose on a per-unit-body-weight basis. The ADD is used for exposure to chemicals with non-carcinogenic or non-chronic effects [17]. The ADD unit is stated in terms of mass/mass-time or mg/kg/day.

Hazard quotient (HQ) method of risk characterization was also used to evaluate non-cancer risk of inhalational exposure to formaldehyde. The hazard quotient (HQ) was computed using Equation (9).

$$HQ = \frac{\text{Intake(mg/kg/d)}}{\text{ReferenceDose(mg/kg/d)}} \quad (9)$$

$$HQ = \frac{\text{ADD(mg/kg/d)}}{\text{RfD(mg/kg/d)}}$$

HQ less than 1.0 (HQ < 1.0) is within safe threshold, while HQ greater 1.0 (HQ > 1.0) is above safe threshold [22]. Reference Dose (RfD) is set up based on health risk assessments.

2.4.2 Modeling Cancer Related Risk (CRR)

The cancer related risk is computed using lifetime average daily doses (LADD). A method of computing the long-term carcinogenic effects of formaldehyde using lifetime average daily dose (LADD) has been recommended [16]. The LADDs for both the public and private mortuaries were computed using Equation (10).

2.4.3 Lifetime average daily dose (LADD)

This is the dose rate averaged over a lifetime. The LADD is used to compute the carcinogenic

or chronic effects of formaldehyde. The LADD unit is also stated in terms of mg/kg/day [17].

$$LADD = \frac{C \times IR \times ED}{BW \times ALT} \quad (10)$$

Where:

C = formaldehyde concentration (mg/m³)
 IR = inhalation rate (16m³/day)
 ED = exposure duration (years)
 BW = body weight (kg)
 ALT = average lifetime (years)

Although the computation of LADD over a lifetime of 70 years has been recommended [17], in this study, a life expectancy of 55 years for male gender in Nigeria as reported [23] was used to compute LADD.

The Cancer related risk (CRR) associated with the inhalation of formaldehyde exposure was computed using the carcinogenic slope factor (CSF) [24] as presented in Equation (11).

$$\text{Cancer Related Risk (CRR)} = \text{Intake (mg/kg/d)} \times \text{carcinogenic slope factor (mg/kg/d)}^{-1}$$

$$CRR = LADD \text{ (mg/kg/day)} \times \text{CSF (mg/kg/day)}^{-1} \quad (11)$$

The non-carcinogenic reference dose (RfD) and carcinogenic slope factor (CSF) are as 0.2 mg/kg/day and 0.021(mg/kg/day)⁻¹ respectively [24].

3. RESULTS

The demographic characteristics of the mortuary workers in the public and private mortuaries are presented in Table 3. The average concentrations of formaldehyde obtained in public and private mortuaries are shown in Table 4. The values of the lethal concentrations for both the public and private mortuaries were estimated as shown in Table 5. Plots of percentage of time the concentrations Equal to or exceeded Threshold concentration in public and private mortuaries are presented in Figs. 1 and 2 respectively. Result of semi-quantitative health risk analysis in public and private mortuaries is presented in Table 6. The computed formaldehyde daily exposure indices for morticians in the mortuaries are shown in Table 7. The computed daily potential dose is shown in Table 8. The results of DEI and DPD normality

test are presented in Figs. 3 and 4 respectively. The variation of daily potential dose with time is presented in Fig. 5.

The demographic characteristics of the mortuary workers in the public and private mortuaries (Table 1) showed that a mean age of 33 years for public and 35 years for private mortuaries. Average length of exposures for workers public

and private mortuaries are 7.5 years and 5.4 years respectively. Average body weights were 76 kg and 74 kg for public and private mortuaries respectively. Workers in the public mortuaries spent an average of 8 hours per day, while workers in the private mortuaries spent an average of 10 hours per day.

Table 3. Demographic characteristics of the mortuary workers

| Characteristics | Public mortuaries | Private mortuaries |
|-------------------------------------|-------------------|--------------------|
| Average Age (years) | 33 | 34.8 |
| Average Employment duration (years) | 7.5 | 5.4 |
| Average Body weight (kg) | 76.4 | 74.6 |
| Average Working time (h/day) | 8 | 10 |

Table 4. Average concentrations of formaldehyde in the mortuaries

| Mortuary category | Minimum (ppm) | Maximum (ppm) | Mean (ppm) | Stdv. (ppm) | OSHA limit |
|--------------------|---------------|---------------|------------|-------------|------------|
| Public Mortuaries | 0.0 | 8.25 | 2.42 | 1.77 | 0.75 |
| Private Mortuaries | 1.18 | 4.58 | 2.52 | 0.99 | 0.75 |

Stdv. = standard deviation

Table 5. Computed lethal concentrations for public and private mortuaries

| | LC ₅₀ (ppm) | LC ₇₅ (ppm) | LC ₉₀ (ppm) | LC ₉₅ (ppm) | OSHA PEL |
|-----------------|------------------------|------------------------|------------------------|------------------------|----------|
| Public Morgues | 3.40 | 2.81 | 2.51 | 2.41 | 0.75 |
| Private Morgues | 3.33 | 2.98 | 2.79 | 2.73 | 0.75 |

Table 6. Result of semi-quantitative health risk analysis

| | Average conc. (mg/m ³) | Exposure level (mg/m ³) | Exposure Rate (ER) | Risk rating | Risk ranking |
|----------------|------------------------------------|-------------------------------------|--------------------|-------------|--------------|
| Public Morgue | 2.97 | 4.24 | 5 | 4.5 | Very high |
| Private Morgue | 3.09 | 2.49 | 5 | 4.5 | Very high |




Fig. 1. Percentage of time equal to or exceeded threshold versus concentrations in public mortuaries




Fig. 2. Percentage of time equal to or exceeded threshold versus concentrations in private mortuaries

Results (presented in Table 4) showed that the average concentrations of formaldehyde obtained in public mortuaries varied between 0.0 ppm and 8.25 ppm with a mean of 2.42 ppm; while concentrations obtained in private mortuaries varied between 1.18 ppm and 4.58 ppm with a mean of 2.52 ppm.

-227.8 respectively for private mortuaries. Substituting the values of α and β into Equation (3), the expressions to estimate lethal concentrations of FA in public and private mortuaries were derived as shown in Equations (12) and (13) respectively.

Table 7. Computed formaldehyde daily exposure index

| Public mortuaries (DEI) | Private mortuaries (DEI) |
|-------------------------|--------------------------|
| 1.61 | 1.4 |
| 1.61 | 1.12 |
| 1.08 | 1.4 |
| 1.61 | 1.12 |
| 1.35 | 1.12 |
| 0.54 | 1.12 |
| 1.61 | 1.68 |
| 0.54 | 1.68 |
| 1.35 | 1.4 |
| 1.08 | 1.4 |
| 1.35 | 1.4 |
| 1.35 | 0.56 |
| 0.54 | 1.4 |
| 1.08 | 1.12 |
| 1.61 | 1.4 |
| 1.08 | 1.12 |
| 1.61 | 1.68 |
| 0.54 | 1.12 |
| 1.61 | 1.68 |
| 0.54 | 1.12 |
| 1.61 | 1.68 |
| 1.35 | 1.68 |

$$LC_x = \text{Exp}\left(\frac{LT_x - 210.71}{-131.3}\right) \tag{12}$$

$$LC_x = \text{Exp}\left(\frac{LT_x - 323.8}{-227.8}\right) \tag{13}$$

The values of the lethal concentrations for both the public and private mortuaries were then estimated from Equations (12) and (13) as shown in Table 5. These values far exceeded the “No Significant Risk Levels (NSRLs)” of 0.0326ppm or 40.0 $\mu\text{g}/\text{m}^3$ [24].

Semi-quantitative analysis shows an exposure rating of 5 for both public and private mortuaries, this gives a risk rating of 4.5 which is ranked as very high as shown in Table 6. Similar result was obtained by [12]. This implies that the formaldehyde exposure in both public and private mortuaries in Rivers State poses very high health risk to morticians/embalmers.

Normality test showed that DEI distribution in the mortuaries does not follow a normal distribution (Fig. 3). The computed formaldehyde daily exposure index for morticians in public mortuaries ranged from 0.54 to 1.61 with a mean deviation of 1.21 \pm 0.42; while the computed DEI for embalmers in private mortuaries ranged from 0.56 to 1.68 with a mean and standard deviation of 1.34 \pm 0.29. The result (Table 7) showed that 40.9% of exposed morticians in the public

The equations to estimate lethal concentrations of FA in the mortuaries were developed from the Figs. 1 and 2, the values of the constants α and β were obtained as 210.71 and -131.3 respectively for public mortuaries; and 323.8 and

mortuaries have daily exposure index between 1.5 and 2.0 rated as very high; 36.36% have DEI between 1.0 and 1.5 rated as high; while, 22.7% have DEI between 0.5 and 1.0 rated as moderate. Similarly, computed daily exposure index showed that 23.53% of exposed morticians in private mortuaries have DEI between 1.5 and 2.0 rated as very high; 64.71% have DEI between 1.0 and 1.5 rated as high; while, 11.76% have DEI between 0.5-1.0 rated as moderate. Generally, 77.2% of workers in the public mortuaries have high daily formaldehyde exposure index, while 88.24% of the workers in the private mortuaries have high daily formaldehyde exposure index.

Analysis of variance indicates that the difference between the DEI in public and private mortuaries was not statistically significant ($p = 0.126$; 95%CI).

Normality test also showed that DPD distribution in the mortuaries does not follow a normal distribution (Fig. 4). The average formaldehyde concentrations, inhalation rate and the duration of exposure and the number of working hours per day were used to calculate the DPD. The results (Table 8) showed that daily potential dose in public mortuaries varies between 7.92 mg/d and 23.76 mg/d with a mean




Fig. 3. DEI normal Q-Q plots of normality test

and standard deviation of 17.82±6.2 mg/d. Also, daily potential dose in private mortuaries varies between 8.24 mg/d and 24.72 mg/d with a mean and standard deviation of 19.66±4.2 mg/d. Analysis of variance indicates that the difference between the DPD in public and private mortuaries was not statistically significant (p = 0.131; 95%CI). It is observed that daily potential dose increases with time of exposure (Fig. 5) Lower daily doses were obtained during the 4-hour exposure, while higher daily doses were obtained during the 12-hour exposure. This showed that the longer the exposure period the higher the dose and hence the more the effects on the exposed workers. These levels of daily dose exposures have been found to cause acute health effects [25]. Thus, morticians in Rivers State are in danger of adverse health effects due to formaldehyde exposure as also reported by [12-26].


The computed average daily doses (ADD) for both the public and private mortuaries were compared with [17] reference dose (RfD) of 0.2 mg/kg/d. The computed average daily doses for public mortuaries ranged from 0.09 to 0.4 mg/kg/d with a mean value of 0.25 mg/kg/d (Fig. 6), while the ADD values for private mortuaries ranged from 0.1 to 0.39 mg/kg/d with a mean value of 0.26 mg/kg/d (Fig. 6). The ADD values for public and private mortuaries exceeded the reference dose by 25% and 30% respectively. These results revealed short-term or acute non-cancerous health effects associated with formaldehyde exposure among the mortuary workers in both the public and private mortuaries in Rivers State. Computed hazard quotient for

both public and private mortuaries are 1.25 and 3.0 respectively (Table 8). These values are greater than 1 (> 1) indicating that there is a considerable or significant non-cancer related risk of formaldehyde exposure in the mortuaries.

Table 8. Computed daily potential dose

| Public mortuaries (DPD, mg/d) | Private mortuaries (DPD, mg/d) |
|-------------------------------|--------------------------------|
| 23.76 | 20.6 |
| 23.76 | 16.48 |
| 15.84 | 20.6 |
| 23.76 | 16.48 |
| 19.8 | 16.48 |
| 7.92 | 16.48 |
| 23.76 | 24.72 |
| 7.92 | 24.72 |
| 19.8 | 20.6 |
| 15.84 | 20.6 |
| 19.8 | 20.6 |
| 19.8 | 8.24 |
| 7.92 | 20.6 |
| 15.84 | 16.48 |
| 23.76 | 20.6 |
| 15.84 | 16.48 |
| 23.76 | 24.72 |
| 7.92 | 16.48 |
| 23.76 | 24.72 |
| 7.92 | 16.48 |
| 23.76 | 24.72 |
| 19.8 | 24.72 |

Computed LADD for public mortuaries ranged from 0.02 mg/kg/d to 0.11mg/kg/d with a mean value of 0.07 mg/kg/d (Fig. 8), while computed






Fig. 4. DPD normal Q-Q plots of normality test




Fig. 5. Computed daily potential dose (DPD)




Fig. 6. Mean average daily doses for public and private mortuaries

Table 9. Computed hazard quotient

| Mortuary category | Mean ADD (mg/kg/d) | HQ |
|--------------------|--------------------|------|
| Public mortuaries | 0.25 | 1.25 |
| Private mortuaries | 0.26 | 1.3 |




Fig. 7. Mean lifetime average daily doses for public and private morticians

Table 10. Computed cancer related risk

| Mortuary category | Mean LADD (mg/kg/d) | CRR | Safe threshold [22] |
|--------------------|---------------------|----------------------|---------------------|
| Public mortuaries | 0.07 | 1.5×10^{-3} | $10^{-4} - 10^{-6}$ |
| Private mortuaries | 0.09 | 1.9×10^{-3} | |

LADD for private mortuaries ranged from 0.04 mg/kg/d to 0.17mg/kg/d with a mean value of 0.09mg/kg/d (Fig. 7). The computed LADD values for both public and private mortuaries are within acceptable reference dose of 0.2 mg/kg/d for formaldehyde exposure [17]. The computed cancer related risk values for public and private mortuaries are 1.5×10^{-3} and 1.9×10^{-3} respectively (Table 10). These values exceeded the threshold target range of $10^{-4} - 10^{-6}$ for cancer risk management [22]. This implies that the mortuary workers/morticians may be at significant cancer risk due to formaldehyde exposure in their workplace environment. Thus these morticians could develop cancer related diseases such as nasal cavity, and nasopharynx, later in life after retirement from service. It has been reported in a study that formaldehyde exposure causes the impairment of the synthetic function of the liver of mortuary workers in Nigeria and also significantly reduced their total globulin level resulting in increased risk of suppressed humoral immunity [26]. However, another study found no association between formaldehyde and either

Hodgkin leukemia or chronic myeloid leukemia [27].

4. DISCUSSION

Previous studies had reported that chronic exposure to FA by male funeral directors revealed three times higher likelihood to die from Amyotrophic lateral sclerosis (ALS), i.e. Lou Gehrig's disease compared with FA unexposed population [11]. Lou Gehrig's disease is a central nervous system (motor neurons) that causes nervous damage and can lead to impairment in movement, eating, talking, breathing and eventual death. Similarly, our present study has also showed that health effects are work duration dependent. The computed cancer related risk for both public and private mortuaries are high and far exceeded the threshold target of 10^{-4} - 10^{-6} for cancer risk management [22]and thus poses a significant cancer risk to morticians with over 20 years of service.

Some studies that evaluated the effects of FA when chronically exposed with high

concentrations of FA have reported that it causes increased prevalence of headache, depression, mood changes, insomnia, irritability, attention deficit and memory loss [25]. FA has been classified as a human carcinogen [28]; its use has not been banned yet. Aside, the central nervous system (CNS) sequelae, it's been reported to have respiratory irritation effects that leads to chest pain, coughing and shortness of breath and asthma [15]. These findings corroborated earlier finding by Obed-Whyte et al. [13].

The results of health risk analysis from this research corroborate with previous case control study among funeral industry workers who had died between 1960 and 1986. That study related cancer risk to duration of employment, work practices and estimated FA exposure levels in the funeral industry and concluded that increased mortality/risk from myeloid leukemia was greatest among those who have worked as morticians for more than 20 years [26].

5. CONCLUSION

The study revealed that embalmers in both public and private mortuaries in Rivers State were exposed to high lethal concentrations and dose of formaldehyde use for the preservation of human cadavers. Results of both semi-quantitative and quantitative analysis indicated very high risk of FA exposure in both public and private mortuaries. The study showed a considerable non-cancer and cancer related health risks in the mortuaries due to the inhalation of formaldehyde gas. Analysis of short-term effect showed significant non-cancer health risk among the mortuary workers. Life-time risk analysis indicated significant carcinogenic health related risk among the mortuary workers. Thus cancer risks and non-cancer risks exist in both public and private mortuaries in the State. Therefore, occupational exposure to FA in mortuaries constitutes a significant health hazards in Rivers State, Nigeria.

6. RECOMMENDATION

Occupational and public health workers should create awareness among mortuaries operators/owners in Rivers State on the health risk faced by the morticians, particularly embalmers so that appropriate action can be taken to minimize exposure to FA. Regular monitoring of FA in all the mortuaries in the State

should be carried out by the National Environmental Standards and Regulations Enforcement Agency (NESREA). The Agency should also enact and enforce laws or guidelines on the use of FA in mortuaries in the State.

Management of mortuaries in the State should engage the services of qualified and registered assessors on Chemical Health Risk to conduct health surveillance on the exposed mortuary workers.

Further studies are therefore recommended to help increase the index of association and help clarify the content analysis of this study and also assess FA effect on the male fertility level of the exposed morticians

CONSENT AND ETHICAL APPROVAL

It is not approval.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar MS, Goud BR, Joseph B, Varghese. Occupational health and safety measures in a mortuary of Private Tertiary Health Care Medical College Hospital, Bangalore. *Indian Journal of Forensic Medicine and Toxicology*. 2016;10(1):98-103.
2. Okoth-Okello AM, Ogonda BA, Ogolla S, Tonui WK, Onyango R. Quality assurance for occupational health and safety administration (OSHA) In the morgue: The impact of SOP domestication on implementation and practice of universal safety precautions in Kenya. *Journal of Biology, Agriculture and Healthcare*. 2013; 3(19):1-10.
3. Work Safe. Guide for the funeral industry. Workplace Health and Safety Queensland. 2013, PN10564 Version 2; 2013. Available:www.worksafe.qld.gov.au
4. Work Safe. Funeral Directors Information and Checklist. Government of Western Australia, Department of Commerce; 2010. Available:https://www.commerce.wa.gov.au/sites/

5. WHO guidelines for indoor air quality: Selected pollutants. The WHO European Centre for Environment and Health, Bonn Office; 2010.
6. NCDOL. A Guide to Formaldehyde. N.C. Department of Labor, Occupational Safety and Health Program; 2013.
7. USEPA. Health and Environmental Effects Profile for Formaldehyde. EPA/600/x-85/362. Environmental Criteria and Assessment Office, Office of Health and Environmental Assessment, Office of Research and Development, Cincinnati, OH; 1988.
8. WHO. Environmental Health Criteria for Formaldehyde. Volume 89. World Health Organization, Geneva, Switzerland; 1989.
9. Hauptmann M, Lubin JH, Stewart PA, Hayes RB, Blair A. Mortality from Lymphohematopoietic malignancies among workers in FA industry. *Journal of the National Cancer Institute*. 2013;95(21): 1615-1623.
10. USDHHS. Registry of toxic effects of chemical substances (RTECS, online database). U.S. Department of Health and Human Services National Toxicology Information Program, National Library of Medicine, Bethesda, MD; 1993.
11. Oaklander M. Why funeral directors may be at higher risk for ALS; 2015. Available:<http://time.com/3956241/funeral-directors-als/>, accessed 25th March 2019
12. Douglas KE, Peterside SS. Assessment of workplace hazards in mortuaries in Port Harcourt. *Port Harcourt Med. J*. 2016; 10(3):102-110.
13. Obed-Whyte R, Douglas KE, Akaranta O. Comparative assessment of formaldehyde concentrations in public and private mortuaries in Rivers State, Nigeria. *Journal of Scientific Research & Reports*; 2019.
14. MOM. A Semi-Quantitative Method Assess occupational exposure to harmful chemicals. Ministry of manpower. Occupational Safety and Health Division; 2010. Available:<http://www.mom.gov.sg>
15. Tang TK. WSH risk management & OH development in Singapore. Conference proceedings, Ministry of Manpower, OSH Specialists Department, Singapore; 2016.
16. USEPA. Guidelines for Exposure Assessment. Risk Assessment Forum, U.S. Environmental Protection Agency Washington, DC. *Federal Register*. 1992; 57(104):22888-22938.
17. USEPA. Exposure factors handbook. National Center for Environmental Assessment, Office of Research and Development; 1997.
18. Beheshti MH, Firoozi CA, Alinaghi LAA, Rostami S. Semi-quantitative risk assessment of health exposure to hazardous chemical agents in a petrochemical plant. *JOHE*, Winter. 2015;4(1).
19. Heydari P, Varmazyar S, Ahmadi S, Alizadeh SS, Fallahi A. Evaluating the semi-quantitative risk of occupational exposure to chemicals in one of the petrochemical industries. *Scientific Journal of Review*. 2016;5(7):435-439.
20. Dazi H, Heydari P, Shokri S, Varmazyar S, Safari Variani A. Semi-quantitative assessment of the health risk of occupational exposure to chemicals and evaluation of spirometry indices on the staff of petrochemical industry. *Arch Hyg Sci*. 2017;6(1):49-57.
21. USEPA. Chapter 6—Inhalation Rates. Exposure Factors Handbook National Center for Environmental Assessment, Office of Research and Development; 2011.
22. USEPA. Risk assessment guidance for superfund, Human Health Evaluation Manual. EPA/540/189/ 002. US Environmental Protection Agency, Washington, DC. 1989;1.
23. National Bureau of Statistics. Statistical Report on Women and Men in Nigeria; 2018. Available:<https://nigerianstat.gov.ng/download/784>
24. OEHHA. Current proposition 65 no significant risk levels (NSRLs) Maximum Allowable Dose Levels (MADLs). Updated April 2018. California Environmental Protection Agency (CalEPA); 2018. Available:<https://oehha.ca.gov/proposition>
25. ATSDR. Addendum to the Toxicological Profile for Formaldehyde. Agency for Toxic Substances and Disease Registry, Division of Toxicology and Environmental Medicine; 2010.
26. Olooto WE. Assessment of the effect of formaldehyde Exposure on the liver in mortuary workers in South Western Nigeria. *Nigerian Medical Practitioner*. 2010;57(4):65-68.
27. Checkoway H, Dell LD, Boffetta P, Gallagher AE, Crawford L, Lees PSJ, Mundt KA. Formaldehyde exposure and mortality risks from acute myeloid

- mleukemia and other lymphohematopoietic malignancies in the US national cancer institute cohort study of workers in formaldehyde industries. American College of Occupational and Environmental Medicine. J. 2015;57(7):785-794.
28. IARC. A review of human carcinogens Part F: Chemical agents and related occupations. International Agency for Research on Cancer Monographs on the evaluation of carcinogenic risks to humans. Lyon: WHO. 2012;100.

© 2019 Roland et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48794>