



Screening for Hypertension and Diabetes in an Underserved Population through Community Outreach; A Case of Rural Community in Enugu State, Nigeria

C. Onodugo Obinna¹, C. Aniwada Elias^{2*} and P. Onodugo Nkeiru¹

¹*Department of Community Medicine, College of Medicine, University of Nigeria, Enugu, Nigeria.*

²*Department of Medicine, College of Medicine, University of Nigeria, Enugu, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors COO and CAE designed the study, authors PON and CAE wrote the protocol, author CAE managed the analyses of the study. Authors COO, CAE and PON managed the literature searches. Authors COO, CAE and PON wrote the first draft of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2019/v15i230116

Editor(s):

- (1) Dr. Nicolas Padilla-Raygoza, Department of Nursing and Obstetrics, Division of Health Sciences and Engineering, Campus Celaya Salvatierra, Mexico.
(2) Dr. Alexandre Sérgio Silva, Professor, Department of Physical Education, Federal University of Paraíba, Brasil.
(3) Dr. Giuseppe Murdaca, Professor, Clinical Immunology Unit, Department of Internal Medicine, University of Genoa, Italy.

Reviewers:

- (1) Ahmed Esmael, Mansoura University Hospital, Egypt.
(2) P. G. Kamal Mehta, Jagdish Chandra D.A.V College, India.
(3) K. D. Mini, Mahatma Gandhi University, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48743>

Original Research Article

Received 02 March 2019

Accepted 11 May 2019

Published 25 May 2019

ABSTRACT

Introduction: Hypertension and Diabetes are the commonest co-morbidity of each other and are among the principal cause of the burden of non-communicable diseases in developing countries. It is important to identify patients with these conditions early in the disease process. This study was to determine the prevalence of elevated Blood Pressure (BP) and elevated Fasting Blood Sugar (FBS) as well as relate it to the characteristics of the study participants in a rural community in Enugu State, Nigeria.

Methods: Community based cross-sectional study in form of outreach was done. The study was conducted over 1 week period among participants aged 18 years and above. Proforma was used

*Corresponding author: E-mail: eaniwada@gmail.com, eaniwada74@gmail.com;

in collecting information on characteristics of participants including age, sex and occupation. Measurements of BP, FBS and BMI were done. Chi square test and Binary Logistic Regression were used for analysis.

Results: Majority of participants were aged > 45 years 127(56.7%), and females 139(62.1%), Mean(SD) 46.89((21.84) Elevated BP 55(24.6%), elevated FBS 42(18.8%), both elevated BP and FBS. 13(5.8%). higher proportion of those aged > 45 years had elevated BP 51(92.7%) and elevated FBS 37(88.1%). More Females had elevated BP 35(63.6%) and elevated FBS 28(66.7%). Predictors were; age >45 years for elevated BP (AOR 18.4; 95% CI 5.7-59.5) and for have elevated FBS (AOR 8.9; 95% CI 3.0-26.5).

Conclusion: Prevalence of raised BP and FBS as well as co-morbid condition was high. It was more among females and older age. Age was a predictor of both raised BP and FBS. This calls for interventional programmes that will assist in limiting the increasing burden of the diseases in rural communities.

Keywords: Raised blood pressure; raised blood sugar; screening; outreach; rural community

1. INTRODUCTION

Non-communicable diseases (NCD), essentially cardiovascular diseases like Hypertension, Diabetes, Cancer and chronic respiratory diseases, are responsible for about 68% (38 million) of the 56 million deaths that occurred globally during 2012 [1]. Practically, 80% of these NCD deaths (29 million) occurred in low and middle-income countries (LMICs) [1]. In addition, the African region of the world is experiencing a double epidemic of both communicable and non-communicable diseases. It is reported that in Sub-Saharan Africa, the menace of NCDs could surpass that of communicable diseases in the nearest future [2].

Hypertension and diabetes are among the important diseases responsible the high burden of non-communicable diseases in developing countries [2]. Both have significantly contributed to medical morbidity and mortality. Hypertension is the commonest co-morbidity of diabetes and diabetes is the commonest co-morbidity of hypertension. They have common genetic predisposition, share risk factors, and have similar environmental influences as causative factors, and are interrelated. They bring about an enormous financial stress on individuals, families, communities and the health system of any country [3]. Worldwide, Diabetes is responsible for about USD 727 billion dollars in health expenditure and 12% of total expenditure on health of adults in 2017 [4]. Fuelling the crisis is the upsurge in unhealthy lifestyle such as tobacco and alcohol consumption, the reduction of physical activity and the changes in dietary intake due to westernization of our culture [2].

In 2012, report according to The World Health Statistics was that one in three adults worldwide has an elevated Blood Pressure (BP) and one in 10 adults have diabetes [5]. The raised BP is responsible for about 50 percent of all deaths from stroke and heart disease [5]. The percentage of deaths attributed to raised blood glucose in those aged 20–69 years in LMICs was 60.5% in men and 45.6% in women [2]. There was doubling in the prevalence of diabetes between 1980 and 2014. Approximately 425 million adults (20-79 years) were living with diabetes in 2017 and is projected that by 2045 this will rise to 629 million [4]. The International Diabetic Federation (IDF) statistics on diabetes reported that Nigeria has the largest population of people living with diabetes and impaired fasting glucose in Africa [6]. Hypertension is also the commonest cardiovascular disease reported in the country [7]. Findings from studies done in Nigeria documented that the prevalence of diabetes varies across different regions of the country with range from 2.2 - 9.8% [8-11]. In line with this, other similar studies reported varying prevalence rates of hypertension in various regions of the country [8,12,13].

To reduce the prevalence and consequences of hypertension and diabetes a complimentary mixture of population-wide and individual interventions is required. Allowing persons to present at health facilities is not yielding desired result as majority of persons do not know that they have these silent killers. Likewise, there might be an enormous burden of hypertension and diabetes mellitus among underserved rural dwellers in the country. It is important to identify patients with these conditions early in the disease process. Periodic outreaches will contribute to early detection of persons with such

conditions. Prompt and effective referral following such detection help reduce morbidity and complications associated with the diseases. This study was to determine the prevalence of elevated Blood Pressure (BP) and elevated Fasting Blood Sugar (FBS) among rural residents of a community in Enugu State Nigeria well as to find out the risk factors associated with these conditions.

2. METHODS

2.1 Study Area

This was at Abor, a rural community in Udi Local government of Enugu state, South East Nigeria. The community is about 30km to Enugu Metropolis They engage in farming, trading as well as civil/public services. There are no well established health facilities even though there is a primary health care facility.

2.2 Study Design and population

A Community based cross-sectional study in form of outreach was done. The study was conducted over 1 week period. All participants aged 18 years and above who gave informed consent were included in the study.

2.3 Sampling Technique and Sample Size Determination

A total study was done. All participants who meet the inclusion criteria were recruited consecutively as they present for screening at the outreach venue throughout the period of study. A total of 224 patients were studied.

2.4 Data Collection Tools and Method

1. Proforma was designed and used in collecting information on characteristics of participants including age, sex, occupation, recording BP, FBS and BMI
2. Measurements of BP, FBS and BMI were done using standard tools and observing standard procedure

Blood pressure: The BP was measured in the sitting position with an appropriate sized cuff encircling the left arm held at the level of the heart. This was measured using the OMRON Arm-type fully Automatic Digital Blood Pressure Monitor, Model BP - 103H. Raised BP was defined using Joint National Committee on Hypertension (JNC) 7 classification as systolic BP ≥ 140 mmHg and/or diastolic BP ≥ 90 mmHg.

Blood sugar: One microliter (1 μ L) of whole blood was collected and tested for blood glucose level using the Accu-chek active test strip and glucometer (Roche Diagnostics GmbH, Mannheim, Germany). Aseptic conditions were maintained throughout the procedure. Diabetes was defined as a fasting blood glucose > 110 mg/dl (6.1mmol/l).

Weight and height: the weight and height of the respondents were measured using a standardized Stadiometer. Weight was measured to the nearest 0.5 kg with the subject standing motionless on the calibrated scale without footwear. Height was measured with the subject standing in an erect position and head positioned so that the top of the external auditory meatus was level with the inferior margin of the bony orbit. The BMI of the subjects was calculated as weight in kilograms divided by height in meters squared.

2.5 Data Collection and Analysis

Patient information were recorded by trained health workers to ensure accuracy of data. Measurements of BP and FBS were done by qualified medical doctors. IBM Statistical Package for Social Sciences Version 21 was used for data entry, editing and analysis. Results were presented in tables. Mean, Standard deviation, proportion and percentages were used as summary measures where appropriate. Chi square test was used to establish associations between characteristics of participants with BP and FBS status. Binary Logistic Regression was done for variables significant variables (age). Level of confidence was at $p < 0.05$.

2.6 Action Taken

Participants found to have elevated BP and/or FBS during the screening were provided education on appropriate lifestyle and dietary modifications, such as salt and fatty reduction as well as need for improved physical activity where not adequate. They were also instructed and referred to tertiary health care facilities.

3. RESULTS

Table 1 shows that majority of participants were aged > 45 years 127(56.7%) with mean age of 46.89 SD of 21.84 were females 139(62.1%), unemployed/students 73(32.6%) followed by Civil/public servants 63(28.1%) and 99(44.2%) had BMI of 18.5-24.9 mg/m²

Table 1. Characteristics of participants

Variables	Frequency	Percent (%)
Age (Yrs)		
≤ 45	97	43.3
>45	127	56.7
Mean(SD)	46.89(21.84)	
Gender		
Female	139	62.1
Male	85	37.9
Occupation		
Civil/public servant	63	28.1
Trading	30	13.4
Farming	40	17.9
Skilled worker	18	8.0
Unemployed/student	73	32.6
BMI		
<18.5	66	29.5
18.5-24.9	99	44.2
25-29.9	57	25.4
≥30	2	.9

Table 2. Screening status of participants

Variables	Frequency	Percent (%)
Blood pressure (BP)		
Normal	169	75.4
Elevated	55	24.6
Fasting Blood Sugar (FBS)		
Normal	182	81.3
Elevated	42	18.8
Both BP and FBS		
No	211	94.2
Elevated	13	5.8

Table 3. Blood pressure and fasting blood sugar disaggregated by characteristics of participants

Variables	Blood pressure		Fasting Blood Sugar	
	Normal Freq(%)	Elevated Freq(%)	Normal Freq(%)	Elevated Freq(%)
Age (Yrs)				
≤ 45	93(55.0)	4(7.3)	92(50.5)	5(11.9)
>45	76(45.0)	51(92.7)	90(49.5)	37(88.1)
Gender				
Female	104(61.5)	35(63.6)	111(61.0)	28(66.7)
Male	65(38.5)	20(36.4)	71(39.0)	14(33.3)
Occupation				
Civil/public servant	41(24.3)	22(40.0)	44(24.2)	19(45.2)
Trading	26(15.4)	4(7.3)	24(13.2)	6(14.3)
Farming	26(15.4)	14(25.5)	33(18.1)	7(16.7)
Skilled worker	14(8.3)	4(7.3)	17(9.3)	1(2.4)
Unemployed/student	62(36.7)	11(20.0)	64(35.2)	9(21.4)
BMI				
<18.5	54(32.0)	12(21.8)	56(30.8)	10(23.8)
18.5-24.9	77(45.6)	22(40.0)	81(44.5)	18(42.9)
25-29.9	37(21.9)	20(36.4)	43(23.6)	14(33.3)
≥30	1(0.6)	1(1.8)	2(1.1)	0(0.0)

Table 4. Relationship of characteristics of participants with blood pressure and fasting blood sugar

Variables	Blood pressure		χ^2 (p value)	AOR (95% CI of AOR)
	Normal Freq(%)	Elevated Freq(%)		
Age (Yrs)				
≤ 45	93(95.9)	4(4.1)	38.547(<0.001)	1
>45	76(59.8)	51(40.2)		18.36(5.66-59.54)
Gender				
Female	104(74.8)	35(25.2)	0.078(0.781)	NA
Male	65(76.5)	20(23.5)		
Occupation				
Civil/public servant	41(65.1)	22(34.9)	11.648(0.020)	NA
Trading	26(86.7)	4(13.3)		
Farming	26(65.0)	14(35.0)		
Skilled worker	14(77.8)	4(22.2)		
Unemployed/student	62(84.9)	11(15.1)		
BMI				
<18.5	54(81.8)	12(18.2)	5.850(0.119)	NA
18.5-24.9	77(77.8)	22(22.2)		
25-29.9	37(64.9)	20(35.1)		
≥30	1(50.0)	1(50.0)		
	Fasting blood sugar			
	Normal	Elevated		
Age (Yrs)				
≤ 45	92(94.8)	5(5.2)	20.757(<0.001)	1
>45	90(70.9)	37(29.1)		8.92(3.00-26.52)
Gender				
Female	111(79.9)	28(20.1)	0.467(0.494)	NA
Male	71(83.5)	14(16.5)		
Occupation				
Civil/public servant	44(69.8)	19(30.2)	9.487(0.050)	NA
Trading	24(80.0)	6(20.0)		
Farming	33(82.5)	7(17.5)		
Skilled worker	17(94.4)	1(5.6)		
Unemployed/student	64(87.7)	9(12.3)		
BMI				
<18.5	56(84.8)	10(15.8)	FT(0.454)	NA
18.5-24.9	81(81.8)	18(18.2)		
25-29.9	43(75.4)	14(24.6)		
≥30	2(100.0)	0(0.0)		

Table 2 shows that 55(24.6%) of participants had elevated Blood Pressure, 42(18.8%) had elevated Fasting Blood Sugar while 13(5.8%) both have elevated BP and FBS.

Table 3 shows that higher proportion of those aged > 45 years had elevated Blood pressure 51(92.7%) and elevated FBS 37(88.1%). More Females had elevated Blood pressure 35(63.6%) and elevated FBS 28(66.7%). More Civil/public servants had elevated Blood pressure 22(40.0%) and elevated FBS 19(45.2%). Higher proportion of those that had BMI of 18.5-24.9 18.5-24.9 mg/m² had elevated Blood pressure 22(40.0%) and elevated FBS 18(42.9%).

Table 4 shows that there were statistically significant association of blood pressure with age ($\chi^2 = 38.547$, $p < 0.001$), Occupation ($\chi^2 = 11.648$, $p = 0.020$). However there were no statistically significant association of blood pressure with gender ($\chi^2 = 0.078$, $p = 0.781$) and BMI ($\chi^2 = 5.850$, $p = 0.119$). Also, there were statistically significant association of Fasting blood sugar with age ($\chi^2 = 20.757$, $p < 0.001$). However there were no statistically significant association of blood pressure with gender ($\chi^2 = 0.467$, $p = 0.494$), Occupation ($\chi^2 = 9.487$, $p = 0.050$) and BMI (FT, $p = 0.454$).

Those aged >45 years were about 18 times (AOR 18.4; 95% CI 5.7-59.5 likely to have

elevated BP than those aged ≤ 45 years. Also those aged >45 years were about 9 times (AOR 8.9; 95% CI 3.0-26.5 likely to have elevated BP than those aged ≤ 45 years.

4. DISCUSSION

Hypertension and Diabetes are the commonest co-morbidity of each other. They have common genetic predisposition, share risk factors, and have similar environmental influences as causative factors, and are interrelated [14]. Both elevated Blood Pressure and elevated Fasting Blood Sugar occur more with advancing age. Result from this study reported that majority of participants were aged > 45 years and females. This is expected as most rural areas are inhabited by retired workers and older persons due urban migration in the country for greener pasture. Majority being were females can be partly explained by the better health seeking behavior of females compared to males.

It was also noted that generally, 24.6% of participants had elevated Blood Pressure, 18.8% had elevated Fasting Blood Sugar and 5.8% had both elevated BP and FBS. The reported prevalence for elevated Blood Pressure was lower than the 42.0%, 44.5% and 46.4% reported respectively different studies in South Eastern Nigeria [15-17]. It is similar to reports from other studies [12,13]. The observed differences may be due differences in sampling technique and location of the study as most of those previous studies were in urban areas whose life style is different from rural communities. However, these finding is revealing as it shows that hypertension, DM and co morbid condition are of high prevalence in rural communities in Nigeria. This calls for interventional programmes including; persistent health education, enlightenment campaigns and community surveillance programmes to aid reduce this growing burden of the diseases in rural communities.

Based on Gender, 25.2% females and 23.5% males had elevated Blood Pressure, 20.1% females and 16.5% males had elevated Fasting Blood Sugar, 5.0% females and 7.1% males had both elevated BP and FBS. More Females equally had elevated Blood pressure 63.6% and elevated FBS 66.7%. The higher prevalence among females were also documented in previous similar studies [12,20-23]. However, there was discordance with many other previous reports. A study involving review of studies on hypertension over five decades reported a similar

prevalence in men and women with range in prevalence of 8% to 46.45% [13]. A Meta analysis of the prevalence of hypertension from population based studies in south western Nigeria reported a higher prevalence in men than women with prevalence ranging from 12.4% to 34.8% [12]. Another study documented prevalence of HTN of 22% (25.9% in males and 20% in females). Similarly other studies had similar findings of higher prevalence among males [18-21]. The findings from this study can partly be explained by fact that women are generally more likely than men to say they are unwell.

Findings also show that those aged >45 years were about 18 times likely to have elevated BP than those aged ≤ 45 years as well as about 9 times likely to have elevated BP than those aged ≤ 45 years. Also Higher proportion of those aged > 45 years had elevated Blood pressure 92.7% and elevated FBS 88.1%. A study done in Mali documented that OR increases with age from 2.06 (30–44 years) to 7.25 (60 and more) [22]. This is similar to finding in other studies in Africa [23-25]. In Ibadan South West Nigeria, a study revealed that hypertension was significantly associated with being in age groups 30-49 years (OR 2.258, 95% CI: 1.311 - 3.884), ≥ 50 years (OR 7.145, 95% CI: 3.644 - 14.011) [26]. In the United States, the estimated percentage of people having diagnosed or undiagnosed diabetes was increasing with age. In the age group of 20-44 years, about 3.7% people had diabetes; in the age group 45-64 years about 13.7%; and age group of ≥ 65 years about 26.9% had diabetes [27]. The study done in Bali showed that the prevalence of raised blood sugar and DM were more than twice higher in the elderly than in the younger age group [28]. A study done in China documented that Fasting and random blood glucose level rose by 0.15 mmol/L, while 2-hour post-prandial blood glucose level rose by 0.26 mmol/L per decade-increase in age [29]. Several reports have documented that age is the strongest risk factor for CVD like Hypertension. In the United States, CVD was the leading cause of death for persons 65 years of age and over in 2007. It was responsible for 28% of deaths in this age group [30].

These findings from current study could possibly be as a result of participants' occupation and residence. Almost all rural dwellers engage in minor farming even if is around their houses since a high proportion of participants were farmers and traders, trekking long distances to

the farm or the farm work itself constituted increased physical activity. Sedentary lifestyle as a key modifiable factor for both diseases may be rare in these rural communities. Age of participants may equally be a factor as these conditions are more with advancing age which is where majority of respondents belong to. The implication of this finding is that since most persons with these conditions do not know that they have the disease, there is the likelihood that they die suddenly with their relatives ascribing it to supernatural things. Then for others that may present at health facility, they are likely to come when complications of their condition has occurred.

Recently the numbers of elders are on the increase worldwide with sharp rise in the developing countries. This has impacted on the prevalence of metabolic diseases (impaired fasting glycemia, DM) and Cardio vascular Diseases including Hypertension. This may be as a result of their age, process of aging itself or remotely through several other age-related risk factors. Some of such documented factors include; central obesity, mitochondrial dysfunction, lipid metabolisms disorders, inflammation, β -cell dysfunction, insulin resistance and metabolic syndrome [28,31].

5. CONCLUSION

Prevalence of raised BP and FBS as well as co-morbid condition was high and in line with findings from other studies. It was more among females and older age. Age was a predictor of both raised BP and FBS. It shows that hypertension, DM and co morbid condition is emerging danger even in rural communities in Nigeria. This call for interventional programmes including mouth outreaches, persistent health education, enlightenment campaigns and community surveillance programmes to aid reduce this growing burden of the diseases in rural communities.

CONSENT

Informed consent was obtained from participants. They were ensured of voluntary participation and confidentiality of their information.

ETHICAL APPROVAL

The Health Research and Ethics committee of University of Nigeria Teaching Hospital, Enugu

gave ethical clearance. Permission was equally obtained from traditional rulers of constituents communities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization World health report: Global status report on noncommunicable disease; 2014. Available:<http://www.who.int/nmh/publications/ncd-status-report-2014/en/> [Accessed on 12/2/2019]
2. Kengne AP, Awah PK, Fezeu L, Mbanya JC. The burden of high blood pressure and related risk factors in urban sub-Saharan Africa: Evidences from Douala in Cameroon. *Afr Health Sci.* 2007;7(1):38–44.
3. Isara AR, Okundia PO. The burden of hypertension and diabetes mellitus in rural communities in southern Nigeria. *Pan African Medical Journal.* 2015;20:103.
4. International Diabetes Federation. Diabetes Fact Sheet; 2018. Available:www.idf.org/webdata/ [Accessed on 12/2/2019]
5. World Health Organization. World health statistics 2012 report. WHO, Geneva; 2012.
6. International Diabetes Federation. Diabetes fact sheet. Available:www.idf.org/webdata/docs/background_info_AFR.pdf. [Accessed on 12/2/2019]
7. Mukadas AO, Misbau U. Incidence and patterns of cardiovascular diseases in north western Nigeria. *Niger Med J.* 2009; 50(3):55-57.
8. Akinkugbe OO. Non communicable diseases in Nigeria: National survey (final report) on hypertension, coronary heart disease, diabetes mellitus, G6PD deficiency and anaemia. National Expert Committee on Non-Communicable Disease. Federal Ministry of Health and Social Services, Lagos.
9. Chinenye S, Ogbera AO. Socio-cultural aspects of diabetes mellitus in Nigeria. *J Soc Health Diabetes.* 2013;1(1):15-21.
10. Omorogiuwa A, Oaikheni GA, Okioya P, Akubueze D, Owobu E, Enahoro I, et al. Diabetes mellitus: Prevalence amongst

- University staff in southern Nigeria and attitude towards routine glycemc/ glucosuric checkup. *Int J Biomed & Hlth Sci.* 2010;6(1):25-29.
11. Nyenwe EA, Odia OJ, Ihekweba AE, Ojule A, Babatunde S. Type 2 diabetes in adult Nigerians: a study of its prevalence in Port Harcourt, Nigeria. *Diabetes Res Clin Pract.* 2003;62:177-185.
 12. Ekwunife OI, Aguwa CN. A meta analysis of prevalence rate of hypertension in Nigerian populations. *Journal of Public Health and Epidemiology.* 2011;3(13):604-607.
 13. Ogah OS, Okpechi I, Chukwuonye II, Akinyemi JO, Onwubere BJC, Falase AO, Stewart S, Sliwa K. Blood pressure, prevalence of hypertension and hypertension related complications in Nigerian Africans: A review. *World J Cardiol.* 2012; 4(12):327-340.
 14. Buchan J, Couper I, Tangcharoensathien V, et al. Early implementation of WHO recommendations for the retention of health workers in remote and rural areas. *Bull World Health Organ.* 2013;91(11): 834–840.
 15. Ulasi II, Ijoma CK, Onwubere BJ, Arodiwe E, Onodogo O, Okafor C. High prevalence and low awareness of hypertension in a market population in Enugu, Nigeria. *Int J Hypertens.* 2011;869675.
 16. Ahaneku GI, Osuji CU, Anisiuba BC, Ikeh VO, Oguejiofor OC, Ahaneku JE. Evaluation of blood pressure and indices of obesity in a typical rural community in eastern Nigeria. *Ann Afr Med.* 2011;10(2): 120-126.
 17. Onwubere BJ, Ejim EC, Okafor CI, Emehel A, Mbah AU, et al. Pattern of blood pressure indices among residents of a rural community in South East Nigeria. *Int J Hypertens.* 2011;621074.
 18. Sahraki R, Mirshekari M, Sahraki H, Mohammadi AR, Sahraki M, Khazaei Feizabad E. Hypertension among 30+ year-old people in Zahedan (Southeast of Iran). *Shiraz E Med J.* 2011;12(3):129–34.
 19. Sarry El-Din A, Erfan M, Kandeel W, Kamal S, El Banna R, Fouad W. Prevalence of pre-hypertension and hypertension in a sample of egyptian adults and its relation to obesity. *Aust J Basic Appl Sci.* 2012;6(13):481–9.
 20. Shapo L, Pomerleau J, McKee M. Epidemiology of hypertension and associated cardiovascular risk factors in a country in transition: A population based survey in Tirana City, Albania. *J Epidemiol Community Health.* 2003;57(9):734–9.
 21. Manandhar K, Koju R, Sinha NP, Humagain S. Prevalence and associated risk factors of hypertension among people aged 50 years and more in Banepa Municipality, Nepal. *Kathmandu Univ Med J (KUMJ).* 2012;10(39):35–8.
 22. Oumar Bâ, Youssouf Camara, Ichaka Menta, et al. Hypertension and associated factors in rural and urban areas Mali: Data from the step 2013 survey. *International Journal of Hypertension.* 2018;Article ID 6959165:7. Available:<https://doi.org/10.1155/2018/6959165>
 23. Awoke A, Awoke T, Alemu S, Megabiaw B. Prevalence and associated factors of hypertension among adults in Gondar, Northwest Ethiopia: A community based cross-sectional study. *BMC Cardiovascular Disorders.* 2012;12:113
 24. Abebe SM, Berhane Y, Worku A, Getachew A. Prevalence and associated factors of hypertension: A cross-sectional community based study in Northwest Ethiopia. *PLoS One.* 2015;10(4): e0125210.
 25. Gebreselassie KZ, Padyab M. Epidemiology of hypertension stages in two countries in sub-sahara africa: Factors associated with hypertension stages. *International Journal of Hypertension.* 2015;12. Available:<http://dx.doi.org/10.1155/2015/959256>
 26. Ajayi IO, Sowemimo IO, Akpa OM, Ossai NE. Prevalence of hypertension and associated factors among residents of Ibadan-North Local Government Area of Nigeria. *Nig J Cardiol.* 2016;13:67-75.
 27. Centers for Disease Control and Prevention. National diabetes fact sheet: National estimates and general information on diabetes and prediabetes in the United States. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2011.
 28. Suastika K, Dwipayana P, Saraswati IMR, Kuswardhani T, Astika N, Putrawan IB, Matsumoto K, Kajiwara N, Taniguchi H. Relationship between age and metabolic disorder in the population of Bali. *J Clin Gerontol Geriatrics.* 2011;30:1-6.
 29. Ko GT, Wai HP, Tang JS. Effects of age on plasma glucose levels in non-diabetic

- Hong Kong Chinese. *Croat Med J.* 2006; 47(5):709-13.
30. National Center for Health Statistics. Health, United States: With special feature on death and dying. Washington DC: US Government Printing Office; 2010. Available:<http://www.cdc.gov/nchs/data/health/hs10.pdf>. 2011
- [Accessed on 12/2/2019]
31. Suastika K, Dwipayana P, Semadi MS, Kuswardhani TA. Age is an important risk factor for type 2 diabetes mellitus and cardiovascular diseases; 2006. Available:<http://dx.doi.org/10.5772/52397> [Accessed on 12/2/2019]

© 2019 Obinna et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48743>