



Optimum K Fertilizer Level for Growth and Yield of Wheat (*Triticum aestivum*) in Cambisols of Northern Ethiopia

Hagos Brhane^{1*} and Kassa Teka²

¹*Mekelle Soil Research Center, Tigray Agricultural Research Institute, Mekelle, Ethiopia.*

²*Department of Land Resources Management and Environmental Protection, Mekelle University, Mekelle, Ethiopia.*

Authors' contributions

This work was carried out in collaboration between both authors. Author HB designed the study, wrote the protocol and the first draft of the manuscript. Author KT reviewed the experimental design all the drafts of the manuscript. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJA/2018/38742

Editor(s):

(1) Bilal Ahmad Wani, Sher-e-Kashmir University of Agricultural Sciences and Technology, Jammu and Kashmir, India.

Reviewers:

(1) Ade Onanuga, Canada.

(2) Gul Ebru Orhun, Canakkale Onsekiz Mart University, Bayramic Vocational College, Turkey.

(3) Hasan Kiliç, Bingol University, Turkey.

Complete Peer review History: <http://www.sciencedomain.org/review-history/23166>

Original Research Article

Received 24th October 2017
Accepted 11th January 2018
Published 13th February 2018

ABSTRACT

A field experiment was conducted in summer to evaluate the response of wheat to different potassium fertilizer rates on Cambisols of Tigray, Northern Ethiopia. The experiments were laid out in Randomized Complete Block Design replicated three times with 4 levels of potassium (0, 30, 60, 90, K₂O kg/ha). Data on yield and yield components of wheat were collected and regression, as well as variance analyses, were done. Results depicted that wheat plant height, spike length, harvest index and 1000 seed weight were all not significantly influenced by K fertilizer rates. However, biological and grain yield of wheat was significantly influenced by potassium levels. Hence, the highest biological yield (straw + grain) and grain yield of wheat were obtained at the rate of 90 K₂O kg/ha. Besides, the highest N and K uptakes by wheat were found at 60 K₂O kg/ha. Similarly, the highest apparent K recovery and agronomic use efficiency were found at 30 K₂O kg/ha. Hence, it

*Corresponding author: E-mail: hagosb7@gmail.com, hagosb@gmail.com;

can be concluded that potassium fertilization is important and the levels in the blended formula did not meet the wheat requirement in the soil reference group. Thus, this study recommends application of potassium.

Keywords: Potassium; blended fertilizer; cambisols; wheat; uptake.

1. INTRODUCTION

Wheat cultivation is a major farming practice in Ethiopia. However, soil degradation and nutrient depletion have gradually increased and become a serious threat to agricultural productivity in the country [1]. In line with this, low production of wheat has been shown in various parts of the country as a result of limited nutrient supply [2].

Increasing soil productivity is absolutely necessary to feed the increasing population in Ethiopia. In addressing this issue balanced fertilization with an optimum application rate is necessary to improve soil fertility and thus increasing the productivity of crops including wheat.

Among others, potassium fertilization has been improved growth and yield of wheat crops in various parts of the world. Several researchers conducted in Bangladesh, Saudi Arabia, Iran and India such as by [3-6] respectively indicated that growth and yield of wheat were increased by application of potassium fertilizer at different levels. On the other hand, fertilizer demonstrations are carried out in Ethiopia by the FAO and the then Ministry of Agriculture through the Freedom from Hunger Campaign conducted in the sixties and early seventies showed that the response to potash fertilization was inconsistent; thus, only urea and DAP were recommended for implementation (Tekalign Mamo, personal communication September 6, 2015). As a result potash fertilization was not practiced for the last many years due to the view that potassium was not deficient in Ethiopian soils. However, recent research findings such as by [7-10] have indicated that potassium was deficient in various areas of the country. In line with this, as one part of nutrient management strategy potassium fertilization has been started in 2014 in the form of blended fertilizers (between 7 and 12 kg per 100kg in the form of K_2O) in different regions of the country. Despite the various efforts made in including K as fertilizer through the introduction of K containing blended fertilizers in Ethiopia, the optimum level and its effect on growth, yield and

nutrient uptake of wheat on specific soil type in the various parts of the country was not studied yet. Besides, there is no adequate evidence which justifies whether the recommended rates of K in the blending formula (between 7 and 12 kg/ha in the form of K_2O) meets the crop demand or not. Thus, this study was designed to investigate potassium fertilization and its optimum level of growth, yield and nutrient uptake of wheat in Cambisol of northern Ethiopia.

2. MATERIALS AND METHODS

2.1 Study Area

The study was conducted in Enderta district, which is located in south eastern zone of Tigray region, northern Ethiopia. Before this experiment the field was covered by barley and tef respectively, for previous two consecutive growing seasons. The district is bounded by Hintalo Wajerat in the south, Seharti -Samre and Degua-Tembien in the west, Kilte-Awulaelo in the north and Afar region in the east. Geographically, the district is located between $13^{\circ}12'55''$ - $13^{\circ}38'38''$ N latitudes and $39^{\circ}16'43''$ - $39^{\circ}48'08''$ longitudes. The average elevation of the area is about 2200 m above sea level [11]. The Wereda falls in SM2-5b Agroecology, characterized by dry climatic conditions and erratic rainfall. Based on meteorological data collected from the nearest meteorological station on the study site, annual rainfall of the latest six years ranges between 258 and 756 mm. The growing season of 2015 had received a relatively lower rainfall compared to the long term average, since it was affected by El-Nino. The mean annual temperature ranges between 11.5 and 24.4°C. The most common soils of the study district are: Cambisol, Calcisols, Vertisols, Kastanozems, Leptosol, Luvisols, Phaozems, Regosols and Fluvisols [11].

2.2 Experimental Design and Procedures

The experiment had 4 levels of potassium (0, 30, 60 and 90 K_2O kg/ha) applied as potassium

chloride (KCl) on top of recommended blended fertilizers. These treatments were laid following Randomized Complete Block Design (RCBD) with three replications. The plot size was 3 m by 3 m with spacing of 1 m between blocks and 0.5 m between plots. On top of the blended fertilizer which contains 15.2% N, 48.8% nitrogen was added to satisfy N wheat requirements (64N kg/ha.) in the area. The blended fertilizer was applied at planting, while the nitrogen and K fertilizers were applied twice during the crop growth stage that is 1/3 of the full dose at planting and the other 2/3 at tillering stage.

The initial experimental field soils were analyzed for texture, pH, organic matter, cation exchange capacity (CEC), total nitrogen, available phosphorus and exchangeable K. The methods used for soil physical and chemical analysis were: Soil pH [12], Organic carbon % [13], soil texture by hydrometer [14], available Phosphorus [15], total nitrogen by Kjeldhal method [16], Neutral Ammonium acetate method [17] for cation exchange capacity and Exchangeable K⁺. After maturity, wheat crop samples were collected and partitioned into grain and straw parts. The grain and straw samples were analyzed for nitrogen and potassium. Plant total nitrogen was analyzed using Kjeldhal method [16] whereas potassium using dry ashing method [18]. In this experiment, picafloor (Kakaba) bread wheat variety was used as a test crop. Data on plant height, spike length, biological yield, grain yield and 1000 seed weight were collected.

The nutrient uptake by straw or grain was calculated by multiplying each nutrient concentration (%) by respective straw or grain yield in kg/ha. Moreover, apparent recovery and K agronomic use efficiency were calculated with the formulas proposed by [19].

$$\text{Apparent K recovery (kg/kg)} = \left(\frac{U_n - U_o}{n} \right) \quad (1)$$

Where; U_n stands for nutrient uptake at 'n' rate of fertilizer and U_o stands for nutrient uptake at control (no fertilizer) and 'n' stands for fertilizer applied.

$$\text{Agronomic K use efficiency (kg/kg)} = \left(\frac{G_n - G_o}{n} \right) \quad (2)$$

Where G_n and G_o stand for grain yield of fertilized plots at 'n' rates of fertilizer and grain

yield of unfertilized plots, respectively, and 'n' stands for nutrient applied.

2.3 Data Analysis

Analyses of variance (ANOVA) were carried out using Statistical Analysis Software (SAS) version 9. Whenever treatment effects were significant, mean separations were made using the least significant difference (LSD) test at the 5% level of probability. Moreover, regression analyses were also done.

3. RESULTS AND DISCUSSION

3.1 Soil Properties before Planting

The physical and chemical properties of the experimental fields before planting are indicated in Table 1.

Table 1. Soil physio- chemical properties of the site before sowing

Parameters	Value
pH _{water} (1:2.5)	7.55
Organic Carbon (%)	0.64
Total N (%)	0.06
P-Olsen (mg/kg)	2.88
Exchangeable K (Cmol/kg)	0.29
CEC (Cmol+/Kg)	23.6
% Sand	55
% Silt	25
% Clay	20
Textural class	Sandy Loam

The site is sandy loam in texture, slightly alkaline in soil pH, low in organic Carbon% and total nitrogen [20], medium in the CEC [17] and Exchangeable K [21] and low in available P [15]. The continuous cultivation without using an organic source of fertilizer may have contributed to the low level of organic carbon and total nitrogen.

3.2 Plant Height and Spike Length

Data presented in Fig. 1 and Table 2 showed that K fertilization had a promoting effect on plant height and spike length. Results showed that average plant height and spike length had increased with K application rates even though the trend was not consistent. The tallest plant

height and spike length were obtained in the treatment which received 60 K₂O kg/ha and it is not statistically different from the other treatments. However, the shortest plant height was measured at control treatments. These findings agreed with the research findings of [22-23].

3.3 Biological Yield (Total above Ground Biomass) and Grain Yield

In both regression analyses and analysis of variances showed that application of K fertilizer rates significantly increased the biological and grain yield of wheat in the study site. In line with this, the highest biological and grain yield of wheat was obtained from treatment that received 90 K₂O kg/ha. Moreover, the lowest biological and grain yields were recorded from the control

treatment and it was significantly lower ($P < 0.05$) as compared with other treatments. The differences in mean biological yield obtained from 0 and 30 K₂O kg/ha in one hand and 30 and 60 K₂O kg/ha on the other hand were not significant. Besides, the application of 60 K₂O kg/ha was significantly ($p \leq 0.05$) increased grain yield of wheat over application of 30 K₂O kg/ha on the study site. However, the difference in wheat grain yield between the application of 60 and 90 K₂O kg/ha was not significant ($p > 0.05$). In this study, treatments which received 90 K₂O kg/ha increased grain yield by 40.2% and 40.74% over treatments which received 30 and 0 K₂O kg/ha. The results are in agreement with the research findings of [24-25] who reported that application of potassium levels had significant effect on the biological and grain yield of wheat.




Fig. 1. Regression analyses of plant height and spike length




Fig. 2. Regression analyses of biological yield and grain yield

3.4 Harvest Index and 1000 Seed Weight

The result showed that harvest index and 1000 seed weight were not significantly affected by the applications of K rates. However, the highest harvest index and 1000 seed weight were recorded at a rate of 60 K₂O kg/ha. The non-significant harvest index result was indicating approximately equal positive effects of potassium on seed and biological yield. The non-significant result found on the 1000 seed weight was agreed with the research findings of [26] who reported that application of K at any level had no significant effect on 1000 seed weight of wheat.

3.5 Nutrient Uptakes, Apparent Recoveries and Agronomic Use Efficiency

3.5.1 K uptake by wheat grain and straw

The result indicated that K uptake by grain and straw were influenced by different K treatment combinations. K uptake by grain and straw of wheat had shown a linear increasing trend with

increasing K rates though there were some inconsistent results. The highest K uptake by grain and straw was found at 60 K₂O kg/ha. The lowest grain and straw K uptake were obtained from the control treatment in the studied soil reference group. This result is in agreement with the research findings of [27] in Kashmir, India.

3.5.2 N uptake by wheat grain and straw

Potassium fertilization had promoted N uptake by grain and straw. Nitrogen uptake by grain and straw of wheat was influenced by various K treatment combinations as indicated in Fig. 4. Grain and straw uptake of nitrogen by wheat increased linearly up to 60 and the lowest N uptake by grain and straw was found in control. This result is consistent with the research findings of [28].

3.5.3 Apparent recovery and agronomic use efficiency of K

The potassium application rate had influenced apparent K recovery and agronomic K use

Table 2. Effect of potassium fertilizer rates on wheat plant height, spike length, biological yield, grain yield, harvest index and 1000 seed weight

Treatment	Plant height (cm)	Spike length (cm)	Biological yield (kg/ha)	Grain yield (kg/ha)	Harvest index	1000 seed weight (g)
Control	47.1	6.1	3048.1 ^c	935.3 ^c	0.31	26.5
RBF	63.0	6.9	4772.2 ^b	1574.7 ^b	0.35	26.1
RBF+ 30 K ₂ okg/ha	66.9	7.1	6044.4 ^{ba}	1580.7 ^b	0.26	22.9
RBF+60 K ₂ okg/ha	67.1	7.2	5981.5 ^{ba}	2203.6 ^a	0.37	28.6
RBF+90 K ₂ okg/ha	66.4	7.2	6531.5 ^a	2216.2 ^a	0.34	27
Lsd(0.05)	ns	ns	1507	551.85	ns	ns
CV	12.2	6.4	15.2	17.2	15.3	14.2

Means followed by the same letter along columns are not significantly different. RBF: recommended blended fertilizer (NPKSZN), Lsd: least significant difference CV: Coefficient of variance, ns: non-significant




Fig. 3. Effect of K rates on K uptake by wheat grain and straw




Fig. 4. Effect of K rates on N uptake by wheat grain and straw

efficiency in the study site. Both apparent recovery and K use efficiency had shown decreasing trend with increasing K rates even though the apparent recovery was not consistent. As a result, the highest agronomic K use efficiency and apparent recovery were obtained at the lowest application rate (30 K₂O kg/ha).

Table 3. Effect of potassium level on apparent recovery and agronomic use efficiency

Level of K (kg/ha)	ARK (kg/kg)	AUE K (kg/kg)
30	1.88	21.51
60	0.41	21.14
90	0.61	14.23

ARK=Apparent recovery of potassium;
AUEK= Agronomic use efficiency of potassium

4. CONCLUSIONS

Potassium fertilization has been started in the country in the form of blended a blanket recommendation in 2014 to improve the productivity of cereals including wheat. However, there were no comprehensive works on its rates and whether the level in the blended formula meets the growth and yield requirement of the crop or not. Thus, the result of this experiment indicated that nutrient uptake, K recovery, K agronomic use efficiency, biological and grain yield of wheat were significantly responded from the additional K levels. Hence, the level of potassium in the blended formula did not meet the growth and yield requirement of wheat on Cambisols of the studied district. In line with this, the highest biological and grain yields of wheat were obtained at 90 K₂O kg/ha. However, the

highest K and N uptake by grain and straw were found at 60 K₂O kg/ha rates. Besides, the highest apparent K recovery and agronomic use efficiency were found at 30 K₂O kg/ha. Therefore, straight application of potassium is recommended rather than incorporating in the blend.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kebede F, Yamoah C. Soil fertility status and numass fertilizer recommendation of typic hapluusterts in the northern highlands of Ethiopia. World Applied Sciences Journal. 2009;6(11):1473-1480.
2. Gebreselassie Y. Selected chemical and physical characteristics of soils of Adet research center and its testing sites in North-western Ethiopia. Ethiopian Journal of Natural Resources; 2002.
3. Saha PK, Hossain ATMS, Miah MAM. Effect of potassium application on wheat (*Triticum aestivum* L.) in old Himalayan piedmont plain. Bangladesh Journal of Agricultural Research. 2010;35(2):207-216.
4. Alderfasi AA, Refay YA. Integrated use of potassium fertilizer and water schedules on growth and yield of two wheat genotypes under arid environment in Saudi Arabia 1-Effect on growth characters. Agric. Environ. Sci. 2010;9(3): 239-247.

5. Malek-Mohammadi M, Maleki A, Siaddat SA, Beigzade M. The effect of zinc and potassium on the quality yield of wheat under drought stress conditions. *International Journal of Agriculture and Crop Sciences*. 2013;6(16):1164.
6. Khan AA. Inamullah and MT Jan. 2014. Impact of various nitrogen and potassium levels and application methods on grain yield and yield attributes of wheat. *Sarhad J. Agric*. 2014;30(1):35-46.
7. Abegaz A. Indigenous soil nutrient supply and effects of fertilizer application on yield, N, P and K uptake, recovery and use efficiency of barley in three soils of Teghane, the Northern Highlands of Ethiopia. *African Journal of Agricultural Research*. 2008;3(10):688-699.
8. Deressa A, Bote B, Legesse H. Evaluation of soil cations in agricultural soils of East Wollega Zone in South Western Ethiopia. *Science, Technology and Arts Research Journal*. 2013;2(1):10.
9. EthioSIS. Report on Soil Fertility Status and Fertilizer Recommendation Atlas for Tigray Regional State, Ministry of Agriculture and Agricultural Transformation Agency (ATA), Addis Ababa, Ethiopia. 2014;76.
10. Wassie H. On -farm verification of potassium fertilizer Effect on the yielded of irish potato grown on Acidic soils of Hagere Selam, Southern Ethiopia. *Ethiopian Journal of Natural Resources*. 2009;11(2):207–221.
11. Gebre T, Kibru T, Tesfaye S, Taye G. Analysis of watershed attributes for water resources management using GIS: The case of chelekot micro-watershed, Tigray, Ethiopia. *Journal of Geographic Information System*. 2015;7(02):177.
12. Rhoades JD. In *methods of soil analysis, Part 2*. Second edition (A. L. Page, Miller and D. R. Keeney, Eds). American Society of Agronomy, Madison, U; 1982.
13. Walkley A, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*. 1934;37(1): 29-38.
14. Bouyoucos GJ. Hydrometer method improved for making particle size analyses of soils. *Agronomy Journal*. 1962;54(5): 464-465.
15. Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. United States Department of Agriculture; Washington; 1954.
16. Bremner JM, Mulvaney CS. Nitrogen—total. *Methods of soil analysis. Part 2. Chemical and microbiological properties. (Methodsofsoilan2)*. 1982;595-624.
17. Landon JR. *Booker tropical soil manual: A Handbook for soil survey and agricultural land evaluation in the tropics and subtropics*. Longman Scientific and Technical, Essex, New York. 1991;474.
18. Chapman HD. Cation-exchange capacity. *Methods of soil analysis. Part 2. Chemical and microbiological properties. (Methodsofsoilanb)*. 1965;891-901.
19. Fageria NK, Baligar VC. Methodology for evaluation of lowland rice genotypes for nitrogen use efficiency. *Journal of Plant Nutrition*. 2003;26(6):1315-1333.
20. Tadesse T, Haque I, Aduayi EA. *Soil, plant, water, fertilizer, animal manure and compost analysis manual*. ILCA/PSD Working Document (ILCA); 1991.
21. Jones Jr. JB. *Agronomic handbook: management of crops, soils and their fertility*. CRC Press; 2002.
22. Khan R, Gurmani AR, Gurmani AH, Zia MS. Effect of potassium application on crop yields under wheat rice system. *Sarhad Journal of Agriculture*. 2007;23(2): 277.
23. Tahir M, Tanveer A, Ali A, Ashraf M, Wasaya A. Growth and yield response of two wheat (*Triticum aestivum* L.) varieties to different potassium levels. *Pak. J. Life Soc. Sci*. 2008;6(2):92-95.
24. Wassie H, Tekalign M. The effect of potassium on the yields of potato and wheat grown on the acidic soils of chench and hagere selam in Southern Ethiopia. In: *International Potash Institute, e-ifc No. 35*. 2013;3-8.
25. Maurya P, Kumar V, Maurya KK, Kumawat N, Kumar R, Yadav MP. Effect of potassium application on growth and yield of wheat varieties. *The Bioscan*. 2014;9(4): 1371-1373.
26. Morshedi A, Farahbakhsh H. Effects of potassium and zinc on grain protein contents and yield of two wheat genotypes under soil and water salinity and alkalinity stresses. *Plant Ecophysiol*. 2010;2:67-72.

27. Wani JA, Malik MA, Dar MA, Farida A, Raina SK. Impact of Method of application and concentration of potassium on yield of wheat. *Journal of Environmental Biology*. 2014;35:623-626.
28. Ashok Singh J, Kumar P, Chauhan N, Singh GR. Effect of the levels of potassium and manganese on the uptake of N, P, and K and yield of wheat. *Journal of Agricultural Physics*. 2009;9:28-32.

© 2018 *Brhane and Teka*; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/23166>