



Pulmonary Function Impairment in Female Workers Exposed to Environments with Varied Ambient Air Pollution in the Central Business Area of Lusaka-Zambia

Lumba Siachingili^{1*} and Fastone M. Goma²

¹Cavendish University, Zambia School of Medicine, Lusaka, Zambia.

²Department of Physiological Sciences, University of Zambia, School of Medicine, Lusaka, Zambia.

Authors' contributions

This work was carried out in collaboration between both authors. Author LS designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author FMG managed and supervised the study. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BJMMR/2016/20719

Editor(s):

(1) Gaetano Santulli, College of Physicians & Surgeons, Columbia University Medical Center, New York, NY, USA.

Reviewers:

(1) Deepak Sharma, Chaudhary Charan Singh University, Meerut, India.

(2) Eugenio Ragazzi, University of Padova, Italy.

Complete Peer review History: <http://sciencedomain.org/review-history/11874>

Original Research Article

Received 6th August 2015
Accepted 1st October 2015
Published 17th October 2015

ABSTRACT

Aims: To determine the lung function of cleaners exposed to particulate matter of aerodynamic diameter less than 2.5 micrometer (PM_{2.5}) in the streets and offices in Lusaka, Zambia.

Study Design: This was a cross sectional study between two groups.

Place and Duration of Study: Lusaka city, central business area, between June and August 2014.

Methodology: The study included women between 18-50 years of age who had been working as street or office cleaners for 6 months or more. Males and individuals in both groups who used to smoke or were currently smokers, as well as those with a history of respiratory related illnesses or had cardiopulmonary conditions were excluded from the study. The cleaners were interviewed to get information on socio-demographic characteristics and other information using a structured interview schedule. The participants' lung volumes, forced expiratory volume in one second (FEV₁),

*Corresponding author: E-mail: lumbablue@gmail.com;

forced vital capacity (FVC) and their ratio (FEV_1/FVC) were measured using a MRI spirometry G spirometer. On the day of the interview, $PM_{2.5}$ in their work environment was sampled using a personal aerosol monitor (SIDEPAK AM510).

Results: Out of the 90 participants, 45 were street sweepers and 45 were office cleaners. More street sweepers had impaired lung function (FEV_1/FVC) 15(75%) than office cleaners 5(25%) $p=0.01$. FEV_1 was also significantly different among street sweepers 12(70.6%) and office cleaners 5(29.4%) $p=0.05$. $PM_{2.5}$ measurements revealed significantly high levels of exposure among street sweepers ($p=0.001$). Participants with impaired lung function ($p=.005$) and those with reduced FEV_1 percent predicted were exposed to significantly high concentrations of $PM_{2.5}$ ($p=0.012$).

Conclusion: Exposure to high $PM_{2.5}$ concentration is associated with pulmonary function impairment and reduced FEV_1 % predicted among cleaners.

Keywords: Ambient pollution; Fine Particulate matter ($PM_{2.5}$); Forced Expiratory Volume in 1 second (FEV_1); Forced Vital Capacity (FVC); lung function status.

DEFINITIONS

Ambient Air: refers to the air in the surrounding environment.

Particulate Pollution: Tiny solid and liquid droplets suspended in the air that when inhaled can cause damage to the lungs.

Fine Particulate matter: is a complex mixture of extremely small particles and liquid droplets. Fine particulate matter is 2.5 micrometres in diameter.

Spirometry: The measurement of how quickly air can be expelled from the Lungs.

Forced Expiratory Volume in 1 Second: The volume of air that can be forcibly exhaled from the lungs in the first second of forced expiration.

Forced Vital Capacity: The total volume of air that can be forcibly exhaled after taking the deepest breath possible.

Lung Function status: refers to how well air flows in and out of the lungs or an FEV_1/FVC ratio of greater than 70%, FEV_1 % predicted greater than 80% in women, or FVC% predicted greater than 80%.

Impaired Lung Function Status: when an individual has an FEV_1/FVC ratio of less than 70% or the loss or distortion or weakening of lung tissue leading to difficulty in air flowing out of the lungs.

Personal Protective Equipment: This is the protective clothing, facemasks, respirators, goggles, or other garment designed to protect the wearer's body from injury by blunt impacts,

electrical hazards, heat, chemicals, and infection, for job-related occupational safety and health purposes.

Exposure: The act of subjecting or an instance of being subjected to an action or an influence, (fine particulate air Pollution).

1. INTRODUCTION

Air pollution is considered a hazard to human health [1]. In the past decades, studies have highlighted the role of ambient air pollution as an important cause of both mortality and morbidity for many different cardiopulmonary diseases [2]. Ambient pollutants include suspended or respirable particulate matter (PM), nitrogen dioxide (NO_2), carbon monoxide (CO), ozone (O_3), and sulphur dioxide (SO_2) [3]. Among these ambient pollutants, respirable particulate matter (PM) is said to have the greatest effect on human health [4]. Airborne PM consists of a mixture of liquid and solid air-suspended particles, which are released straight into the atmosphere or after the transformation of gas into particles from natural or human-induced processes [5]. Some of the important sources of fine particulate matter include burning fuels emitted from vehicles, open air burning of house hold wastes and biomass cooking fuels such as charcoal and fire wood [6]. Studies conducted in many developing countries have reported an increase in $PM_{2.5}$ burden and its constituents [7,8]. In Africa, the increase in the burden of $PM_{2.5}$ is due to the growing ownership of motor vehicles, unpaved roads as well as continued use of biomass (firewood and charcoal) as a major domestic energy source [9]. Indeed combustion of biomass fuels is usually incomplete and is said to release several pollutants among them Particulate Matter.

The most important parameter for defining the toxicity of PM is particle diameter and composition [10]. The United States Environmental Protection Agency (US EPA) and other agencies that regulate air pollution have three main categories for PM: PM_{0.1} (ultrafine particles) PM_{2.5} (fine particulate matter) and PM₁₀ (coarse particulate matter), which refer to particles with aerodynamic diameter smaller than 0.1, 2.5 and 10 micrometres (µm), respectively [11]. Studies show that it is the fine (PM_{2.5}) and ultrafine (PM_{0.1}) fraction that are capable of penetrating deep into lung tissue and induce oxidative stress that are more harmful [12]. Furthermore, studies in electron microscopy show that most of the effectively retained particles in the lung parenchyma are PM_{2.5} [13]. Therefore, particle size and the ability to penetrate into the lung tissue and subsequent retention of the fine particles play an important role in causing lung function impairment [14].

When inhaled, air pollutants cause obstructive, restrictive or both types of functional impairment of the respiratory system manifested by reduced functional vital capacity (FVC), forced expiratory volume in one second (FEV₁) and their ratio FEV₁/ FVC [15]. PM_{2.5} induces cell injury and death of respiratory epithelial cells; it also leads to decreases in immunity defences through the destruction of macrophages [16]. Damage to lung vascular tissue creates a need for angiogenesis; In PM_{2.5} related pathology, angiogenesis is said to be induced by the ability of PM_{2.5} to produce Reactive Oxygen Species (ROS) that have been implicated in the activation of pro-inflammatory mediators, especially cytokines such as vascular endothelial growth factor, transforming growth factors (TGF) and so on [17,18]. These cytokines and the resulting inflammation stimulate angiogenesis. Angiogenesis, in lung vascular homeostasis, maintains an ideal number of capillaries per unit of lung volume, thus, it is an important mechanism in reestablishing a functional gas-exchange interface [19]. However, the profuse repair that results from excessive production of angiogenic stimulators, can result in lung remodeling that may progress to chronic fibrotic lung disease and persistent respiratory insufficiency or impaired Lung Function [20].

Inhalation of PM_{2.5} also increases airway reactivity and induces allergic symptoms [21]. Presence of allergies has been associated with impaired lung function status among susceptible occupational groups such as street sweepers, steel plant workers and so on [22,23].

Spirometry is an important as well as simple tool, in assessing the functioning ability of the lungs [24]. Spirometric measures of lung function, namely maximum forced vital capacity (FVC) and maximum forced expiratory volume in 1 s (FEV₁) have been described as early indicators of chronic respiratory and systemic inflammation [25]. The lowering of both the FEV₁ (FEV₁< 75% predicted for age and height) and FVC (FVC<80% predicted for age and height) indicates a restrictive lung impairment while the ratio thereof maybe greater than 70% [26,27]. In obstructive impairment, the FVC may be normal but FEV₁ is reduced [27].

1.2 Occupational Exposure to Fine Particulate Matter (Pm_{2.5}) Pollution

Occupation plays an important role on the level of personal exposure to pollutants as demonstrated in a study that showed that female street sweepers exposed to high concentrations of dust had lower lung function values compared to females of the same category working in an office [28]. The study further revealed that use of personal protective equipment (PPE) was essential in preventing this. Other studies have equally revealed that street sweepers by virtue of their exposure to dust were more likely to have a FEV₁/FVC ratio less than 60% [16,29]. Office cleaners are also at risk of developing lung function impairment due to exposure to indoor sources of PM_{2.5} like chemical detergents and fungal spores from furniture [30].

Despite belonging to an organised workforce, cleaners in Zambia like many other African countries are not sufficiently taken care of in terms of periodical health check-ups and provision of personal protective equipment such as gloves, facemasks and respirators. Sometimes, due to poor sensitization on the need to use this equipment others feel there is no need to use it, whilst the protective equipment maybe worn out and/or not replaced in good time [31].

Air pollution is a hazard to lung function but this has not been documented in cleaners in Lusaka. There is no air quality monitoring hence the levels of PM_{2.5} in ambient Lusaka air are not known and health-based limits for dust control in the various work places are lacking. Although studies on lung impairment have been carried out on specific occupational groups in Zambia such as miners and stone crushers [32,33], no study has been carried out on other occupational groups including sweepers.

This study was therefore, aimed at determining the level of air pollution particularly $PM_{2.5}$ in the work environments and the possible effects of this pollutant on the lung function of individuals that are exposed to these pollutants such as street sweepers and office cleaners. The data obtained would be useful as an advocate tool for provision of protective equipment for the cleaners. It will also provide insights on the possible effects of $PM_{2.5}$ on lung function to policy makers, health care providers and researchers and provide a baseline for further study. We envisage that this study will help improve enforcement and implementation of air quality regulations around the city.

2. MATERIALS AND METHODS

The study was conducted in the central business area of Lusaka Zambia. The study sites included all the 12 streets of the central business area. The indoor study sites along these streets were purposively sampled and included offices that had as little outdoor air circulation as possible.

2.1 Selection of Participants

The study population included female cleaners working within the central business area and these were divided into two groups according to their job category, 45 office cleaners and 45 street sweepers. Females aged between 18 and 50 years of age who had been working as street or office cleaners for 6 months or more were invited to participate in the study. Individuals in both groups who used to smoke or were currently smokers, as well as those with a history of respiratory related illnesses or had cardiopulmonary conditions were excluded from the study. Males were also excluded from the study as they constitute a very small proportion of the people in this sector. The participants in both groups were identified using a list of employees provided by the supervisor. The lists were used as a sampling frame for which random sampling technique was employed to select the participants. Informed consent was obtained from those selected who agreed to participate in the study. The study protocol was cleared by the University of Zambia Biomedical Ethics Committee (UNZABREC).

2.2 Data Collection

Data collection was carried out between the months of June and August 2014. A structured

interview schedule, with questions adapted from the American Thoracic Society (ATS) respiratory questionnaire, was used to collect demographic information and to record Spirometry data from participants. Prior to its use, the interview schedule was pilot tested on fifteen (15) randomly selected female cleaners to ascertain the levels of understanding. The questions were administered in commonly spoken language. The language used was simple and the cleaners had no difficulty understanding the questions. Information pertaining to the use of protective wear, cooking fuel, smoking history, occupation history, allergies and history of respiratory diseases were captured using the interview schedule.

2.3 Measurement of Lung Function

The lung function tests were carried out, using a portable MRI spirometry G spirometer (Medical Research International, Spirometry G, Rome, Italy). The device allowed for calibration and it had the software to predict lung function indices for age, sex and height. The tests were taken with participants in the sitting position by a trained spirometry technician. The procedure was explained to the participants who were urged to seal their lips tightly around the mouthpiece, breathe in fully (maximal inspiration) at the start of the test, immediately blast air out as fast and as far as possible until the lungs were completely empty.

Three maneuvers were done at 5-minute intervals and the best of the three results was recorded. The predicted FEV_1 , FVC were determined using height and age of the participants. Lung function status of each participant was determined using the FEV_1/FVC ratio. The lung function measures were stored on the device and also recorded.

2.4 Measurement of Fine Particulate Matter ($PM_{2.5}$)

A TSI SidePak AM510 Personal Aerosol Monitor (TSI incorporation St. Paul, MN United States of America) was used to sample and record the levels of fine particulate matter ($PM_{2.5}$) in the air. The aerosol monitor was attached to the participant and the sampling tube placed near the participants' breathing zone. The built in impactors of the aerosol monitor were set on the 2.5 cut off in order to sample only $PM_{2.5}$ concentrations in mg/m^3 then converted to

$\mu\text{g}/\text{m}^3$. The SidePak was zero-calibrated prior to each use by attaching a zero filter according to the instructions provided in the user guide. Measurements of $\text{PM}_{2.5}$ for both indoor and outdoor areas were taken in the morning, midday, and in the afternoon during cleaning for 30-60 minutes. $\text{PM}_{2.5}$ readings were stored in the sampling device and manually transferred to a data sheet for analysis.

2.5 Data Analysis

The Independent Samples Mann-Whitney U test for non-parametric data was used to compare medians of $\text{PM}_{2.5}$ across the indoor and outdoor cleaning sites. Explorative statistics using the Independent Samples Median Test were used to determine the association of $\text{PM}_{2.5}$ with lung function characteristics (predicted $\text{FEV}_1\%$ and $\text{FVC}\%$ and the ratio FEV_1/FVC).

Chi-square was used to determine the association between the dependent variable (lung function status) and the independent variables (age, cooking fuel, presence of allergies and use of PPE). It was also used to determine association of pulmonary function in the two groups of cleaners. Statistics were done at the 5% level of significance. Data analysis was done using IBM SPSS Statistics for Windows Version 20.0 (IBM Corp. Armonk, NY, USA).

3. RESULTS

The study consisted of 90 participants divided into two groups; 45 street sweepers and 45 office cleaners, all female, none were tobacco smokers or had previously smoked tobacco, all working within the central business area of Lusaka.

3.1 Lung Function Status of Participants

Fig. 1 is a description of lung function status of the two groups of cleaners. Median of FEV_1/FVC was 84%, minimum 64% and maximum was 100% for office cleaners while for street sweepers the median was 88%, minimum 43% and maximum was 100%. The medians of FEV_1/FVC were significantly different between the two groups of cleaners $p < 0.05$.

Table 1 describes the lung function characteristics of the participants. 15 (75% within lung function) street sweepers had impaired lung function compared to 5 (25%) office cleaners. This difference was statistically significant at $p < 0.01$.

3.2 Contributing Factors to $\text{PM}_{2.5}$ Exposure

Table 2 shows factors that could possibly contribute to participants' exposure to $\text{PM}_{2.5}$ such as use of PPE, cooking fuel, previous occupation and allergy symptoms and their association with lung function status.

Based on these characteristics, 58 (64.4%) of the 90 participants reported using PPE sometimes or not at all. Among those with impaired lung function 12 (60%) reported not using PPE consistently while 8 (40%) reported always using PPE. There was no significant difference in lung function between those always using PPE and those that used PPE occasionally or never ($p = .792$).

There was no significant difference in lung function status between participants who used charcoal as cooking fuel and those who used electricity ($p = .355$). However, among those that reported using charcoal as a cooking fuel 14 (70%) had impaired lung function status while among those that reported using electricity as cooking fuel, only 6 (30%) had impaired lung function status. Most (70%) of the participants with impaired lung function status used charcoal as cooking fuel.

About half (53%) of the participants reported that they suffered from allergic symptoms while at work. However, most of those with impaired lung function status (75%) reported having allergy symptoms. Table 2, shows lung function status and allergy symptoms among these participants. Among the participants with normal lung function status, 33 (47.1%) had allergies while among those with impaired lung function status 15 (75%) had allergies. There was a statistically significant association between presence of allergy symptoms and pulmonary function impairment ($p < 0.05$).

The data in Table 2 on the two groups of cleaners, reveals that, 29 (41.4%) of the participants with normal lung function were street sweepers, and 41 (58.6%) were office cleaners. 80% of those with impaired lung function, were street sweepers and 20% were office cleaners. There was a statistically significant association between cleaning location and pulmonary function $p = 0.002$.




Fig. 1. Medians of FEV₁/FVC between the cleaners

3.3 PM_{2.5} concentrations in the study Areas

The highest observed value for PM_{2.5} outdoors, was 398 µg/m³ whilst indoors it was 53 µg/m³. The lowest observed value outdoors was 123 µg/m³; while indoors it was 10 µg/m³. These values varied between morning and afternoon cleaning times the average of the two readings was recorded (Fig. 2). The PM_{2.5} concentration between the indoor and outdoor study sites was significantly different (p=0.0001).

3.4 PM_{2.5} Concentrations and Pulmonary Function Indices (FEV₁/FVC FEV₁ and FVC)

The Fig. 3 shows the PM_{2.5} exposures among the participants and the percent predicted values of FEV₁/FVC. The participants in the two lung function categories were exposed to significantly different concentrations of PM_{2.5}. (p=0.001). The median of PM_{2.5} concentration among those with normal lung function status was 44.0 µg/m³ while among the participants with impaired lung function the median of PM_{2.5} concentration was 171.0 µg/m³.

Participants with reduced FEV₁ percent predicted were exposed to significantly higher

concentrations of PM_{2.5} (200 µg/m³) in comparison to those with normal FEV₁ percentage predicted for age and height (45 µg/m³) p=0.03. (Fig. 4)

Participants with normal predicted FVC were exposed to lower PM_{2.5} concentrations (49 µg/m³) compared to those with reduced FVC (260 µg/m³) predicted for age and height. There was no significant difference in FVC percent predicted between those exposed to high PM_{2.5} concentrations and those exposed to low PM_{2.5} concentrations p=0.121. (Fig. 5)

3.5 Predictors of Lung Function Status

Two separate logistic regression analyses were conducted to predict PM_{2.5} exposure and cleaning group as predictors of lung function. A test of each model against a constant only model was statistically significant (chi square = 5.018 and 6.429 respectively, p < .05 with df = 1). Prediction success for both PM_{2.5} and cleaning group was 77.8%. The Wald criterion demonstrated that both PM_{2.5} and cleaning group made a significant contribution to prediction (p = .025 and p=.015 respectively). An increase in PM_{2.5} by one unit increases the odds ratio of having impaired lung function by 1.




Fig. 2. Distribution of PM_{2.5} concentrations in indoor and outdoor locations




Fig. 3. Medians of PM_{2.5} concentration (µg/m³) FEV₁/FVC

Table 1. Lung function status of participants

Lung function variable	Street sweepers no (%)	Office cleaners no (%)	p-value
Lung function status (FEV₁/FVC)			
Normal (FEV ₁ /FVC > 70%)	30(42.9)	40(57.1)	0.01*
Impaired (FEV ₁ /FVC < 70%)	15(75.0)	5(25.0)	
FEV₁ percent predicted			
Normal (FEV ₁ %predicted > 80%)	33(45.2)	40(54.8)	0.059
Reduced (FEV ₁ % predicted < 80%)	12(70.6)	5(29.4)	
FVC percent Predicted			
Normal (FVC% predicted > 80%)	36(46.2)	42(53.8)	0.06
Reduced (FVC %predicted < 80%)	9(75.0)	3(25.0)	

^pPearson's Chi-Squared Test (2-sided), *Indicates a p-value at significance level <0.05."

Table 2. Lung function status by key characteristics of participants

	Lung function status		P-value*
	Normal (n=70, FEV ₁ /FVC>70%)	Impaired (n = 20, FEV ₁ /FVC < 70%)	
	No (%)	No (%)	
Use of PPE			
Always	24 (34.3)	8 (40.0)	0.79
Sometimes /never	46 (65.7)	12 (60.0)	
Cooking fuel			
Charcoal	41 (58.6)	14 (70.0)	0.35
Electricity	29 (41.4)	6 (30.0)	
Cleaning location^p			
Indoor -office cleaners	41 (58.6)	4 (20.0)	0.002[*]
Outdoor -street sweepers	29 (41.4)	16 (80.0)	
Allergies^p			
No	37 (52.9)	5 (25.0)	0.02[*]
Yes	33 (47.1)	15 (75.0)	

^pPearson's Chi-Squared Test (2-sided), *Indicates a p-value at significance level <0.05."




Fig. 4. Medians of PM_{2.5} concentration (µg/m³) and predicted values of FEV₁




Fig. 5. Medians of PM_{2.5} concentration (µg/m³) and predicted values of FVC

4. DISCUSSION

4.1 Lung Function Characteristics of Participants

The study confirmed the presence of impaired lung function status among cleaners exposed to fine particulate air pollution (PM_{2.5}) in ambient Lusaka air. The street sweepers showed significantly impaired lung function and higher odds of having impaired lung function than office cleaners. This shows that street sweepers are more likely to have impaired pulmonary function than office cleaners. Indeed other studies have associated street sweeping and exposures to large amounts of dust with respiratory conditions [22, 29].

4.2 Contributing Factors to PM_{2.5} Exposure

The levels of PM_{2.5} depend on several factors. Biomass fuel use is said to be a major driver of respiratory and cardiovascular disease [34]. Most (61.1%) of the participants in the present study used charcoal as a cooking fuel. Although there was no statistically significant difference in Cooking fuel used between the 2 groups, there were more (60%) individuals with impaired lung function who used charcoal compared to those who used electricity ($p > 0.05$). Indeed, there has

been a reported association between biomass (Charcoal and firewood) cooking fuels and impaired lung function [6].

The cleaners recruited to the study had low frequency of PPE use. This could be a possible explanation for the observed lung function impairment in the two groups. Studies show that street sweeping without precautionary measures such as proper use of personal protective equipment in the form of facemasks and respirators may predispose to respiratory conditions [29,34]. This is based on the observation that the use of precautionary measures such as PPE tend to reduce the levels of inhaled particulates and subsequent pathological effects [35].

The current study showed a relationship between presence of allergies and lung function status. This finding was consistent with studies that revealed that allergy symptoms such as those present in asthmatic individuals are worsened in cases of exposure to pollutants and these further indicate a decrease in lung function measurements especially FEV₁ [13,21,25]. In the current study, 75% of those who reported having allergy symptoms had impaired lung function status. The presence of allergy symptoms may be an indicator of increased susceptibility to the effects of PM_{2.5} exposure [25].

Table 3. Predictors of lung function status

Characteristics/factors	Study sample N(%)	Odds ratio (95% CI)	P-value
Particulate matter	90 (100)	0.99 (0.992-0.999)	0.025*
PM _{2.5}			
Cleaning group	45 (50.00)	1.0	
Office cleaners			
Street sweeper	45 (50.00)	0.25 (0.08-0.76)	0.015*

*Indicates a p-value at significance level <0.05"

4.3 PM_{2.5} Concentration ($\mu\text{g}/\text{m}^3$) in the Study Areas

The results in the present study show that real-time PM_{2.5} concentrations were significantly elevated in both indoor and outdoor areas. The measured PM_{2.5} concentrations outdoors were high, ranging from 123 $\mu\text{g}/\text{m}^3$ -398 $\mu\text{g}/\text{m}^3$ compared to 10 $\mu\text{g}/\text{m}^3$ - 53 $\mu\text{g}/\text{m}^3$ indoors. These findings are consistent with other studies carried out in other African cities, which revealed that air pollution levels particularly PM_{2.5} concentrations were quite high and that they exceeded international guidelines [7,8]. In indoor areas air circulation is controlled by the presence of air conditioners hence the small variations in the PM_{2.5} however; this does not prevent increases in PM_{2.5} either [30]. Because of these elevated concentrations, 80% of those with lung function impairment were street sweepers and only 20% were office cleaners.

4.4 PM_{2.5} and Pulmonary Function indices (FEV₁/FVC FEV₁ and FVC)

A significant association was observed between PM_{2.5} concentration and lung impairment (FEV₁/FVC). This finding collaborated with those that reported that interquartile increases in PM_{2.5} exposure results in increased respiratory impairment [21].

The results of this study have shown that the concentration of PM_{2.5} across the lung function status categories were different. Higher concentrations of PM_{2.5} (median 171.00) were associated with impaired lung function status. The level of lung function impairment is related to the dosage of PM an individual is exposed to [9]. The ability for the fine particulate matter (PM_{2.5}) to penetrate the alveoli and cause endothelial damage by release of inflammatory mediators such as chemokines and cytokines causes reduction in lung function [5,10].

In line with other studies [21,25] the current study further revealed that PM_{2.5} concentration had an

effect on the FEV₁ percent predicted and not on the FVC percent predicted. FEV₁ reduces because the inhaled particulates cause irritation in the airways resulting in over production of mucus and proinflammatory mediators that block the airways. The tendency to resist airflow under forced conditions in blocked airways reduced the volume of air that could be forcibly expired as a result FEV₁ is reduced but not the FVC.

5. CONCLUSION

The environments in which both groups of women worked were quite polluted as seen from the rather high real-time PM_{2.5} concentrations which were higher in outdoor environments (streets) compared to indoor environments (offices). Exposure to higher PM_{2.5} concentrations was associated with impaired lung function. This was observed more in those workers who did not consistently use personal protective equipment.

6. LIMITATIONS

This study, being cross sectional, could not determine causality. There were also many other factors which were confounders in this study which needed to be controlled for before conclusions can be made. These require a more detailed study to be done to exclude the influence of factors such as use of wood fuel and charcoal, age and pre-existence of allergies which have significant effect on pulmonary function.

Furthermore, in the current study, only real-time measurements of the levels of PM_{2.5} were done. There are proposals suggesting the use of PM_{2.5} measurements obtained in centrally located stations.

CONSENT

An informed consent form prepared according to the Research Ethics Committee guidelines was given to the participants in order to guarantee

voluntary participation. The contents of the information sheet were translated into the commonly spoken language. Simple language was used in providing the participants with sufficient knowledge to ensure the decision to take part is a well-informed one. Contents such as the purpose of the study, its nature and methods to be used in the study were explained. The information sheet was kept by the participants, while the consent forms were kept by the principal investigator. Participants gave consent either through written or using the thumb prints for those who could not write. The participants were allowed to ask questions pertaining to the study and were free to withdraw from the study at any time if they felt uncomfortable without any penalty or loss.

ETHICAL APPROVAL

Approval to carry out the study was granted in writing by The University of Zambia Biomedical Research Ethics Committee (Assurance No. FWA 00000338, IRB 00001131 of IORG 0000774, Ref: 013-03-14). Permission to conduct the study was obtained from the various employers of the groups of cleaners and the Lusaka City Council.

ACKNOWLEDGEMENTS

This project was supported by grant # 5R24TW008873 administrated by the Fogarty International Center of the National Institutes of Health and funded by OGAC and OAR.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World health organisation air quality guidelines on particulate matter, ozone, nitrogen dioxide and sulphur dioxide, global update 2005, summary of risk assessment; 2005.
Available:<http://www.euro.who.int/Document/E87950.pdf>
(Accessed 10th November 2013)
2. Valavanidis A, Fiotakis K, Vlachogianni T. Airborne particulate matter and human health: Toxicological assessment and importance of size and composition of particles for oxidative damage and carcinogenic mechanisms. *Journal of Environmental Science and Health, Part C: Environmental Carcinogenesis and Ecotoxicology Reviews.* 2008;26(4): 339-362.
DOI: 10.1080/10590500802494538.
3. Lusaka city council and environmental council of Zambia Lusaka city state of the environment outlook report. 2011;75-76.
4. Ghio AJ, Kim C, Devlin RB. Concentrated ambient air particles induce mild pulmonary inflammation in healthy human volunteers. *American Journal of Respiratory Critical Care Medicine.* 2000;162(3pt1):981-8.
5. Daigle C, Chalupa D, Gibb R, Morrow P, Oberdörster G, Utell M, Frampton M. Ultrafine particle deposition in humans during rest and exercise. *Journal of Inhalation Toxicology.* 2003;15:539-52.
6. Umoh VA, Peters E. The relationship between lung function and indoor air pollution among rural women in the Niger Delta region of Nigeria. *Lung India.* 2014;31(2):110-115.
7. Petkova E, Darby J, Kinney P. Particulate matter in African cities air qual atmos health. 2013;6:603-614.
DOI: 10.1007/s11869-013-0199-6.
8. Gree A, Odeshi TA, Sridhar M, Ige M. Outdoor respirable particulate matter and the lung function status of residents of selected communities in Ibadan, Nigeria. DOI: 10.1177/1757913913494152.
Perspectives in Public Health. 2013; 1757913913494152.
9. Ekpenyong E, Etebong O, Akpan E, Samson T, Nyebuk D. Urban city transportation mode and respiratory health effect of air pollution: A cross-sectional study among transit and non-transit workers in Nigeria. *BMJ Open.* 2012;2(5): e001210.1136/bmjopen-2012-001253PMCID: PMC3488752.
10. Nodari S, Corulli A, Manerba A, Metra A, Apostoli P, Cas L. Endothelial Damage Due To Air Pollution Heart International. 2006;2(2):115-125.
11. USA-EPA. Air quality criteria for particulate matter; 2012.
Available:http://www.epa.gov/ttn/naaqs/standards/pm/s_pm_index.html
(Accessed 12th August 2014)
12. Götschi T, Sunyer J, Chann S, De Marco R, Forsberg B, Garcia-Esteban R, et al. Air pollution and lung function in the European community respiratory health survey. *International Journal of Epidemiology;* 2008. PMCID: PMC2734069.

13. Trenga CA, Sullivan JH, Schildcrout JS, Shepherd KP, Shapiro GG, Liu LJ, et al. Effect of particulate air pollution on lung function in adult and paediatric subjects in a seattle panel study. *Chest*. 2006; 29(6):1614-1622.
14. Giradot SP, Smith S, Davis WT, et al. Ozone and PM2.5 exposure and acute pulmonary health effects: A study of hikers in the Great Smoky mountains national park environ health perspect. 2006;114: 1044–1052.
DOI: 10.1289/ehp.8637.
PMCID: PMC1513325.
Available:<http://dx.doi.org>
(Accessed 20th September 2013)
15. Lagorio S, Forastiere F, Pistelli R, et al. Air pollution and lung function among susceptible adult subjects: A panel study *Environmental Health: A Global Access Science Source*. 5:11.
DOI: 10.1186/1476-069X-5-11.
16. Ma"kela" R, Kauppi P, Suuronen K, Tuppurainen M, Hannu T. Occupational asthma in professional cleaning works: A clinical study. *Occupational Medicine*. 2011;61:121–126.
DOI: 10.1093/occmed/kqq192.
Available:<http://occmed.oxfordjournals.org>
(Accessed 13th October 2014)
17. Dysart MM, Galvis BR, Russell AG, et al. Environmental particulate (PM2.5) Augments Stiffness-Induced Alveolar Epithelial Cell Mechanoactivation of Transforming Growth Factor Beta. *PLoS One*. 2014;9(9):e106821.
DOI: [10.1371/journal.pone.0106821](https://doi.org/10.1371/journal.pone.0106821)
PMCID: PMC4167324
(Accessed 20th September 2015)
18. Valavanidis A, Vlachogianni T, Fiotakis K et al. Review pulmonary oxidative stress, Inflammation and Cancer: Respirable particulate matter, fibrous dusts and ozone as major causes of lung carcinogenesis through reactive oxygen species mechanisms. *International Journal of Environmental Research and Public Health*. ISSN 1660-4601.
Available:www.mdpi.com/journal/ijerph
(Accessed 20th September 2015)
19. Matarese A, Santulli G. Angiogenesis in chronic obstructive pulmonary disease: A Translational Appraisal *Transl Med UniSa*. 2012;3:49–56. PMCID: PMC3728789.
Available:<http://www.ncbi.nlm.gov>
(Accessed 20th September 2015)
20. Voekel NF, Douglas IS, Nicolls M, Angiogenesis in Chronic Lung Disease. *Chest*. 2007;131(3):874–879.
DOI: 10.1378/chest.06-2453.
PMCID: PMC4396181.
NIHMSID: NIHMS50137.
Available:<http://www.ncbi.nlm.gov>
(Accessed 20th September 2015)
21. Dales R, Chen AM, Frescura AM, Liu L, Villeneuve PJ. Acute effects of outdoor air pollution on forced expiratory volume in 1 s: A panel study of schoolchildren with asthma. *Eur Respir J*. 2009;34:316–323.
DOI: 10.1183/09031936.00138908.
22. Sabde YD, Zodpey SP. Respiratory morbidity among street sweepers working at Hanumannaga zone of Nagpur Municipal Maharashtra. *Indian Journal of Public Health*. 2008;52:3.
23. Singh LP, Bhardwaj A, Deepak KK. Occupational exposure to respirable suspended particulate matter and lung functions deterioration of steel workers: An exploratory study in India; 2013. Article ID 325410.
Available:<http://dx.doi.org/10.1155/2013/325410>
(Accessed 11th May 2014)
24. Verma S, Sharma Y, Shikha A. Multivariate study of some lung function tests at different age groups in healthy Indian males. *Indian Journal of Chest Disease and Allied Science*. 2002;44: 850–91.
25. Lewis TC, Robins GT, Dvonch TJ, Keeler GJ, Fuyuen Y, Yip FY, et al. Air pollution–associated changes in lung function among asthmatic children in detroit environ health perspect. 2005;113:1068–1075.
DOI: 10.1289/ehp.7533.
Available:<http://dx.doi.org>
(Accessed 14th August 2014)
26. Barreiro J, Perillo L. An approach to interpreting spirometry American family physician. 2013;69(5):11074.
27. Levy M, Quanjer P, Booker K. et al. Diagnostic spirometry in primary care-pro; 2009.
Available:www.thepcrj.com
(Accessed 21st March 2014)
28. Nku C, Peters E, Eshiet A. Effect of exposure to dust on lung function, oxygen saturation and symptoms among female street sweepers in Calabar. *Nigerian*

- Journal of Physiological Sciences. 2005; 24:30–9.
29. Khurshid SA, Mehmood N, Nasim N, Khurshid M, Khurshid B. Sweeper's lung disease: A cross-sectional study of an overlooked illness among sweepers of Pakistan. International Journal of COPD. 2013;8:193-197. Available:<http://dx.doi.org/10.2147/COPD.S40468>"<http://dx.doi.org/10.2147/COPD.S40468> (Accessed 11th May 2014)
30. Zock JP. World at work: Cleaners- multiple occupational hazards in a large service sector Occup Environ Med. 2005;62: 581–584.
31. Muula AS, Rudatsikira E, et al. Occupational illnesses in the 2009 Zambian labour force survey. BMC Res Notes. 2010;3:272.
32. Laima C. Prevalence and correlates of lung function impairment among open-pit miners at Nchanga in Zambia; 2013. Available:<http://hdl.handle.net/12456789/2530>
33. Siziya S. Associations of cement dust with occurrence of respiratory conditions and Lung function. East Africa Journal Public health. 2005;2:1-5.
34. Piddock K, Gordon S, Ngwira A, Msukwa M, Nadeau G, Davis KJ, Nyirenda MJ, et al. A cross-sectional study of household biomass fuel use among a periurban population in Malawi Ann Am Thorac Soc. 2014;11(6):915-2.
35. Nagoda M, Okpapi JU, Babashani M. Assessment of respiratory symptoms and lung function among textile workers at Kano Textile Mills, Kano, Nigeria. Niger J Clin Pract. 2012;15:373-9. Available:<http://www.njcponline.com/text.asp?2012/15/4/373/104505> (Accessed 16th October 2014)

© 2016 Siachingili and Goma; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciedomain.org/review-history/11874>