



Lymphangioma of Nasopharynx in an Adult: A Rare Case

Kartik Parelkar^{1*}, Keya Shah², Smita Nagle¹ and Reshma Hanwate¹

¹Department of ENT, Grant Govt. Medical College and Sir J J Group of Hospitals, Byculla, Mumbai, Maharashtra, India.

²Department of ENT, Seth G.S. Medical College and KEM Hospital, Mumbai, India.

Authors' contributions

This work was carried out in collaboration between all authors. Author KP designed and wrote the first draft of the manuscript. Authors SN and KP were the operating surgeons. Authors KS and RH managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BJMMR/2016/29476

Editor(s):

(1) Gauri Mankekar, ENT Department, PD Hinduja Hospital, Mumbai, India.

Reviewers:

(1) Ramesh Parajuli, Tribhuvan University, Nepal.

(2) Oğuzhan Güven Gümüştas Burtom Imaging Center, Turkey.

(3) Fawzia Butt, University of Nairobi, Kenya.

(4) Shraddha Mukerji, University of Texas Medical Branch, USA.

Complete Peer review History: <http://www.sciencedomain.org/review-history/16762>

Case Report

Received 13th September 2016
Accepted 22nd October 2016
Published 5th November 2016

ABSTRACT

Lymphangiomas are rare benign congenital malformations of the lymphatics, most often encountered during childhood. Generally they are found in the head and neck region, isolated involvement of the nasopharynx is very rare. Our patient had a nasopharyngeal lymphangioma which aggravated his chronic otitis media. The surgical excision of mass was performed by a combined transoral and transnasal endoscopic approach. Histopathological evaluation revealed the diagnosis as lymphangioma. After a follow-up of 1 year the patient is free of recurrence.

Keywords: Lymphangioma; nasopharynx; IHC-immunohistochemistry.

*Corresponding author: E-mail: kartikparelkar@gmail.com;

1. INTRODUCTION

Lymphangiomas are congenital malformations of the lymphatic system. Lymphangiomas are classified as microcystic, macrocystic and cystic hygromas according to the size of the lymphatic cavities incorporated [1].

They commonly present in the head and neck region, with over 90% of lesions manifesting before 2 years of age. The onset of the lesion after 2 years of age is rare [2].

Isolated involvement of the nasopharynx with lymphangioma is very rare. Literature search for nasopharyngeal lymphangioma displayed two unique cases that had been reported in Russia in 1966 and 1969 (no abstract available); after which two more cases have been reported, one in Turkey and the other in India, in 2013 [2,3].

Any nasopharyngeal pathology can precipitate chronic otitis media (COM) and nasopharyngeal lymphangioma is no exception.

2. CASE REPORT

A 34 year old male presented with left chronic otitis media for 10 to 12 years; patient had noticed a small mass in the oral cavity on the left side for last 4 to 5 years.

On otoscopy, the left tympanic membrane showed a large central perforation with

mucopurulent discharge in the middle ear. Patient had intermittent mucopurulent ear discharge which would subside on topical antibiotic medications temporarily.

Oral examination revealed a 3x3 cm pinkish cystic mass in the left supratonsillar fossa. On probing it was soft in consistency and did not bleed on touch. It was attached to the posterior tonsillar pillar and extending up into the nasopharynx (Figs. 1 and 2).



Fig. 1. Cystic mass in the oropharynx



Fig. 2. Zero degree view of the mass in the oropharynx

Diagnostic nasal endoscopy showed a polypoidal mass, which was obliterating the eustachian tube (ET) opening on the left side. It was attached to the supero-lateral wall of the nasopharynx, the posterior lip of the ET opening (Fig. 3) and in continuity with the small cystic mass seen in the oropharynx, through the posterior pillar of the tonsil.



Fig. 3. Zero degree view through the left nostril showing the nasopharyngeal mass (*)

A contrast enhanced CT (CECT) neck reported a minimally enhancing (27 Hounsfield units on plain and 31 Hounsfield units on postcontrast) soft tissue mass measuring 22 x 21 mm. There was minimal obliteration of the left fossa of Rosenmuller and the mass extended into the oropharynx involving the left posterior tonsillar pillar and supratonsillar fossa. (Figs. 4 A and B)

A magnetic resonance imaging (MRI) showed a submucosal soft tissue mass with no neurovascular extension (Fig. 4C).

After routine blood investigations, pre-anesthetic checkup and informed written consent, patient was prepared for surgery. A combined approach of transnasal and transoral routes was used. A simple rubber catheter was inserted through the right nostril and used to retract the soft palate for better visualization. A 4mm rigid zero degree endoscope along with a insulated sickle knife, on which unipolar cautery was applied, was used to gradually separate the nasopharyngeal part. Care was taken to avoid cauterization of the eustachian tube opening. Once the nasopharyngeal part was separated from the lateral wall, the oropharyngeal part was excised in a manner similar to tonsillectomy, by placing the patient in the Rose's position. The oropharyngeal extension was removed along with the left posterior pillar of the tonsil using a bipolar cautery (Fig. 5). Hemostasis was achieved, rest of the post-operative period was uneventful. Histopathology showed, multiple cystically dilated channels lined by a flattened endothelium. Immunohistochemistry (IHC) was positive for lymphatic lineage markers D 2-40 (Fig. 6). The histopathological diagnosis was lymphangioma.

3. DISCUSSION

Lymphangiomas are uncommon, congenital benign malformations of the lymphatic system. They are slowly progressive vascular hamartomas.




Fig. 4. Radiological imaging



Fig. 5. Excisional biopsy specimen

Recently Catalfamo et al. [4] reported nine tongue lymphangiomas in adults. Naidu and Mc Calla [5] reported a comprehensive review of studies from 1828 to 2000 years on lymphatic malformation in adults. They found 91 adult cases of lymphangiomas, located on the neck. Our adult male patient had a lymphangioma in the nasopharynx, which is an unusual site. Lymphangiomas are diagnosed on the basis of history, examination findings and radiological investigations. Lymphangiomas at oral cavity, tongue, and larynx can present with symptoms of dysphagia, dyspnea, foreign body sensation, sore throat and tonsillar mass. When located at the nasopharynx, they may cause nasal obstruction or may aggravate the COM without symptoms of nasal obstructions per se. Though our patient had a visible mass in the oropharynx,

smaller sized nasopharyngeal pathology can go undetected if nasal endoscopy is not performed. Hence from our experience, it is essential that every patient with COM be screened with nasal endoscopy to rule out any nasal/nasopharyngeal pathology.

MRI can demarcate the mass from the surrounding tissues and can give the extent of the lesion. CECT scan can also be helpful in investigating these lesions.

Antrochoanal polyps, nasopharyngeal carcinoma, angiofibroma, cystic lesions, and other benign masses should be included in the differential diagnosis. Lymphangiomas can get infected or can cause hemorrhage and the mass effect associated with rapid growth of the lesion. Also in some patients, lymphocytopenia, a precipitating cause of infection, can be documented [6]. Lymphangiomas can be managed by namely surgical excision; laser debulking, sclerotherapy and corticosteroids [7]. It is suggested that the desire for complete excision must be weighed against the need to keep vital structures intact. If the larynx is involved tracheostomy is almost always required [8].

Though in our case we used cautery, surgical laser is considered particularly suitable because it allows precise ablation and recontouring of the tissue so that the functions can be preserved [9]. Coblation too would be similar to laser or perhaps even better, but there is no literature available currently regarding its use for such a




Fig. 6. IHC with D2-40 marker for lymphangioma

case. Sclerosing agents can be used but they act by causing inflammatory reaction hence, their use can result in damage to adjacent structures. Radiotherapy is not effective in treatment of lymphangiomas [10].

In our case we were able to excise the lesion without causing any damage to surrounding vital anatomical structures using a transoral and transnasal endoscopic approach.

We reported this case because of the unusual site of involvement and unusual age of presentation.

4. CONCLUSION

Although lymphangioma of the nasopharynx is very rare, it can cause ET obstruction and aggravate COM. Complete surgical excision followed by a tympanoplasty (if required) would be the ideal way to go about. Since there is a risk of recurrence, regular follow up would be essential.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

ACKNOWLEDGEMENT

We would like to acknowledge the contribution of our pathologist Dr. Mrudula Sampat, she has done an accurate histopathological reporting for us.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bloom DC, Perkins JA, Manning SC. Management of lymphatic malformations. *Curr Opin Otolaryngol Head Neck Surg.* 2004;12:500–504.
2. Haksever M, Akduman D, Aslan S, Yazla S, Haksever H. Nasopharyngeal lymphangioma in an adult: A rarity. *Laryngoscope.* 2013;123(12):2972-5. DOI: 10.1002/lary.24214 Epub 2013 Jun 11. PubMed PMID: 23712707
3. Verma R, Verma RR, Verma RR, Sardana NK. Isolated lymphangiomatous polyp nasopharynx in an adult first case report in English literature. *Indian J Otolaryngol Head Neck Surg.* 2014;66(4):460-3. DOI: 10.1007/s12070-013-0681-3 Epub 2013 Oct 18. PubMed PMID: 26396962. PubMed Central PMCID: PMC4571466
4. Catalfamo L, Nava C, Lombardo G, Iudicello V, Siniscalchi EN, Saveriade PF. Tongue lymphangioma in adult. *J Craniofac Surg.* 2012;23:1920–1922.
5. Naidu SI, McCalla MR. Lymphatic malformations of the head and neck in adults: A case report and review of the literature. *Ann Otol Rhinol Laryngol.* 2004;113:218–222.
6. Stanescu L, Georgescu EF, Simionescu C, Georgescu I. Lymphangioma of the oral cavity. *Rom J Morphol Em Bryol.* 2006;47:373–377.
7. Orvidas LJ, Kasperbauer JL. Pediatric lymphangiomas of the head and neck. *Ann Otol Rhinol Laryngol.* 2000;109(4):411–421.
8. Cohen SR, Thompson JW. Lymphangiomas of the larynx in infants and children: a survey of pediatric lymphangioma. *Ann Otol Rhinol Laryngol.* 1986;95(suppl 127):1–20.
9. Papsin BC, Evans JNG. Isolated laryngeal lymphangioma: A rare cause of airway obstruction in infants. *J Laryngol Otol.* 1996;110:969–972.
10. Sobol SE, Manoukian JJ. Acute airway obstruction from a laryngeal lymphangioma in a child. *Int J Pediatr Otorhinolaryngol.* 2001;58:255–257.

© 2016 Parelkar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/16762>