

Differential Enthalpy, Factor of Cardiac Power and Precursor of Work Power from the Nodal Tissue

A. B. Kunyima^{1*}, S. N. Lusamba^{1,2}, M. B. Kunyima³ and C. N. Kabele²

¹Department of Chemistry, Laboratory of Physical Organic and Food Chemistry, Faculty of Sciences, University of Kinshasa, P.O.Box 190 Kinshasa XI, Democratic Republic of Congo.

²Department of Chemistry, Laboratory of Analytical Chemistry and Quality Control, Faculty of Sciences, University of Kinshasa, P.O.Box 190 Kinshasa XI, Democratic Republic of Congo.

³Higher Institute of Medical Techniques of Kinshasa, P.O.Box 774 Kinshasa XI, Democratic Republic of Congo.

Authors' contributions

This work was carried out in collaboration between all authors. Author ABK designed the study, wrote the protocol, and wrote the first draft of the manuscript. Author SNL is the experimentalist of the results of this paper and managed the literature searches and author MBK analyzed and discussed the study and author CNK managed the proof reading and correction of manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CA/2016/26708

Editor(s):

- (1) Francesco Pelliccia, Department of Heart and Great Vessels University La Sapienza, Rome, Italy.
(2) Fatih Yalcin, Fulbright Visiting Professor, School of Medicine, Department of Cardiology, Johns Hopkins University, USA.

Reviewers:

- (1) Marina Ostojic, Clinical Center of Serbia, Serbia.
(2) Anonymous, University of British Columbia, Canada.
(3) Safinaz Ebrahim El-Toukhy, National Research center, Egypt.
(4) Salma Abdelmalek, Sousse Faculty of Medicine, Sousse, Tunisia.
(5) Luciana Moisés Camilo, Federal Institute of Education, Science and Technology of Rio de Janeiro, Brazil.
(6) Loc Nguyen, University of Science, Ho Chi Minh, Vietnam.

Complete Peer review History: <http://sciencedomain.org/review-history/15218>

Original Research Article

Received 29th April 2016
Accepted 14th June 2016
Published 30th June 2016

ABSTRACT

Background: The blood can be considered as a real fluid flowing in fixed vibrating myocardial canal. Its ejection during the left ventricular contraction is the phenomenon hereby studied. The left ventricle has been taken as a checking volume encompassing one part of aorta. The limit of this thermodynamic system is dotted and situated under endocardial tissue. The blood in this closed ventricle is considered in stationary state. A vent connects the left ventricle to the aorta and the system receives the work power from nodal tissue.

*Corresponding author: E-mail: anaclet.kunyima@unikin.ac.cd;

Aim and Objective: The study aims to dial differential enthalpy and to show that it can be used to quantify the cardiac power.

Methodology: The study has been performed on healthy persons, men and women, of Democratic Republic of Congo. The sample of 20.000 cases of healthy persons whose age varies between 13-73 years has been investigated. The blood pressure measurements and calculations are our methodology of work. The figures have been plotted by means of origin 8 program.

Results: In each cross section of life the parameter in title has been calculated and correlated at different ages. Also the cardiac power has been calculated at different cardiac frequencies. Interpretations of the results are commented.

Conclusion: A new concept of differential enthalpy is introduced in this paper to show that this original parameter has twofold roll. Firstly it can be used to calculate cardiac power and secondly it can help to evaluate the work power from Keith Flack node. This last assertion constitutes the content of our research on Keith Flack impetus assessment in heart acting.

Keywords: Differential enthalpy; cardiac power; work power; total massic differential enthalpy; differential pressure.

ABBREVIATIONS

Variation of the total differential enthalpy (ΔH_D); Cardiac frequency (f_c); Total internal energy (U_{cz}); Variation of the total massic differential enthalpy (Δh_{czb}); Total massic internal energy (μ_{czb}); Cardiac power (P_C); Differential pressure (P_D).

1. INTRODUCTION

The blood plasma and the cytoplasm are not the simple liquid media. They contain many molecules belonging to all the possible types and which give to the solution a prominent viscosity particularly in the case of the cytoplasm, considered as a gel.

The viscosity is the outcome of the interactions between the solvent molecules and the molecules of dissolved substances. Each movement of a molecule in solution entails friction with the solvent molecules and with the others dissolved molecules. Should the molecule is not spherical, for example should it is a fibrillar protein, the frictions are enhanced and the diffusion is slowed, but also it happens in privileged direction and not in all the directions [1].

The molecules have tendency to diffuse in all the space offered to them and afterwards they move permanently in all directions in the totality of this space with the velocities depending on their size and on the linked water crown they entail. Hence the blood can be considered as a real fluid flowing in fixed vibrating myocarde canal. Its ejection during the left ventricular contraction is the phenomenon hereby studied [1,2,3,4].

The left ventricle has been taken as a checking volume encompassing one part of the aorta. The

limit of this thermodynamic system is dotted and situated under endocard. The blood in this closed ventricle is considered in stationary state. A vent connects the left ventricle to the aorta and the system receives the work power from nodal tissue. When the left ventricle is filled up by the blood during the diastolic period, this blood is not in displacement and it has a total internal energy $U_{czb} = \mu_{czb} dM_b^+$, where dM_b^+ is the elementary mass of blood received by the system and μ_{czb} is the total massic internal energy of the blood [1,4,5,6,7].

When the blood pressure in left ventricle becomes higher, a certain amount of blood is poured in aorta, in this case displacement energy should be added in order to take into account the total energy relative to the mass of blood in displacement in the opened system, that means $\mu_{czb} dM_b^+ + P_D v dM_b^+ = (\mu_{czb} + P_D v) dM_b^+ = h_{czb} dM_b^+$, where P_D is the differential pressure at the origin of the left ejection, v is the massic volume of blood and h_{czb} is the massic total enthalpy of blood in displacement. The total internal energy U_{cz} of the system defined by the border is then $dU_{cz} = \delta E^+ + \delta Q^+ + h_{czb} dM_b^+$. It will be demonstrated in our research on Keith Flack impetus assessment in heart acting that $\delta Q^+ = 0$ and δE^+ is the technical work energy received by the system corresponding to the work Power subtitle to be dialed. It can be uptaken that massic total differential enthalpy is

equal to $h_{czb} \left(\frac{\Delta H_D}{M} = \Delta h_{czb} \right)$ and it will be used to dial work power.

Unfortunately the total work power received by the system can not be calculated because it serves not only to electrical activity of myocarde but also to the blood displacement. Therefore the first part of work power will never be known in recent future perhaps, whereas the last one will be calculated in next paper on the chosen thermodynamic system because the ejected volume of blood is given for healthy persons [1,5,8,9]. In the mean time research is continuing in the laboratory to deal with the investment of work power in myocarde electrical activity.

2. MATERIALS AND METHODS

The sphygmomanometer (Manual, type Aneroid 767 Tycos murale de Welch Allyn) has been used [10,11]. The observation and calculations [12,13] are our methodology of work. The figures have been plotted by means of origin 8 program.

The sample of 20.000 cases of healthy persons (in regular check-up) whose age varies between 13-73 years has been investigated as follow. The measures of blood pressure were taken in seated position, at rest.

Group 1: 10.000 men with such subgroups

- 1500 men with average of 13 years old.
- 1500 men with average of 23 years old.
- 1500 men with average of 33 years old.
- 1500 men with average of 43 years old.
- 1500 men with average of 53 years old.
- 1500 men with average of 63 years old.
- 1000 men with average of 73 years old.

Group 2: 10.000 women with the same subgroups.

The subgroup of 13 years old contains for example the persons with 13 years and one month, 13 years and few days, 13 years and five months, 13 years and six months. The reasoning is the same for the other ages until 73 years. After fastidious calculations and measures during 2.5 years (April 2013-December 2015), it has been observed that the minor error was of the same order of magnitude than the implement error. So it has been decided to take the implement error as precision of all the measures of P_D .

Consequently the blood pressure of a healthy person of 13 years old will be find around the value mentioned in the table for this age. Exception can of course occur. Hence the age mentioned in the table is for group and individual. It should noted that the study has been performed in Kinshasa precisely in Kinshasa Reference General and Provincial Hospital, in University Clinics of Kinshasa and in our Laboratory (LACOPA). The procedure of experimentation has remained the same [12].

The following equation has been used for the blood which is the real fluid [1,8,14] in order to verify the hypothesis in title.

$$\int dH = \int C_p dT + \int \left[V - T \left(\frac{\partial V}{\partial T} \right)_p \right] dT$$

Human body temperature is constant [1,4].

$$\text{So } dT = 0 \text{ and } \left(\frac{\partial V}{\partial T} \right)_p = 0$$

$$\int dH = \int_1^2 V dp$$

The ejected volume at left ventricle contraction is constant (80 ml) [1,4].

$$\Delta H_D = V(P_2 - P_1) = -V(P_1 - P_2) = -V(P_s - P_d)$$

$P_1 = P_s =$ Systolic pressure.

$P_2 = P_d =$ Diastolic pressure.

It has been decided to call this energy total differential enthalpy because it depends on differential pressure ($P_D = P_s - P_d$).

This parameter can be used to calculate cardiac Power ($P_c = -\Delta H_D \cdot f_c$), where f_c is the cardiac frequency and the minus sign means that the system has given this power. In the continuation it will be relinquished specially in the figures. The same reasoning is proposed for ΔH_D .

Converted in total massic differential enthalpy (h_{czb}) it becomes the energy accompanying the blood movement through the virtual border between left ventricle and aorta.

$$\text{Indeed } \Delta H_D = + \frac{M}{\rho} (P_d - P_s)$$

$$h_{czb} = \frac{\Delta H_D}{M} = + \frac{1}{\rho} (P_d - P_s)$$

M is the blood mass and ρ is the blood volumic mass.

Δh_{czb} is the variation of the total massic differential enthalpy ($h_{czb} = h + \frac{c^2}{2} + gz$).

When a thermodynamic system is a seat of work transfers, heat transfers and mass transfers, a distinction should be made between global work energy received by the system A^+ and technical work energy E^+ received only at the mobile parts of the system [8].

$$\delta A^+ = \delta E^+ + \delta E_b^+$$

δE^+ is the technical energy at mobile parts.

δE_b^+ is the energy at the blood.

The total internal energy U_{cz} of the system can be written: $dU_{cz} = \delta A^+ + \delta Q^+$, where $U_{cz} = U + \frac{Mc^2}{2} + Mgz$; $u_{cz} = u + \frac{c^2}{2} + gz$

$$dU_{cz} = \delta E^+ + \delta Q^+ + \delta E_b^+$$

where $\delta E_b^+ = \delta U_{czb} = u_{czb} \cdot dM_b^+$ (for stationary blood).

$\delta E_b^+ = F_b C_b dt = P_D S_b C_b dt$ (for blood in movement), where C_b is the mean velocity of the blood in the section S_b ; P_D is blood pressure through the S surface = P_D .

For the stationary blood:

$$dU_{cz} = \delta E^+ + \delta Q^+ + u_{czb} \cdot dM_b^+$$

For the blood in displacement $\dot{M}_b^+ = \frac{dM_b^+}{dt} = \frac{C_b \cdot S_b}{v_b}$, where v_b is massic volume of blood through the S_b surface: $v_b \frac{dM_b^+}{dt} = S_b C_b$.

This last relation in the expression of the blood in movement gives the contribution of the blood in displacement to its energy:

$$\delta E_b^+ = P_D v_b \frac{dM_b^+}{dt} dt = P_D v_b dM_b^+$$

Globally it can be written

$$\delta E_b^+ = u_{czb} \cdot dM_b^+ + P_D v_b dM_b^+ = (u_{czb} + P_D v_b) dM_b^+$$

$$\delta E_b^+ = h_{czb} dM_b^+$$

$$dU_{cz} = \delta E^+ + \delta Q^+ + h_{czb} \cdot dM_b^+$$

In the calculation the kinetic energy and the potential energy will be neglected.

3. RESULTS AND DISCUSSION

Epidemiologic studies have shown clearly that the cardiac frequency is a risk factor of morbidity and cardiovascular death; more for men than for women [15].

The Impact of the cardiac frequency remains indeed meaningful after adjusting on the age and many factors of cardiovascular risk such as body mass index, the glycaemia, the cholesterol, the triglycerides and arterial pressure. The effect of high cardiac frequency has been found as well in the hypertensive group as in the healthy persons even though in the case of the hypertensive the impact is high. It has been observed that for the women the effect of cardiac frequency is weak [16].

It is generally admitted that the cardiac frequency decreases with the age. Indeed the most observed cardiovascular manifestation for the old persons is the decrease of maximum cardiac frequency with the effort.

On the contrary, at the rest, the decrease of the cardiac frequency with the age is less clean. Study has been led on a large fraction of French population, the decrease of the cardiac frequency with the age has not been observed as it is shown in Fig. 1 where it can be seen that on a very large fraction of population, the cardiac frequency is almost constant between 15 and 75 years [17].




Fig. 1. Cardiac frequency (f_c) as a function of gender and age

Moreover the women have a cardiac frequency more of almost 4 bpm than man cardiac frequency.

A difference of cardiac frequency between genders on this French population has been reported. It has been pointed out that the women cardiac frequencies are higher than the men cardiac frequencies of about 5-7 bpm [17,18].

In this paper the study of cardiac frequency between genders has not been performed. It has been however observed that in each cross section of live all the cardiac frequencies have been found between 60-100 bpm.

This scattering of cardiac frequencies in our population of Democratic Republic of Congo can be explained in short by the lack of informations on nutrition science in general and the absence of the sporting activities.

The insufficiency of better medical taking charge of population and its growing impoverishment are also at the origin of the situation [13,19,20].

Any way in this work it has been decided to make tables and figures with systolic and diastolic pressures measured at different cardiac frequencies found in each gender and each cross section of life.

It is useful before continuing to remember the representation of the heart acting in the plane as it can be seen in Fig. 2 (this is a modelisation). This figure is made in Laboratory of Physical Organic and Food Chemistry (LACOPA), Faculty of Sciences, University of Kinshasa, Democratic Republic of Congo. It should be branded on the memory that this figure is done in the teaching purpose.

The Tables 1 and 2 are relative to healthy men and healthy women of 13 years old until 73 years old with a discard of ten years (one generation).

Table 1. Vital parameters for men

Age (years)	P_s (mm Hg)	P_d (mm Hg)	$P_D \pm \Delta P_D$ (mm Hg)	$\Delta H_D \pm \Delta \Delta H_D$ (J)	fc (bpm)	$P_c \pm \Delta P_c$ (W)
13	118	68	$50 \pm 0,5$	$- 0,526 \pm 0,005$	60	$-0,526 \pm 0,005$
					65	$-0,570 \pm 0,005$
					70	$-0,614 \pm 0,006$
					75	$-0,657 \pm 0,006$
					80	$-0,701 \pm 0,007$
					85	$-0,745 \pm 0,007$
					90	$-0,789 \pm 0,008$
					95	$-0,833 \pm 0,008$
					100	$-0,877 \pm 0,008$
					23	123
65	$-0,547 \pm 0,005$					
70	$-0,589 \pm 0,006$					
75	$-0,631 \pm 0,006$					
80	$-0,673 \pm 0,007$					
85	$-0,715 \pm 0,007$					
90	$-0,757 \pm 0,008$					
95	$-0,800 \pm 0,008$					
100	$-0,842 \pm 0,008$					
33	120	73	$47 \pm 0,5$	$-0,495 \pm 0,005$		
					65	$-0,536 \pm 0,005$
					70	$-0,577 \pm 0,006$
					75	$-0,618 \pm 0,006$
					80	$-0,660 \pm 0,007$
					85	$-0,701 \pm 0,007$
					90	$-0,742 \pm 0,008$
95	$-0,783 \pm 0,008$					

					100	-0,825 ± 0,008
					60	-0,495 ± 0,005
					65	-0,536 ± 0,005
					70	-0,577 ± 0,006
43	122	75	47 ± 0,5	-0,495 ± 0,005	75	-0,618 ± 0,006
					80	-0,660 ± 0,007
					85	-0,701 ± 0,007
					90	-0,742 ± 0,008
					95	-0,783 ± 0,008
					100	-0,825 ± 0,008
					60	-0,526 ± 0,005
					65	-0,570 ± 0,005
					70	-0,614 ± 0,006
53	132	82	50 ± 0,5	-0,526±0,005	75	-0,657 ± 0,006
					80	-0,701 ± 0,007
					85	-0,745 ± 0,007
					90	-0,789 ± 0,008
					95	-0,833 ± 0,008
					100	-0,877 ± 0,008
					60	-0,557 ± 0,005
					65	-0,604 ± 0,005
					70	-0,650 ± 0,006
63	140	87	53 ± 0,5	-0,557±0,005	75	-0,697 ± 0,006
					80	-0,743 ± 0,007
					85	-0,790 ± 0,007
					90	-0,836 ± 0,008
					95	-0,883 ± 0,008
					100	-0,929 ± 0,008
					60	-0,642 ± 0,005
					65	-0,695 ± 0,005
					70	-0,749 ± 0,006
73	142	81	61 ± 0,5	-0,642±0,005	75	-0,802 ± 0,006
					80	-0,856 ± 0,007
					85	-0,909 ± 0,007
					90	-0,963 ± 0,008
					95	-1,016 ± 0,008
					100	-1,070 ± 0,009

It can be seen in this table that at a given ΔH_D the cardiac power increases linearly with increasing cardiac frequency as it is shown in Fig. 2 [13,14].

When the cardiac power is followed as a function of age at a given cardiac frequency, the figures exhibit all of them the decrease of the cardiac power from 13-43 years; stagnation between 43-53 years and afterwards an increase until 73 years (see for example Fig. 4).

The same pace is followed by the differential pressure whose the relation with cardiac power has been established [13].

Note that the heart acting is better should the given cardiac power and its yield are low.

ΔH_D also decreases with the encreasing age, becomes stagnant between 33-43 years and afterwards increases as it is observed in Fig. 5a.

When cardiac power at a given cardiac frequency in each cross section of life is correlated at differential enthalpy through the ages a straight line is observed as it can be seen in Figs. 5b and 5c for example. Note that the slopes of these figures determine exactly the frequencies in question to show the efficiency and the precision of measures.




Fig. 2. Schematic representation of the cardiac systems in a plane. Made in LACOPA/UNIKIN, DRC




Fig. 3. Cardiac power (P_c) versus cardiac frequency (f_c)




Fig. 4. Cardiac power (P_c) versus Age




Fig. 5a. Variation of the total differential enthalpy(ΔH_D) versus Age




Fig. 5b. Variation of cardiac power (P_c) at $f_c = 60$ bpm as a function of the total differential enthalpy (ΔH_D)




Fig. 5c. Variation of cardiac power (P_C) at $f_C = 100$ bpm as a function of the total differential enthalpy (ΔH_D)

The Table 2 shows the differential enthalpy values and the cardiac power at different cardiac frequencies for the women at the same ages (13-73).

When one has a look to this table the observation is the same.

The cardiac power is correlated to cardiac frequency as it is shown in Fig. 6.

It can be seen the difference with Fig. 3 in the behavior of age women.

The cardiac power as a function of age at a given cardiac frequency shows the following pace.

When this is compared with the man the behavior is quite different.

When ΔH_D is followed as a function of age, it is observed in this group the curve quite different compared to men as it can be remarked in Fig. 8a.

Table 2. Vital parameters for women

Age (years)	P_{sys} (mm Hg)	P_{dias} (mm Hg)	$P_D \pm \Delta P_D$ (mm Hg)	$\Delta H_D \pm \Delta \Delta H_D$ (J)	fc (bpm)	$P_c \pm \Delta P_c$ (W)
13	117	71	$46 \pm 0,5$	$-0,484 \pm 0,005$	60	$-0,484 \pm 0,005$
					65	$-0,524 \pm 0,006$
					70	$-0,564 \pm 0,006$
					75	$-0,605 \pm 0,006$
					80-	$-0,645 \pm 0,007$
					85	$-0,685 \pm 0,007$
					90	$-0,726 \pm 0,008$
					95	$-0,766 \pm 0,008$
					100	$-0,807 \pm 0,009$
					60	$-0,473 \pm 0,005$
					65	$-0,513 \pm 0,006$
					70	$-0,552 \pm 0,006$

23	118	73	45 ± 0,5	-0,473±0,005	75	-0,592 ± 0,006
					80	-0,631 ± 0,007
					85	-0,671 ± 0,007
					90	-0,710 ± 0,008
					95	-0,750 ± 0,008
					100	-0,789 ± 0,009
33	118	72	46 ± 0,5	-0,484 ± 0,005	60	-0,484 ± 0,005
					65	-0,524 ± 0,006
					70	-0,564 ± 0,006
					75	-0,605 ± 0,006
					80	-0,645 ± 0,007
					85	-0,685 ± 0,007
43	131	81	50 ± 0,5	-0,526 ± 0,005	90	-0,726 ± 0,008
					95	-0,766 ± 0,008
					100	-0,807 ± 0,009
					60	-0,526 ± 0,005
					65	-0,570 ± 0,005
					70	-0,614 ± 0,006
53	135	83	52 ± 0,5	-0,547 ± 0,005	75	-0,657 ± 0,006
					80	-0,701 ± 0,007
					85	-0,745 ± 0,007
					90	-0,789 ± 0,008
					95	-0,833 ± 0,008
					100	-0,877 ± 0,008
63	138	87	51 ± 0,5	-0,547 ± 0,005	60	-0,547 ± 0,005
					65	-0,592 ± 0,006
					70	-0,638 ± 0,006
					75	-0,684 ± 0,006
					80	-0,729 ± 0,007
					85	-0,775 ± 0,007
73	140	89	51 ± 0,5	-0,536 ± 0,005	90	-0,821 ± 0,008
					95	-0,866 ± 0,008
					100	-0,912 ± 0,009
					60	-0,536 ± 0,005
					65	-0,581 ± 0,006
					70	-0,626 ± 0,006
73	140	89	51 ± 0,5	-0,536 ± 0,005	75	-0,671 ± 0,006
					80	-0,715 ± 0,007
					85	-0,760 ± 0,007
					90	-0,805 ± 0,008
					95	-0,850 ± 0,008
					100	-0,894 ± 0,009




Fig. 6. Cardiac power (P_c) versus cardiac frequency (f_c)




Fig. 7. Cardiac power (P_c) versus Age

There is physiological difference between men and women. P_D for men is higher than for women except at the ages 43 of and 53 years where the inversion is observed. Comment is the same with ΔH_D .

Also in this case when cardiac power at a given cardiac frequency in each cross section of life is correlated at differential enthalpy through the ages a straight line is observed (Figs. 8b and 8c), straight line whose slope determines exactly the frequency in question.




Fig. 8a. Variation of the total differential enthalpy(ΔH_D) versus Age




Fig. 8b. Variation of cardiac power (P_C) at $f_c = 60$ bpm as a function of the total differential enthalpy (ΔH_D)




Fig. 8c. Variation of cardiac power (P_C) at $f_C= 100$ bpm as a function of the total differential enthalpy (ΔH_D)

4. CONCLUSION

The overall purpose is to present the new concept of differential pressure allowing to calculate cardiac power and through which the work power from Keith-Flack node will be deemed [1,8].

ΔH_D does not depend on cardiac frequency, it depends on age. For men the value of ΔH_D at 13 years (0.526 J) after decreasing with age is reached again at 53years (0.526 J) and increases until 73 years. For women however the value of ΔH_D at 13 years (0.484 J) after decreasing with age is reached earlier at 33 years (0.484 J) and increases until 63 years where it remains constant until 73 years showing thus men and women are really physiologically quite different [17].

Also men ΔH_D values are higher than women ΔH_D values except at 43 and 53 years old where inversion is observed meaning that at a certain moment of life the inversion can occur between two genders. The observation is the same with respect to the cardiac power at given cardiac frequency.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kunyima A. et Kunyima M. Biophysique cardiaque: Traitement thermoexergetique du système cardiaque. Edition Académique Express Press; 2007.
2. Jawad Mazhar, Mary Mashicharan, Ahmad Farshid. Predictors and outcome of no-reflow post primary percutaneous coronary intervention for ST elevation myocardial infarction. IJC Heart & Vasculature. Elsevier; 2015.
3. Longa Kaluba, Bellington Vwalika, Methuselah Jere, Fastone M. Goma. Carotid-radial pulse wave form and velocity in normotensive and hypertensive Pregnant Women at University Teaching Hospital (UTH), Lusaka, Zambia, Cardiology and Angiology: An International Journal. 2015;4(3):107-114.
4. Silbernagl S. et Despopoulos A. Atlas de poche de physiologie. 3rd edition, Médecine- Sciences, Flammarion; 2002.

5. Bejan A. Advanced engineering thermodynamics. éd. J. Wiley, New York; 1997.
6. Aurora Bakalli, Ljubicab Georgievska-Ismail, Bedri Zahiti, Dardan Koçinaj, Agreta Gecaj-Gashi, Ardiana Murtezani. Relationship of mitral annular plane systolic excursion (MAPSE) to left ventricular thrombus formation. *Cardiology and Angiology: An International Journal*. 2014;2(4):238-246.
7. Ahlam Kadhim Al Hamdany. Changes of pre ejection period and left ventricular ejection time during head up tilt. *Cardiology and Angiology: An International Journal*. 2015;3(1):12-16.
8. Lucien Borel et Daniel Favrat. Thermodynamique energetique : De l'energie a l'exergie. 2nd edition, Presses Polytechniques et Universitaires Romandes. Lausanne. 2011;1.
9. Jimmy T. Efirid, Wesley T. O'Neal, Stephen W. Davies, Whitney L. Kennedy, Lada N. Alger, Jason B. O'Neal T. Bruce Ferguson, Alan P. Kypson. Long-term mortality of 306,868 patients with multi-vessel coronary artery disease: CABG versus PCI. *British Journal of Medicine & Medical Research*. 2013;3(4):1248-1257.
10. Gallavardin. La tension artérielle en clinique, sa nature, sa valeur sémiologique. 2nd edition, 1 vol. in-8-, 200 figures, Masson et Cie, 1920; Sciences.gloubik.info, The date of putting on line: Saturday 8th February; 2014.
11. Simon Vézina. Applications de l'hydrodynamique. Physique XXI Tome A. Mécanique. 1st edition. Collège de Maisonneuve. Québec; 2015.
12. Kunyima AB, Lusamba SN, Malumba AM, Kabele CN. About physical bases of heart acting. Case of Healthy Women in Democratic Republic of Congo. *Cardiology and Angiology: An International Journal*. 2016;5(1):1-14.
13. Lusamba S. Ntumba. Bases physiques du fonctionnement cardiaque: Application de "Kunyima equation" aux femmes saines de la République Démocratique du Congo. DEA (Master). UNIKIN; 2016.
14. Jean-Noël Foussard et Edmond Julien. Thermodynamique, fondements et applications - Thermodynamique, bases et applications. éd. Dunod, Paris; 2005.
15. Palatini P, Benetos A, Julius S. Impact of increased heart rate on clinical outcomes in hypertension: Implications for antihypertensive drug therapy. *Drugs*. 2006;66:133-44.
16. Benetos A, Thomas F, Bean K, et al. Resting heart rate in older people: A predictor of survival to age 85. *J. Am Geriatricsoc*. 2003;51:284-5.
17. Morcet J, Safar M, Thomas F, Guize L, Benetos A. Associations between heart rate and other risk factors in a large French population. *J Hypertens*. 1999;17:1671-6.
18. Palatini P, Julius S. Heart rate and the cardiovascular risk. *J Hypertens*. 1997; 15:3-17.
19. Pamplona Roger. Guide des plantes médicinales, Tome 1, éd. Vie et Santé; 2000.
20. Kabele Ngiefu Christophe et Lusamba S. Ntumba. Contraintes à l'innovation et opportunités de relance de l'industrie alimentaire à base des produits locaux. Colloque International sur la Biodiversité et changements globaux: Valorisation des effluents des industries, des résidus agropastoraux et forestiers. Université de Ngaoundéré. Cameroun. From 21 to 23 July; 2015.

© 2016 Kunyima et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/15218>